4/6

C. Hall

C. Hall.

A TREATISE

ON

ELEMENTARY TRIGONOMETRY.

A TREATISE

ON

ELEMENTARY TRIGONOMETRY

BY THE

REV. J. B. LOCK, M.A.

FELLOW OF GONVILLE AND CAIUS COLLEGE, CAMBRIDGE;
FORMERLY MASTER AT ETON.

STEREOTYPED EDITION

London:
MACMILLAN AND CO.
AND NEW YORK.
1891

[*The Right of Translation is reserved.*]

Printed 1882.
New edition 1884, *new edition* 1885, *new edition* 1886,
new edition 1887, *new edition* 1888, *reprinted* 1889, 1890
1891.

FROM THE PREFACE TO THE FIRST EDITION.

THE present Work on ELEMENTARY TRIGONOMETRY contains that part of the subject which can conveniently be explained without the use of infinite series. It is intended either for class-teaching or for private study. Accordingly the Examples are numerous and for the most part easy. Those which are not original have been selected from the Cambridge and Army Examination Papers of the last few years.

The Miscellaneous Examples are somewhat more difficult, and should in most cases be postponed until the student reads the subject for the second time.

The order in which the chapters are read may be varied at the discretion of the Teacher.

* * * *

J. B. L.

ETON,
March, 1882.

IN the SECOND EDITION a short course has been indicated for the use of Students who wish to read the Solution of Triangles as early as possible. Such

Students are advised to omit every article that is marked with an asterisk.

A double asterisk has been placed before those articles which should be omitted by all Students until they are reading the subject for the second time.

A Key by Mr CARR is now ready.

CONTENTS

CHAP.		PAGE
I.	On Measurement	1
II.	On Incommensurable Quantities	9
III.	On the Relation between the Circumference of a Circle and its Diameter	14
IV.	On the Measurement of Angles	23
V.	The Trigonometrical Ratios	44
VI.	On the Ratios of Certain Angles	53
VII.	On the Trigonometrical Ratios of the same Angle	66
VIII.	On the Use of the Signs + and −	86
IX.	On the Use of + and − in Trigonometry	90
X.	On Angles unlimited in Magnitude. I.	101
XI.	On the Ratios of Two Angles	117
XII.	On the Ratios of Multiple Angles	133
XIII.	On Angles unlimited in Magnitude. II.	144
XIV.	On Logarithms	161
XV.	On the Use of Mathematical Tables	177
XVI.	On the Relations between the Sides and Angles of a Triangle	191
XVII.	On the Solution of Triangles	206
XVIII.	On the Measurement of Heights and Distances	221
XIX.	On Triangles and Circles	230
XX.	On the Area of the Circle, the Construction of Trigonometrical Tables, etc.	241
	Examples for Exercise	253
	Examination Papers	275
	Answers to Examples	291

CHAPTER I.

On Measurement.

1. It is usual to say that we have measured any concrete quantity, when we have found out *how many* times it contains some familiar quantity of the same kind.

We say for example, that we have measured a line, when we have found out *how many* feet it contains. We say that we have measured a field, when we have found out *how many* acres or how many square yards it contains.

2. To know the measurement of any quantity then, we must have two things. First, we must have a *unit*, or standard of reference, of the *same kind* as the thing measured. Secondly, we must have the *measure*, or the *number of times* the thing measured contains the unit, or standard quantity.

3. Hence, the **measure** of a quantity is the **number,** and the **unit** is the **concrete quantity,** by means of which it is measured.

Example 1. A line contains 261 feet. Here the *measure* or number is 261 and the *unit* a foot.

Example 2. What is the measure of $2\frac{1}{2}$ miles when a yard is the unit?

$$2\tfrac{1}{2} \text{ miles} = \tfrac{5}{2} \times 1760 \text{ yards},$$
$$= 4400 \text{ yards} = 4400 \times 1 \text{ yard},$$

therefore the measure is 4400 when a yard is the unit.

Example 3. What is the unit when the measure of a field of 10 acres is 242?

$$10 \text{ acres} = 242 \times \frac{10 \text{ acres}}{242},$$

$$\therefore \text{ the unit is } \frac{10 \text{ acres}}{242} = 200 \text{ square yards.}$$

Example 4. If the unit be a yards, what is the measure of b miles?

$$b \text{ miles} = b \times 1760 \text{ yards},$$

$$\therefore b \text{ miles} = \frac{b \times 1760}{a} \times a \text{ yards},$$

$$\therefore \text{ the measure required is } \frac{b \times 1760}{a}.$$

EXAMPLES. I.

(1) What is the measure of 1 mile when a chain of 66 feet is the unit?

(2) What is the measure of an acre when a square whose side is 22 yards is the unit?

(3) What is the measure of a ton when a weight of 10 stone is the unit?

(4) The length of an Atlantic cable is 2300 miles and the length of the cable from England to France is 21 miles. Express the length of the first in terms of the second as unit.

(5) The measure of a certain field is 22 and the unit 1100 square yards: express the area of the field in acres.

(6) Find the measure of a miles when b yards is the unit.

(7) The measure of a certain distance is a when the unit is c feet. Express the distance in yards.

(8) A certain sum of money has for its measures 24, 240, 960 when three different coins are units respectively. If the first coin is half a sovereign, what are the others?

* 4. The measure of a quantity is the number of times which that quantity contains the unit.

We may express the same thing in different ways,

(i) The *measure* of a quantity is the **ratio** of that quantity to the unit.

(ii) The *measure* of a quantity is the **fraction** that the quantity is of the unit.

* 5. This last statement is in the language of Arithmetic, and the word 'fraction' is to include whole numbers, and improper fractions.

* 6. The following are therefore different forms of the same question :

(i) What is the measure of 4 miles when 66 feet is the unit?

(ii) How many times does 4 miles contain 66 feet?

(iii) What is the ratio of 4 miles to 66 feet?

(iv) What fraction of 66 feet is 4 miles?

Example (i) What is the measure of a yards when b feet is the unit?

$$a \text{ yards} = 3a \text{ feet} = \frac{3a}{b} \times b \text{ feet},$$

$$\therefore \text{ the measure is } \frac{3a}{b}.$$

(ii) How many times does a yards contain b feet?

$$\text{As in (i), } a \text{ yards} = \frac{3a}{b} \times b \text{ feet.}$$

Answer. $\frac{3a}{b}$ times.

(iii) What is the ratio of a yards to b feet?

$$\text{As in (i), } a \text{ yards} = \frac{3a}{b} \times b \text{ feet,}$$

$$\therefore \frac{a \text{ yards}}{b \text{ feet}} = \frac{3a}{b},$$

$$\therefore \text{ the required ratio is } \frac{3a}{b}.$$

(iv) What fraction is a yards of b feet?

As in (iii), $\frac{a \text{ yards}}{b \text{ feet}} = \frac{3a}{b}$,

$\therefore$ the required fraction is $\frac{3a}{b}$.

* EXAMPLES. II.

(1) The ratio of the area of one field to that of another is 20 : 1, and the area of the first is half a square mile. Find the number of square yards in the second.

(2) The ratio of the heights of two persons is 9 : 8, and the height of the second is 5 ft. 4 in. What is the height of the first?

(3) The measure of a field with 3 acres for unit is 65. Find the ratio of the field to an acre.

(4) One field contains a second of $2\frac{1}{2}$ acres, $6\frac{3}{4}$ times.

What is the measure of the first field in terms of the second?

What is the ratio of the first field to the second?

Express the first as the fraction of the second.

(5) A certain weight is 3·125 of a ton.

What is its measure in terms of 4 cwt.?

How many times does it contain 4 cwt.?

What is its ratio to 4 cwt.?

What fraction is it of 4 cwt.?

(6) The ratio of a certain sum of money to 3 guineas is $\frac{2}{7}$.

Find its measure in terms of one pound.

Find the unit when its measure is 22.

(7) What is the measure of a miles when b chains is the unit?

How many times do a miles contain c chains?

What is the ratio of a miles to d chains?

What fraction is a miles of k chains?

(8) What is the unit when the measure of £20 is $\frac{400}{21}$?

£25 contains a certain sum $\frac{500}{21}$ times. What is that sum?

The ratio of £30 to a certain sum is $\frac{600}{21}$. What is that sum?

The fraction which £10 is of a certain sum is $\frac{200}{21}$. What is that sum?

7. It is explained in Arithmetic, in the application of square measure, that the measure of the area of a rectangle is found in terms of a square unit, by multiplying together the measures of the sides in terms of the corresponding linear unit.

Example. Find in square feet the measure of a square surface whose side is 12 feet.

The area is 12×12 square feet $= 144 \times 1$ square foot,

$\therefore$ the measure required is 144.

8. We shall apply this result to Euclid I. 47.

Example 1. The sides containing the right angle of a right-angled triangle are 3 ft. and 4 ft. respectively; find the length of the hypotenuse.

Let x be the number of feet in the hypotenuse.

Then by Euclid I. 47, the square described on the side of x feet = the sum of the squares described on the sides of 3 feet and 4 feet respectively,



$$\therefore x^2 \text{ square feet} = 9 \text{ square feet} + 16 \text{ square feet}$$
$$= 25 \text{ square feet},$$
$$\therefore x^2 = 25,$$
$$\therefore x = 5.$$

Therefore the length of the hypotenuse is 5 feet.

Example 2. Find the length of the perpendicular drawn from the vertex to the base of an isosceles triangle whose equal sides are 10 feet each, and whose base is 12 feet.

Let ABC be the isosceles triangle such that AB is 10 feet, AC is 10 feet and BC is 12 feet.



Draw AD perpendicular to BC.

Then because the triangle ABC is isosceles AD will bisect the base BC in D; therefore BD is 6 feet.

Let AD contain x feet.

Then by Euclid I. 47, the square on AB = the sum of the squares on BD and AD.

$$\therefore 10^2 \text{ sq. ft.} = 6^2 \text{ sq. ft.} + x^2 \text{ sq. ft.}$$
$$\therefore 10^2 = 6^2 + x^2,$$
$$\therefore x^2 = 100 - 36,$$
$$\therefore x^2 = 64,$$
$$\therefore x = 8.$$

Therefore the required length of AD is 8 feet.

Example 3. Find the length of the diameter of the square one of whose sides contains a feet.



Let $ABCD$ be the square, so that AB is a feet, and AD is a feet.

Let the diameter BD be x feet.

Then the square on DB = the sum of the squares on DA and AB.

$$\therefore\ x^2 \text{ sq. ft.} = a^2 \text{ sq. ft.} + a^2 \text{ sq. ft.}$$

$$\therefore\ x^2 = a^2 + a^2,$$

$$\therefore\ x^2 = 2a^2,$$

$$\therefore\ x = \sqrt{2}\,.\,a.$$

Therefore the required length of the diameter is $\sqrt{2}\,.\,a$ feet.

EXAMPLES. III.

(1) Find the length of the hypotenuse of a right-angled triangle whose sides are 6 feet and 8 feet respectively.

(2) The hypotenuse of a right-angled triangle is 100 yards and one side is 60 yards : find the length of the other side.

(3) One end of a rope 52 feet long is tied to the top of a pole 48 feet high and the other end is fastened to a peg in the ground. If the pole be vertical and the rope tight, find how far the peg is from the foot of the pole.

(4) The houses in a certain street are 40 feet high and the street 30 feet wide : find the length of the ladder which will reach from the top of one of the houses to the opposite side of the street.

(5) A wall 72 feet high is built at one edge of a moat 54 feet wide; how long must scaling ladders be to reach from the other edge of the moat to the top of the wall?

(6) A field is a quarter of a mile long and three-sixteenths of a mile wide: how many cubic yards of gravel would be required to make a path 2 feet wide to join two opposite corners, the depth of the gravel being 2 inches?

(7) The sides of a rectangular field are $4a$ feet and $3a$ feet respectively. Find the length of its diameter.

(8) If the sides of an isosceles triangle be each $13a$ yards and the base $10a$ yards, what is the length of the perpendicular drawn from the vertex to the base?

(9) Show that the perpendicular drawn from the right angle to the hypotenuse in an isosceles right-angled triangle, each of whose equal sides contains a feet, is $\frac{\sqrt{2}}{2} . a$ ft.

(10) If the hypotenuse of a right-angled isosceles triangle be a yards, what is the length of each side?

(11) Show that the perpendicular drawn from an angular point to the opposite side of an equilateral triangle, each of whose sides contains a feet, is $\frac{\sqrt{3}}{2} . a$ ft.

(12) If in an equilateral triangle the length of the perpendicular drawn from an angular point to the opposite side be a feet, what is the length of the side of the triangle?

(13) Find the ratio of the side of a square inscribed in a circle to the diameter of the circle.

(14) Find the distance from the centre of a circle of radius 10 feet, of a chord whose length is 8 feet.

(15) Find the length of a chord of a circle of radius a yards, which is distant b feet from the centre.

(16) The three sides of a right-angled triangle, whose hypotenuse contains $5a$ feet, are in arithmetical progression: prove that the other two sides contain $4a$ feet and $3a$ feet respectively.

* CHAPTER II.

On Incommensurable Quantities.

9. Two numbers are said to be **commensurable** when their ratio can be expressed as an *arithmetical* fraction: that is, as a fraction whose numerator and denominator are both whole numbers.

Example. $4{\cdot}9\dot{3}$ and $81\frac{3}{4}$ are two commensurable numbers.

Their ratio is $4\frac{84}{90} \div 81\frac{3}{4} = \frac{444 \times 4}{90 \times 327}$.

10. Two numbers are said to be **incommensurable** when their ratio *cannot* be expressed as an arithmetical fraction.

Example. $\sqrt{2}$ and 1 are two incommensurable numbers. For $\frac{\sqrt{2}}{1}$ cannot be expressed exactly as an arithmetical quantity. So $\sqrt{3}$ and $\sqrt{2}$ are two incommensurable numbers.

11. One number alone is said to be *an incommensurable number* when it is incommensurable with *unity*. So that an incommensurable number cannot be expressed as an arithmetical fraction.

Example. $\sqrt{2}$ and $\sqrt{3}$, and all surd numbers, are incommensurable numbers.

12. Two *quantities* are said to be commensurable when their *measures* referred to a common unit are commensurable.

Example. A mile and a thousand yards are two commensurable quantities. Their measures with a yard for unit are 1760 and 1000; and these are commensurable numbers.

13. Two quantities are said to be incommensurable when their measures referred to a common unit are incommensurable.

Example. The side of a square and its diameter are two incommensurable quantities. For if the side of a square contain a feet, the diameter (see Example 3, p. 7) contains $\sqrt{2} \cdot a$ feet, and therefore the ratio of their measures is $1 : \sqrt{2}$. So that their measures are incommensurable.

14. There is *no practical difficulty* in dealing with incommensurable quantities. We can always find for their measures arithmetical expressions sufficiently accurate for all practical purposes.

15. A little consideration will convince the student that no measurement can in *practice* be made with *absolute* accuracy.

For example: A skilful mechanic is probably satisfied if in measuring some material two or three feet in length the error in his measurement is less than the thirty-second part of an inch. (The thirty-second part of an inch is less than half the height of the smallest letter on this page.) That is to say, he is satisfied if he make no greater error than about a *thousandth part of the whole length* to be measured. He would record such a measurement thus,

$$2 \text{ ft. } 3\tfrac{27}{32} \text{ inches,}$$

which is 891 thirty-second parts of an inch.

16. We will suppose that the length thus measured is the side of a square, and that the workman wishes to know to the same degree of accuracy as his measurement, what is the length of the diameter of the square.

He can find it thus.

The diameter of a square $= \sqrt{2} \times$ its side,

$\therefore$ the diameter of this square $= \sqrt{2} \times 891$ thirty-second parts of an inch.

Also $\sqrt{2} = 1{\cdot}414$ nearly,

$\therefore \sqrt{2} \times 891 = 1{\cdot}414 \times 891$

$= 1259{\cdot}8.$

$\therefore$ the required diameter $= 1260$ thirty-second parts of an inch, nearly.

The error being less than one thirty-second part of an inch.

17. The student will be able to see from the above example, that if the value of an incommensurable number is found to 4 figures, a very considerable degree of accuracy is attained. Also that a much greater degree of accuracy is attained for every additional figure.

18. It is no advantage in calculations such as the above to have the value of such quantities as $\sqrt{2}$ calculated to any greater degree of accuracy than the observed measurement. For instance, our calculation being correct as far as the whole numbers are concerned we should gain nothing by using $1{\cdot}4142$ instead of $1{\cdot}414$ for $\sqrt{2}$. This would give us the answer $1260{\cdot}0$ instead of $1259{\cdot}8$; the difference being a fifth of a thirty-second part of an inch, a quantity by hypothesis too small to be of any importance.

Example 1. The side of an equilateral triangle contains 2 feet; find, correct to the ten-thousandth part of a foot, the length of the perpendicular drawn from an angular point to the opposite side.

Here (as in Example 2, p. 6), let the perpendicular contain x feet, then



$$x^2 \text{ sq. ft.} = 2^2 \text{ sq. ft.} - 1^2 \text{ sq. ft.},$$
$$\therefore\ x^2 = 4 - 1 = 3,$$
$$\therefore\ x = \sqrt{3},$$
$$= 1{\cdot}7320 \text{ \&c.};$$

$\therefore$ the length of the perpendicular $= 1{\cdot}7320$ feet.

Example 2. Find the length, correct to the ten-thousandth part of a foot, of the side of the square described upon a diameter whose length is a feet.

Let x be the number of feet in each of the sides of the square.



Then x^2 sq. ft. $+ x^2$ sq. ft. $= a^2$ sq. ft.

$$\therefore x^2 + x^2 = a^2,$$

$$\therefore 2x^2 = a^2,$$

$$\therefore \sqrt{2} \,.\, x = a,$$

$$\therefore x = \frac{a}{\sqrt{2}} = \frac{a\sqrt{2}}{2}$$

$$= \frac{a}{2} \times (1{\cdot}4142) = a \times {\cdot}7071\ldots,$$

$\therefore$ the length of the perpendicular $= a \times {\cdot}7071\ldots$feet.

* EXAMPLES. IV.

(1) Find, correct to the thousandth part of a foot, the length of the diameter of a square whose side is 7 feet.

(2) Find, correct to a yard, the length of the diameter of a square whose side is one mile.

(3) Find, correct to the hundredth part of an inch, the hypotenuse of a right-angled triangle whose sides are 3 ft. $6\frac{1}{2}$ in. and 3 ft. 4 in. respectively.

(4) Find, to the nearest inch, the side of a square whose area is 1000 square yards.

(5) Find, correct to the tenth part of a foot, the diameter of a square field whose area is ten acres.

(6) Find, to the nearest inch, the side of a square field whose area is one acre.

(7) Find the height of an equilateral triangle whose side is 10 feet.

(8) Find the height of an equilateral triangle whose side is 5·32 feet.

(9) The top of a table measures 24·6 inches and 41·3 inches along two adjacent sides. What should the diameter measure if the table is rectangular?

(10) Find, to the nearest inch, the diameter of a lawn-tennis court whose length is 78 feet, and breadth 36 feet, supposing that it is properly marked out.

CHAPTER III.

On the Relation between the Circumference of a Circle and its Diameter.

19. The circumference of a circle is a line, and therefore it has length.

We might imagine the circumference of a circle to consist of a flexible wire; if the circular wire were cut at one point and straightened, we should have a straight line of the same length as the circumference of the circle.

20. A *polygon* is a figure enclosed by any number of straight lines.

21. A *regular* polygon has all its sides equal and all its angles equal.

22. The *perimeter* of a polygon is the sum of its sides.

23. If we have two circles in which the diameter of the first is greater than the diameter of the second, it is evident that the circumference of the first will be greater than the circumference of the second.



24. It seems, therefore, not unlikely that, if the diameter of the first circle be twice that of the second, the circumference of the first will be twice that of the second.

25. And also not unlikely that whatever be the ratio of the circumference of the first circle to its diameter, the same will be the ratio of the circumference of the second circle to its diameter.

This suggests that it is not unlikely that the circumference of a circle $= k$ times its diameter, where k is some *number* which is the same for all circles.

We shall presently prove that this is the case.

26. But although we can prove that

$$\frac{\text{the circumference of a circle}}{\text{its diameter}} = \text{a fixed numerical quantity,}$$

the method of calculating the value of this number is beyond the limits of an elementary treatise.

27. We shall therefore simply state here, what is proved in the Higher Trigonometry,

(i) that this numerical value is incommensurable,

(ii) that it is approximately 3·14159265 &c.

28. When we say that this number is incommensurable we mean (cf. Chapter II.) that its exact value cannot be stated as an *arithmetical* fraction.

It also happens that we have no short *algebraical* expression such as a surd, or combination of surds, which represents it exactly.

So that we have no *numerical* expression whatever, arithmetical nor algebraical, to represent *exactly* the ratio of the circumference of a circle to its diameter.

Hence the universal custom has arisen, of denoting its *exact* value by the letter π.

29. Thus π stands *always* for **the exact value of a certain incommensurable number,** whose approximate value is **3·14159265,** which number is **the ratio of the circumference of any circle to its diameter.**

It cannot be too carefully impressed on the student's memory that π stands for this number 3·14159265...&c., and for nothing else; just as 180 stands for the number one hundred and eighty, and for nothing else.

* 30. We proceed to prove that the ratio of the circumference of a circle to its diameter is the same for all circles.

The proof depends on the following important principle;

The length of the circumference of a circle is that to which the length of the perimeter of a regular inscribed polygon approaches, as the number of its sides is continually increased.

* 31. We take for granted that the *straight* line is the *shortest* line that can join two points.

* 32. Let *ABCDEF* be any regular polygon inscribed in a circle.



Then the side AB is shorter than any other line joining AB, so that the side AB is less than the arc AB;

therefore the perimeter of the polygon, viz.

$$AB + BC + CD + DE + EF + FA$$

is less than the sum of the arcs, that is, is less than the circumference of the circle.

Now let each of the *arcs* AB, BC, etc. be bisected in α, β, γ, δ, ϵ, η, and let the lines $A\alpha$, αB, $B\beta$, βC, etc. be joined;

then the figure $A\alpha B\beta C\gamma$ etc. is a regular polygon of twice as many sides as the first polygon.

And since the sides $A\alpha + \alpha B$ are together greater than the side AB,

it follows that the perimeter of the second polygon, viz.

$$A\alpha + \alpha B + B\beta + \beta C + C\gamma + \gamma D + \text{etc.}$$

is greater than the perimeter of the first.

But the perimeter of the second polygon is less than the circumference of the circle,

because each side is less than the corresponding arc.

Hence the perimeter of the second polygon is *nearer* the circumference, but is less than the circumference.

By bisecting the arcs of the second polygon we should get a third polygon, whose perimeter is nearer the circumference of the circle than the second.

It is clear that by continuing this process, we can get a polygon whose perimeter is as near as we please to the circumference of the circle. Hence,

The length of the circumference of a circle is that to which the length of the perimeter of a regular inscribed polygon approaches, as the number of its sides is continually increased.

* 33. PROP. *The ratio of the circumference of a circle to its diameter is the same for all circles.*

Let *ABCDEF*, *abcdef* be any two circles.

Let a *regular polygon* of any, the same number of sides be inscribed in each of them.

Join *A*, *B*, *C*, etc., *a*, *b*, *c*, etc. the angular points of the polygons to the centres *O*, *o* respectively.





Then *AOB* any one of the isosceles triangles in the first figure is equiangular with*, and therefore similar to, *aob* any one of the isosceles triangles in the second figure.

$$\therefore\ AB : OA = ab : oa, \qquad \text{[Euc. VI. 4.]}$$

and $$BC : OA = bc : oa,$$

and so on.

$$\therefore\ AB + BC + CD + \text{etc.} : OA = ab + bc + cd + \text{etc.} : oa;$$

or, the perimeter of the first polygon is to the radius of the

* For at *O* and at *o*, four right angles are each divided into the *same* number of equal angles, so that the *vertical* angles *AOB*, *aob* of the isosceles triangles *AOB*, *aob* are equal.

first circle as the perimeter of the second polygon is to the radius of its circle.

This is true *whatever* be the number of the sides of the two polygons.

And therefore it is true *however great* be the number of sides of the two polygons.

But the *circumferences* of the circles are what the *perimeters* of the polygons become, when the number of the sides is indefinitely increased. Therefore the circumference of the first circle is to its radius OA as the circumference of the second circle is to its radius oa.

Thus the ratio of the circumference of any circle to its radius is equal to the ratio of the circumference of any other circle to its radius.

So that the ratio $\frac{\text{circumference}}{\text{diameter}}$ is the same numerical quantity for all circles. Q. E. D.

34. We said above (Art. 29) that this number is 3·14159 etc., and that it is denoted by π.

Hence the circumference of any circle of radius r

$$= (3{\cdot}14159265 \text{ etc.}) \times \text{its diameter} = 2\pi r.$$

35. We may notice that $\frac{22}{7} = 3{\cdot}\dot{1}42857\dot{}$.

So that $\frac{22}{7}$ and π differ by less than a thousandth part of their value.

Also $\frac{355}{113} = 3{\cdot}1415929$ etc. So that $\frac{355}{113}$ may be used for π with sufficient accuracy for any practical purpose.

Hence $\frac{22}{7}$, 3·14159 and $\frac{355}{113}$ are each used for π according to the degree of accuracy required.

36. Of these 3·14159 is the most frequently used. The student should notice however that in *dividing* by π it will be more convenient to use $\frac{3\,5\,5}{1\,1\,3}$ than 3·14159.

37. The following results are instructive.

The ratio of the perimeter of a regular polygon inscribed in a circle to the diameter of the circle, when the polygon has

four sides, is $2\sqrt{2}$ $=2{\cdot}8284\ldots$

six sides, is 3 $=3$

eight sides, is $4\sqrt{(2-\sqrt{2})}$ $=3{\cdot}0614\ldots$

ten sides, is $\frac{5}{2}(\sqrt{5}-1)$ $=3{\cdot}0901\ldots$

twelve sides, is $3\sqrt{2}(\sqrt{3}-1)$ $=3{\cdot}1058\ldots$

twenty sides, is $5\{\sqrt{(3+\sqrt{5})}-\sqrt{(5-\sqrt{5})}\}$ $=3{\cdot}1287\ldots$

sixty sides, is $\dfrac{15}{2\sqrt{2}}\{(\sqrt{5}-1)(\sqrt{3}+1)$

$$-\sqrt{(10+2\sqrt{5})}\,(\sqrt{3}-1)\}=3{\cdot}1401\ldots$$

These numbers approach 3·14159 as the number of sides is increased, while the surd expression becomes more complicated.

The first of these results the student will be able to verify; the second is proved in Euclid IV. 15. The rest will be proved later on.

Example 1. The driving wheel of a locomotive engine is 5 ft. 6 in. high. What is its circumference?

Here we have a circle whose diameter is $5\frac{1}{2}$ feet;

$\therefore$ its circumference $=\pi\times 5{\cdot}5$ feet,

$=(3{\cdot}14159\ldots)\times 5{\cdot}5$ feet,

$=17{\cdot}278\ldots$ feet.

The circumference is 17 ft. 3 in. approximately.

Example 2. If a piece of wire 1 foot long be bent into the form of a circle, what will be the diameter of the circle?

Here the circumference $=1$ foot,

that is $\pi\times$ diameter $=1$ foot,

$$\therefore \text{ diameter} = \frac{1 \text{ foot}}{\pi} = \frac{7}{22} \times 1 \text{ foot}$$

$$= \frac{84}{22} \text{ inches}$$

$$= 3{\cdot}8 \text{ inches, nearly.}$$

If a greater degree of accuracy be desired, we must use $\frac{355}{113}$ instead of $\frac{22}{7}$.

We then get, the diameter $= 3{\cdot}7699$ inches,

$= 3{\cdot}77$ inches very nearly.

EXAMPLES. V.

In the answers of the first 12 of the following examples $\frac{22}{7}$ is used for π.

(1) Find the circumference of a circle whose diameter is one yard.

(2) Find the circumference of a circle whose radius is 4 feet.

(3) Find the circumference of a 48 inch bicycle wheel.

(4) The circumference of a circle is 10 feet; find its diameter.

(5) What must be the diameter of a locomotive driving wheel, that it may make 220 revolutions per mile?

(6) How many revolutions does a 36 inch bicycle wheel make per mile?

(7) How many more revolutions per mile does a 50 inch bicycle wheel make than one of 52 inches?

(8) A locomotive whose driving wheel is 5 feet high has an instrument to record the number of revolutions made. What number will the instrument record in running 100 miles?

(9) If the instrument in Question 8 indicates 3 revolutions per second, how many miles per hour is the engine running?

(10) What is the diameter of the driving wheel of a locomotive engine which makes 4 revolutions per second when the engine is going at the rate of 60 miles per hour?

(11) The large hand of the Westminster clock is 11 feet long; how many yards per day does its extremity travel? How far does the extremity move in a minute?

(12) The diameter of the whispering gallery in St Paul's is 108 feet; what is its circumference?

(13) Find the number of inches of wire necessary to construct a figure consisting of a circle with a regular hexagon inscribed in it, one of whose sides is 3 feet.

(14) How many inches of wire would be necessary in a figure similar to that in Question (13), if the circumference of the circle were ten feet?

(15) Find how many inches of wire are necessary to make a figure consisting of a circle and a square inscribed in it, when each side of the square is 2 feet.

(16) How many inches of wire are necessary for a figure similar to that in Question (15), when the circumference of the circle is 12 feet?

(17) Find the length of string necessary to string the handle of a cricket bat; having given the diameter of the handle $= 1\frac{1}{4}$ in., the length of the handle $= 12$ in., the diameter of the string $= \frac{1}{40}$th of an inch.

CHAPTER IV.

On the Measurement of Angles.

38. In elementary Geometry (Euclid I.—VI.) the angles considered are each always less than two right angles.

For example, in speaking of the angle *ROP* in Euclid we should always mean the angle less than two right angles,



not an angle measured in the opposite direction greater than two right angles.

39. In Trigonometry, by *the angle ROP* is meant, not the *present inclination* of the two lines *OR*, *OP* but, the **amount of turning** which *OP* has gone through when, starting from the position *OR*, it has turned about *O* into the position *OP*.

Example. Suppose a race run round a circular course. The position of any one of the competitors would be known, if we remark that he has described a certain angle about the centre of the course. Thus, if the distance to be run is three times round, the line joining each competitor to the centre would have to describe an angle of 12 right angles.

When we remark that a competitor has described an angle of $6\frac{2}{3}$ right angles, we record not only his present position, but the

total distance he has gone. He would in such a case have gone a little more than one and a half times round the course.

40. Definition. The **angle** between two lines OR, OP is the **amount of turning** about the point O which one of



the lines OP has gone through in turning from the position OR into the position OP.

41. The angle ROP may be the *geometrical* representative of an unlimited number of Trigonometrical angles.

(i) The angle ROP may represent the angle less than two right angles as in Euclid.

In this case OP has turned from the position OR into the position OP by turning about O in the direction *contrary* to that of the hands of a watch.

(ii) The angle ROP may represent the angle described by OP in turning from the position OR into the position OP in the *same* direction as the hands of a watch.

In the first case it is usual to say that the angle ROP is described in the *positive* direction, in the second that the angle is described in the *negative* direction.

(iii) The angle ROP may be the geometrical representation of any of the Trigonometrical angles formed by any number of complete revolutions in the *positive* or in the *negative* direction, added to either of the first two angles. (We shall return to this subject in Chapter IX.)

EXAMPLES. VI.

Give a geometrical representation of each of the following angles, the starting line being drawn in each case from the turning point towards the right.

1. $+3$ right angles.
2. $+5$ right angles.
3. $+4\frac{1}{2}$ right angles.
4. $+7\frac{1}{4}$ right angles.
5. -1 right angle.
6. $10\frac{2}{3}$ right angles.
7. $-10\frac{1}{3}$ right angles.
8. $+4$ right angles.
9. -4 right angles.
10. $4n$ right angles.
11. $(4n+2)$ right angles.
12. $-(4n+\frac{1}{2})$ right angles.

---

42. There are two methods of measuring angles.

(i) The rectangular measure.

(ii) The circular measure.

Rectangular Measure.

43. Angles are always measured *in practice* with the **right angle** (or part of the right angle) as unit.

44. The reasons why the right angle is chosen for a unit are:

(i) All right angles are equal to one another.

(ii) A right angle is practically easy to draw.

(iii) It is an angle whose size is very familiar.

45. The right angle is a large angle, and it is therefore subdivided for practical purposes.

It is usual to explain two methods of subdivision.

(i) The sexagesimal method.

(ii) The centesimal, or decimal, method *.

* See Art. 55.

I. *The Sexagesimal Method.*

46. In this method the right angle is divided into 90 equal parts, each of which is called a **degree**; each degree is subdivided into 60 equal parts, each of which is called a **minute**; and each minute is again subdivided into 60 equal parts, each of which is called a **second**.

Instruments used for measuring angles are subdivided accordingly; and the size of an angle is known when, with such an instrument, it has been observed that the angle contains a certain number of degrees, and a certain number of minutes beyond the number of complete degrees, and a certain number of seconds beyond the number of complete minutes.

Thus an angle might be recorded as containing 79 degrees + 18 minutes + 36·4 seconds.

Degrees, minutes, and seconds are indicated respectively by the symbols °, ′, ″, and the above angle would be written

$$79^\circ \,.\, 18' \,.\, 36{\cdot}4''.$$

* II. *The Centesimal or Decimal Method.*

47. The other method of subdivision is the Centesimal or Decimal. Here each right angle is divided into 100 equal parts each of which is called a **grade**; each grade is subdivided into 100 equal parts, each of which is called a **minute**; and each minute is again subdivided into 100 equal parts, each of which is called a **second**.

Instruments of observation would be subdivided accordingly, and any observed angle would be recorded as containing so many grades + so many minutes + so many seconds.

Grades, minutes and seconds are indicated respectively by the symbols ^{g}, $'$, $''$. So that an angle of 26 grades + 19 minutes + 34·2 seconds would be written

$$36^g\ 19'\ 34{\cdot}2''.$$

*48. It will be observed that this method is simply that of the decimal system of notation.

The above angle for example $= \frac{36}{100} + \frac{19}{10000} + \frac{34{\cdot}2}{1000000}$ of a right angle.

That is ·3619342 of a right angle.

This is equal to 36·19342 of a grade.

Also to 3619·342 of a minute.

Also to 361934·2 of a second.

Example. Express $302^g\ 2'\ 4{\cdot}6''$ as the decimal of a right angle.

$$\text{This angle} = \frac{302}{100} + \frac{2}{10000} + \frac{4{\cdot}6}{1000000} \text{ of a right angle}$$
$$= 3{\cdot}0202046 \text{ of a right angle.}$$

*49. Hence, to express an angle given in grades, minutes and seconds as the decimal of a right angle, we have only to observe that the *first and second* decimal places are occupied by the grades, the *third and fourth* decimal places are occupied by the minutes, and the *fifth and sixth* decimal places are occupied by the seconds.

*50. The same observation will enable us to express in grades, minutes, and seconds an angle given as the decimal of a right angle.

Example. Express 3·4650023 of a right angle in grades, minutes, and seconds.

3 right angles = 300 grades.

·46 of a right angle = 46 grades.

·00,50 of a right angle = 50 minutes.

·00,00,02,3 of a right angle = 2·3 seconds.

Therefore the angle is $346^g\ 50'\ 2{\cdot}3''$.

* EXAMPLES. VII.

Express as the decimal of a right angle,

(1) $63^g\ 21'\ 18''$.
(2) $104^g\ 26'\ 99{\cdot}1''$.
(3) $2^g\ 18'\ 27''$.
(4) $3^g\ 29'\ 48{\cdot}94''$.
(5) $62'\ 41''$.
(6) $1000^g\ 8'\ 12''$.
(7) $32^g\ 4'\ 5{\cdot}2'$
(8) $1^g\ 2'\ 3{\cdot}4''$
(9) $69^g\ 0'\ 7{\cdot}1''$.
(10) $119^g\ 3'\ 0{\cdot}45''$.
(11) $1006^g\ 18'\ 1''$
(12) $2^g\ 26'\ 4{\cdot}8''$.

Express in grades, minutes and seconds,

(13) ·367891 of a right angle.
(14) 1·043021 of a right angle.
(15) ·012003 of a right angle.
(16) ·00102 of a right angle.
(17) ·0625 of a right angle.
(18) 3·02125 of a right angle.
(19) 1·001 of a right angle.
(20) ·0101001 of a right angle.
(21) 6·451 of a right angle.
(22) ·023 of a right angle.
(23) ·00011 of a right angle.
(24) ·00001 of a right angle.

---

51. An angle given in degrees, minutes, and seconds may be expressed as the decimal of a right angle by the usual method.

Example. Express $39^0\ 4'\ 27''$ as the decimal of a right angle.

60) 27 seconds
60) 4·45 minutes
90) 39·07416666 etc. degrees
·43415740740 etc. right angles

Answer. ·43415$\dot{7}$4$\dot{0}$ of a right angle.

52. An angle given as the decimal of a right angle may be expressed in degrees, minutes, and seconds by the converse of the above.

Example. Express ·43415$\dot{7}$4$\dot{0}$ of a right angle in degrees, minutes, and seconds.

·43415740740 etc. right angles
90
39·07416666600 degrees

The last two figures would be 66 if we were to write down the recurring part to more figures.

This gives 39·07416666666 etc. degrees
60
4·4499999960 minutes
that is 4·44$\dot{9}$ minutes
or 4·45 minutes
60
27·00 seconds.

The result is 39° 4′ 27″.

*53. We have seen that an angle expressed as the decimal of a right angle can be at once expressed in grades, minutes, and seconds.

Hence an angle expressed in degrees, minutes, and seconds, can be expressed in grades, etc. by first reducing the angle to the decimal of a right angle.

Example. Express 39° 4′ 27″ in grades, minutes, and seconds.

This angle is ·43415$\dot{7}$4$\dot{0}$ of a right angle, by Art. 51
and this =43ᵍ 41ᶜ 57·$\dot{4}$0$\dot{7}$ᶜᶜ.

*54. An angle given in grades, minutes, and seconds can be expressed in degrees, minutes, and seconds by first expressing the angle as the decimal of a right angle.

Example. Express $\cdot 43^g\ 41'\ 57\cdot 40\dot{7}''$ in degrees, minutes, and seconds.

This angle is $\cdot 43415\dot{7}40$ of a right angle, which is $39^0\ 4'\ 27''$ from Art. 52.

* EXAMPLES. VIII.

Express each of the following angles (i) as the decimal of a right angle, (ii) in grades, minutes, and seconds;

(1) $8^0\ 15'\ 27''$.
(2) $6^0\ 4'\ 30''$.
(3) $97^0\ 5'\ 15''$.
(4) $16^0\ 14'\ 19''$.
(5) $132^0\ 6'$.
(6) 49^0.

Express in degrees, minutes and seconds,

(7) $1^g\ 37'\ 50''$.
(8) $8^g\ 75'$.
(9) $170^g\ 45'\ 35''$.
(10) $24^g\ 0'\ 25''$.
(11) $18^g\ 1'\ 15''$.
(12) 35^g.

55. The decimal or centesimal system of subdividing a right angle was proposed by the French at the commencement of the present century; but, although it possesses many advantages over the established method, no one has been found willing to undertake the great expense that would have to be incurred in rearranging all tables and all books of reference, and all the records of observations, which would have to be transferred from the old system to the new, before the advantages of the decimal system could be felt. Thus the decimal system of angular measurement has never been used even in France, and in all probability never will be used in practical work.

On Circular Measure.

56. By the following construction we get an angle of great importance in Trigonometry.

On the circumference of a circle whose centre is O



let an *arc* RS be measured so that its length is equal to the *radius* of the circle, and let R and S be joined to the centre.

57. We are about to prove (Art. 60) that this angle ROS is a *fixed* fraction of a right angle, so that all such angles are equal to one another.

58. We may state the same thing thus—We are about to prove that if we take any number of different circles, and measure on the circumference of each an arc equal in length to its radius, then the angles at the centres of these circles which stand on these arcs respectively, will be all of the same size.

59. Definition. The angle which at the centre of a circle stands on an arc equal in length to the radius of the circle is called a **Radian.**

60. *To prove that all Radians are equal to one another.*

Since the Radian at the centre of a circle stands on an arc equal in length to the radius,

and an angle of two right angles at the centre of a circle stands on half the circumference,

and since angles at the centre of a circle are to one another as the arcs on which they stand (Euc. VI. 33),

therefore the radian is to an angle of two right angles as the radius is to half the circumference;

that is, as the diameter is to the whole circumference;

that is, in the constant ratio $1 : \pi$.

Therefore the radian $= \dfrac{2 \text{ right angles}}{\pi}$.

That is, the radian is a fixed fraction of a right angle.

But all right angles are equal to one another.

Therefore all radians are equal to one another. Q. E. D.

61. Thus the radian possesses the qualification most essential in a unit, viz. it is always the same.

The student will find, in the theoretical part of Trigonometry, that many expressions can be written more shortly when a radian is used for the unit of angle, than when any other unit is used.

62. Thus the reasons why a radian is used as a unit are :

(i) All radians are equal to one another.

(ii) Its use simplifies many formulæ in Theoretical Trigonometry.

63. The system of angular measurement in which a **radian** is the unit is called **Circular Measure.**

Therefore the *circular measure of an angle* is the *number of radians* which the angle contains.

64. A radian $= \dfrac{1}{\pi} \times 2$ right angles,

$= \frac{1}{3}\frac{1}{5}\frac{3}{5}$ of 180° nearly,

$= 57{\cdot}2957 \ldots .$ degrees.

The student should notice that a radian is a little less than an angle of an equilateral triangle.

65. Circular measure is, as we have said, used in *theoretical* investigations, in which the angle under consideration is almost always expressed by a *letter*. This is usually one of the *Greek* letters $\alpha, \beta, \gamma \ldots, \phi, \theta, \psi \ldots$.

Strictly speaking these letters represent *numbers*, i.e. measures; so that some unit of *angle* must be *understood* in such an expression as 'the angle θ.' (Art. 2.)

For this reason, when an angle is denoted by a *Greek* letter such as α, β, γ, etc., θ, ϕ, ψ, etc., it is understood that *circular measure* is the measure used, unless the contrary is *expressly stated*.

So that 'the angle θ' means 'θ radians.'

Similarly 'the angle π' means 'π radians' or '3·14159... radians,' that is two right angles.

[*Note*. It will also be convenient, in using such letters as A, B, $C \ldots S$, T, etc. to represent angles, to agree that the *unit* understood with this kind of letter shall be a *degree*, so that when A stands for an angle, that angle contains A degrees.]

66. In *numerical* examples it will be necessary to use some letter (c suppose) to denote a radian.

Thus 2^c denotes 'two radians.'

67. Strictly speaking then 'the angle θ' should be written θ^c. (Just as in speaking of 'the angle ninety,' we ought to say ninety *degrees*.) But if it is clearly understood that 'the angle θ' means θ *radians*, there can be no ambiguity in the expression.

68. The student cannot too carefully notice, that unless an *angle* is obviously referred to, the letters θ, ϕ,... α, β,... stand for *mere numbers*.

Thus as we have said above (29) π *stands for a number and a number only*, viz. 3·14159....., but in the expression 'the angle π' that is 'the angle 3·14159.....' there must be *some unit* understood. The *unit understood* here is a radian, and therefore 'the angle π' stands for 3·14159.....', that is *two right angles*.

Hence, *when an angle is referred to, π is a very convenient abbreviation for* **two right angles**.

69. To express in degrees or grades an angle given in radians, we first express the angle in right angles, remembering that

2 right angles $= \pi$ radians.

Example. How many degrees are there in the angle whose circular measure is 2?

$$\text{This angle} = 2 \text{ radians} = 2 \times \frac{2 \text{ right angles}}{\pi} = \frac{4}{\pi} \text{ right angles},$$

$$= \frac{4 \times 90^0}{\pi} = \frac{360^0}{\pi},$$

$$\therefore \text{ the angle contains } \frac{360}{\pi} \text{ degrees.}$$

*70. If D, G and α be the number of degrees, grades and radians respectively in any angle, then

$$\frac{D}{180} = \frac{G}{200} = \frac{\alpha}{\pi}.$$

For each fraction is the ratio of the angle to two right angles.

Example. Find the number of degrees in two radians.

Let D be the number, then

$$\frac{D}{180} = \frac{2}{\pi},$$

$$\therefore\ D = \frac{360}{\pi}.$$

EXAMPLES. IX.

1. Express the following angles in rectangular measure.

(1) π. (2) $\frac{3\pi}{4}$. (3) 1^c.

(4) 3^c. (5) $3{\cdot}14159265^c$ etc. (6) $\frac{2^c}{\pi}$.

(7) θ. (8) $\cdot00314159^c$ etc. (9) 10π.

2. Express the following angles in circular measure.

(1) 180^0. (2) 360^0. (3) 60^0.

(4) $22\frac{1}{2}^0$. (5) 1^0. (6) $57{\cdot}295^0$ etc.

(7) n^0. (8) $\frac{90^0}{\pi}$. (9) A.

* 3. Express the following angles in circular measure.

(1) $33^g\ 33'\ 33{\cdot}3''$. (2) 50^g. (3) $16{\cdot}6^g$.

(4) 1^g. (5) $1'$. (6) $10''$.

(7) n^g. (8) $\frac{200^g}{\pi}$. (9) 1000^g.

* 4. Find the ratio of

(1) 45^0 to $\frac{3\pi}{4}$. (2) 60^0 to 60^g.

(3) 25^g to $22^0\ 30'$. (4) 24^g to 2^c.

(5) $1{\cdot}75^c$ to $\frac{100^0}{\pi}$. (6) 1^0 to 1^c.

*71. *To prove that the measure of an angle at the centre of a circle in radians* (i.e. *in Circular Measure*) *is the ratio of the arc on which it stands to the radius of the circle.*



Since angles at the centre of a circle are to one another as the arcs on which they stand (Euc. VI. 33),

Therefore any angle ROP at the centre of a circle is to the radian as its arc RP is to the arc of the radian.

But the arc of the radian is equal to the radius,

Therefore $\frac{\text{any angle } ROP}{\text{the radian}}$ is equal to $\frac{\text{its arc } RP}{\text{the radius}}$.

And therefore any angle ROP is equal to $\frac{\text{its arc}}{\text{radius}} \times$ (a radian).

That is, the *measure* α of an angle *in radians* is the ratio $\frac{\text{arc}}{\text{radius}}$; or, $\alpha = \frac{\text{arc}}{\text{radius}}$.

Hence (cf. Art. 70) $\frac{D}{180} = \frac{G}{200} = \frac{\alpha}{\pi} = \frac{\text{arc}}{\text{radius}} \times \frac{1}{\pi}$.

Example. Find the number of grades in the angle subtended by an arc 46 ft. 9 in. long, at the centre of a circle whose radius is 25 feet.

The angle stands on an arc of $46\frac{3}{4}$ ft. and the radian, at the centre of the same circle, stands on an arc of 25 feet.

$$\therefore \text{ the angle} = \frac{46\frac{3}{4}}{25} \text{ radians}, \quad = \frac{187}{100} \times \frac{2 \text{ right angles}}{\pi},$$

$$= \frac{187}{100} \times \frac{200^g}{\pi}, = 119^g \text{ nearly}.$$

* EXAMPLES. X.

(In the answers $\frac{22}{7}$ is used for π.)

(1) Find the number of radians in an angle at the centre of a circle of radius 25 feet, which stands on an arc of $37\frac{1}{2}$ feet.

(2) Find the number of degrees in an angle at the centre of a circle of radius 10 feet, which stands on an arc of 5π feet.

(3) Find the number of right angles in the angle at the centre of a circle of radius $3\frac{2}{11}$ inches, which stands on an arc of 2 feet.

(4) Find the number of French minutes in the angle at the centre of a circle of radius 8 ft. 4 inches, which stands on an arc of 1 inch.

(5) Find the length of the arc subtending an angle of $4\frac{1}{2}$ radians at the centre of a circle whose radius is 25 feet.

(6) Find the length of an arc of eighty degrees on a circle of 4 feet radius.

(7) Find the length of an arc of sixty grades on a circle of ten feet radius.

(8) The angle subtended by the diameter of the Sun at the eye of an observer is 32′; find approximately the diameter of the Sun if its distance from the observer be 90,000,000 miles.

(9) A railway train is travelling on a curve of half a mile radius at the rate of 20 miles an hour; through what angle has it turned in 10 seconds?

(10) A railway train is travelling on a curve of two-thirds of a mile radius, at the rate of 60 miles an hour; through what angle has it turned in a quarter of a minute?

(11) Find approximately the number of English seconds contained in the angle which subtends an arc one mile in length at the centre of a circle whose radius is 4000 miles.

(12) If the radius of a circle be 4000 miles, find the length of an arc which subtends an angle of 1″ at the centre of the circle.

(13) If in a circle whose radius is 12 ft. 6 in. an arc whose length is ·6545 of a foot subtends an angle of 3 degrees, what is the ratio of the diameter of a circle to its circumference?

(14) If an arc 1·309 feet long subtend an angle of $7\frac{1}{2}$ degrees at the centre of a circle whose radius is 10 feet, find the ratio of the circumference of a circle to its diameter.

(15) On a circle 80 feet in radius it was found that an angle of $22^0\ 30'$ at the centre was subtended by an arc 31 ft. 5 in. in length; hence calculate to four decimal places the numerical value of the ratio of the circumference of a circle to its diameter.

(16) If the diameter of the moon subtend an angle of 30′, at the eye of an observer, and the diameter of the sun an angle of 32′, and if the distance of the sun be 375 times the distance of the moon, find the ratio of the diameter of the sun to that of the moon.

(17) Find the number of radians in (i.e. the circular measure of) 10″ correct to 3 significant figures. (Use $\frac{355}{113}$ for π.)

(18) Find the radius of a globe such that the distance measured upon its surface between two places in the same meridian, whose latitudes differ by $1^0\ 10'$, may be one inch.

(19) Two circles touch the base of an isosceles triangle at its middle point, one having its centre at, and the other passing through the vertex. If the arc of the greater circle included within the triangle be equal to the arc of the lesser circle without the triangle, find the vertical angle of the triangle.

(20) By the construction in Euc. I. 1, prove that the unit of circular measure is less than 60^0.

(21) On the 31st December the Sun subtends an angle of 32′ 36″, and on 1st July an angle of 31′ 32″; find the ratio of the distances of the Sun from the observer on those two days.

(22) Show that the measure of the angle at the centre of a circle of radius r, which stands on an arc a, is $\frac{k.a}{r}$, where k depends solely on the unit of angle employed.

Find k when the unit is (i) a radian, (ii) a degree.

*72. Questions concerning angles expressed in different systems of measurement are easily solved by *expressing each angle in right angles.*

Example 1. The sum of the measure of an angle in degrees and twice its measure in radians is $23\frac{2}{7}$, find its measure in degrees ($\pi=\frac{22}{7}$).

Let the angle contain x right angles.

Then the measure of the angle in degrees $=90x$,

" " " " " radians $=\frac{\pi}{2}x$.

$$\therefore\ 90x+2.\frac{\pi}{2}x=23\tfrac{2}{7},$$

$$\therefore\ 90x+\tfrac{22}{7}x=1\tfrac{63}{7},$$

$$\therefore\ 652x=163,$$

$$\therefore\ x=\tfrac{1}{4}.$$

The angle is $\frac{1}{4}$ of a right angle, that is $22\frac{1}{2}^0$, nearly.

Example 2. The three angles of a triangle are in arithmetical progression, and the measure of the least in grades is to that of the greatest in circular measure as $120 : \pi$. Express each angle in degrees.

Let the angles contain $x-y$, x, $x+y$ **right angles** respectively; they are then in A.P.

Their *sum* is $3x$ right angles; but since they are the angles of a triangle, their *sum* is 2 right angles;

$$\therefore 3x=2,$$

$$\therefore x=\tfrac{2}{3}.$$

Again, the least angle contains $(x-y)\times 100$ grades, and the greatest angle contains $(x+y)\frac{\pi}{2}$ radians,

$$\therefore 100\,(x-y) : \frac{\pi}{2}(x+y)=120 : \pi.$$

$$\therefore 100\,(x-y)=60\,(x+y),$$

$$\text{or, } 40x=160y,$$

$$\text{or, } x=4y.$$

$$\therefore 4y=\tfrac{2}{3},$$

$$\text{because } x=\tfrac{2}{3},$$

$$\text{or, } y=\tfrac{1}{6}.$$

$$\therefore x-y=\tfrac{2}{3}-\tfrac{1}{6}=\tfrac{1}{2},\quad x=\tfrac{2}{3},\quad x+y=\tfrac{2}{3}+\tfrac{1}{6}=\tfrac{5}{6}.$$

Thus the angles contain $\frac{1}{2}$, $\frac{2}{3}$, and $\frac{5}{6}$ right angles respectively;

therefore the angles are 45^0, 60^0, 75^0.

** EXAMPLES. XI.

(In the following examples the answers will be given in terms of π.)

(1) The sum of the degrees and of the grades in a certain angle is 38; find its circular measure.

(2) The difference of two angles is 20^g, and their sum is 48^0; find them.

(3) One angle is double of a second, and the sum of their measures in degrees and in grades respectively is 140; express the angles in degrees.

(4) Two angles are in the ratio of 4 : 5, and the difference of their measures in grades and in degrees respectively is $2\frac{1}{2}$; find the angles in degrees.

(5) The difference between two angles is $\frac{\pi}{9}$, and their sum is 56 degrees; find the angles.

(6) If the three angles of a triangle are in arithmetical progression, show that the mean angle is 60^0.

(7) The three angles of a triangle are in arithmetical progression, and the number of grades in the least is to the number of degrees in the mean as 5 : 6. Find the angles in degrees.

(8) The three angles of a triangle are in arithmetical progression, and the number of grades in the greatest is to the number of degrees in the sum of the other two as 10 : 11. Find the angles in degrees.

(9) The three angles of a triangle are in arithmetical progression, and the number of grades in the least is to the number of radians in the greatest as $200 : 3\pi$. Express the angles in grades.

(10) If D be the number of degrees and G the number of grades in any angle, prove that $G - D = \frac{1}{9}D$.

(11) If M be the number of English minutes and m the number of French minutes in any angle, prove that

$$2M - m = \tfrac{4}{27}M.$$

(12) If G, D and C be the number of grades, degrees and radians in any angle, prove that $G - D = \dfrac{20C}{\pi}$.

(13) If an angle be expressed in French minutes, show that it will be transferred to English minutes by multiplying by ·54.

(14) Divide $33^0\ 6'$ into two parts so that the number of English seconds in one part may be equal to the number of French seconds in the other part.

(15) Find the ratio of $9^0\ 27'$ to $12^g\ 50'$.

(16) Find the number of radians in an angle of n English minutes.

(17) Express in each of the three systems of angular measurement the angles

(i) of a regular hexagon,

(ii) of a regular octagon,

(iii) of a regular quindecagon.

(18) Show that the number of degrees in an angle of a regular decagon is to the number of grades in an angle of a regular pentagon in the ratio of 6 : 5.

(19) Show that the number of grades in an angle of a regular pentagon is equal to the number of degrees in an angle of a regular hexagon.

(20) Find in English minutes the difference between the angle of a regular polygon of 48 sides and two right angles.

(21) If we take for unit the angle between a side of a regular quindecagon and the next side produced, find the measures (i) of a right angle, (ii) of a radian.

(22) Find the unit when the sum of the measures of a degree and of a grade is 1.

(23) What is the unit when the sum of the measures of 9^0 and of 5^g is $\frac{3}{20}$?

(24) If the measure of b grades is a, find the measure of c degrees.

(25) What is the unit when the sum of the measures of a grades and of b degrees is c?

(26) The number of grades in a certain angle exceeds the number of degrees in it by $\frac{1}{10}$ of the number of degrees in a radian. If this angle be taken as unit, what is the measure of a right angle?

(27) The three numbers which express the three angles of a triangle are all equal, and the units of angle in each are respectively a degree, a grade and the sum of a degree and a grade; express each of the angles in circular measure.

(28) The three angles of a triangle have the same measure when expressed in degrees, grades and radians respectively; find this measure.

(29) The measures of the angles of a triangle in degrees, grades and radians respectively are in the ratio of 1 : 10 : 100; find the number of radians in the smallest angle.

(30) The interior angles of an irregular polygon are in A.P.; the least angle is 120^0; and the common difference 5^0: find the number of sides.

CHAPTER V.

The Trigonometrical Ratios.

73. Let *ROE* be any angle (see the figure in Art. 83). In one of the lines containing the angle take any point *P*, and from *P* draw *PM* perpendicular to the other line *OR*.

Then, in the right-angled triangle *OPM*, formed from the angle *ROE*,

(i) the side *MP*, which is *opposite the angle under consideration*, is called the **perpendicular**;

(ii) the side *OP*, which is *opposite the right angle*, is called the **hypotenuse**;

(iii) the *third* side *OM*, which is adjacent to the right angle and to the angle under consideration, is called the **base.**

From these three,—perpendicular, hypotenuse, base,—we can form *three* different sets containing two each.

The **ratios** or *fractions* formed from these sets, viz.

(i) $\frac{\text{perpendicular}}{\text{hypotenuse}}$, (ii) $\frac{\text{base}}{\text{hypotenuse}}$, (iii) $\frac{\text{perpendicular}}{\text{base}}$,

and the ratios formed by inverting each of them, viz.

(iv) $\frac{\text{hypotenuse}}{\text{perpendicular}}$, (v) $\frac{\text{hypotenuse}}{\text{base}}$, (vi) $\frac{\text{base}}{\text{perpendicular}}$,

will be found to be of great importance in treating of any angle *ROE*. Accordingly to each of these six ratios has been given a separate *name* (Art. 75).

74. The student should observe carefully

(i) that each *ratio*, such as $\frac{\text{perpendicular}}{\text{hypotenuse}}$, is a mere *number*;

(ii) that, as we shall prove in Art. 83, these ratios remain unchanged as long as the angle remains unchanged;

(iii) that if the angle be altered ever so slightly, there is a consequent alteration in the value of these ratios.

[For, let ROE, ROE' be two angles which are nearly equal;



Let $OP = OP'$; then OM is *not* $= OM'$, and therefore the ratios $\frac{OM}{OP}$ and $\frac{OM'}{OP'}$ are *not* equal: also MP is *not* $= M'P'$ and therefore the ratios $\frac{MP}{OP}$ and $\frac{M'P'}{OP'}$ are *not* equal.]

(iv) that by giving **names** to these ratios we are enabled to apply the methods of Algebra to the Geometry of Euclid VI., just as in Chapter I. we applied the methods of Algebra to Euc. I. 47.

The student is recommended to pay careful attention to the following definitions. He should be able to write them out in the exact words in which they are printed.

75. DEFINITION. *To define the three principal Trigonometrical Ratios of an angle.*



Let ROE be an angle.

In OE one of the lines containing the angle take any point P, and from P draw PM perpendicular to the other line OR, or, if necessary, to RO produced.

Then, in the right-angled triangle OPM, the side MP, which is *opposite the angle under consideration*, is called the *perpendicular*.

The side OP, which is *opposite the right angle*, is called the *hypotenuse*.

The *third* side OM (which is adjacent to the right angle and to the angle under consideration) is called the *base*.

Then the ratio

(i) $\dfrac{MP}{OP} = \dfrac{\text{perpendicular}}{\text{hypotenuse}}$ is called the **sine** of the angle ROE.

(ii) $\dfrac{OM}{OP} = \dfrac{\text{base}}{\text{hypotenuse}}$ „ **cosine** „

(iii) $\dfrac{MP}{OM} = \dfrac{\text{perpendicular}}{\text{base}}$ „ **tangent** „

NOTE. The order of the letters, in MP, OM and OP, indicates direction and decides their algebraical signs. [Art. 132.]

76. If A stand for the angle ROE, these ratios are called sine A, cosine A and tangent A, and are usually abbreviated thus:

$$\sin A, \quad \cos A, \quad \tan A.$$

77. There are three other Trigonometrical Ratios, formed by *inverting* the sine, cosine and tangent respectively, which are called the cosecant, secant, and cotangent respectively.

78. *To define the three other Trigonometrical Ratios of any angle.*

The same construction and figure as in Art. 75 being made, then the ratio

(iv) $\dfrac{OP}{MP} = \dfrac{\text{hypotenuse}}{\text{perpendicular}}$ is called the **cosecant** of the angle ROE.

(v) $\dfrac{OP}{OM} = \dfrac{\text{hypotenuse}}{\text{base}}$ „ **secant** „

(vi) $\dfrac{OM}{MP} = \dfrac{\text{base}}{\text{perpendicular}}$ „ **cotangent** „

79. Thus if A stand as before for the angle ROE, these ratios are called cosecant A, secant A, and cotangent A. They are abbreviated thus,

$$\text{cosec } A, \quad \sec A, \quad \cot A.$$

80. From the definition it is clear that

$$\text{cosec } A = \frac{1}{\sin A},$$

$$\sec A = \frac{1}{\cos A},$$

$$\cot A = \frac{1}{\tan A}.$$

81. The above definitions **apply to an angle of any magnitude.** (We shall return to this subject in Chapter X.)

For the present the student may confine his attention to angles which are each less than a right angle.

82. The powers of the Trigonometrical Ratios are expressed as follows:

$$(\sin A)^2, \text{ i.e. } \left(\frac{\text{perpendicular}}{\text{hypotenuse}}\right)^2, \text{ is written } \sin^2 A,$$

$$(\cos A)^3, \text{ i.e. } \left(\frac{\text{base}}{\text{hypotenuse}}\right)^3, \text{ is written } \cos^3 A,$$

and so on.

The student must notice that '$\sin A$' is a *single symbol.* It is the *name* of a *number*, or *fraction*, belonging to the angle A; and if it be at any time convenient, we may denote *sin A* by a *single letter*, such as s or x. Also $\sin^2 A$ is an abbreviation for $(\sin A)^2$, that is for $(\sin A) \times (\sin A)$. Such abbreviations are used because they are *convenient.*

83. *The Trigonometrical Ratios are always the same for the same angle.*



Take any angle ROE; let P be any point in OE one of the lines containing the angle, and let P', P'' be any two points in OR the other line containing the angle. Draw PM perpendicular to OR, and $P'M'$, $P''M''$ perpendiculars to OE.

Then the three triangles OMP, $OM'P'$, $OM''P''$ each contain a right angle, and they have the angle at O common; therefore their third angles must be equal.

Thus the three triangles are equiangular.

Therefore the ratios $\frac{MP}{OP}$, $\frac{M'P'}{OP'}$, $\frac{M''P''}{OP''}$ are all equal. (Eu. VI. 4.)

But each of these ratios is $\frac{\text{perpendicular}}{\text{hypotenuse}}$ *with reference to the angle at* O; that is, they are each *sin* ROE.

Thus, sin ROE is the same *whatever* be the position of the point P on *either* of the lines containing the angle ROE.

Therefore sin ROE is always the same.

84. A similar proof holds good for each of the other ratios.

85. Also if two angles are equal, it is clear that the numerical values of their Trigonometrical Ratios will be the same.

We have already shown (Art. 74), that the values of these ratios are different for different angles.

Hence for each particular value of A, sin A, cos A, tan A, etc. have *definite numerical values*.

Example. We shall prove (Art. 92) that

$$\sin 30^0 = \tfrac{1}{2} = \cdot 5, \quad \cos 30^0 = \frac{\sqrt{3}}{2} = \cdot 8660\ldots, \quad \tan 30^0 = \frac{1}{\sqrt{3}} = \cdot 577\ldots$$

86. In the following examples the student should notice

(i) the *angle* referred to,

(ii) that there is a *right angle* in the same triangle as the angle referred to,

(iii) the *perpendicular*, which is opposite the angle referred to, and is perpendicular to one of the lines containing the angle,

(iv) the *hypotenuse*, which is opposite the right angle,

(v) the *base*, the third side of the triangle.

Example. In the second figure on the next page, in which BDA is a right angle, find sin DBA and cos DBA.

In this case

(i) DBA is the *angle*.

(ii) BDA is a *right angle* in the same triangle as the angle DBA.

(iii) DA is the *perpendicular*, for it is opposite DBA and is perpendicular to BD.

(iv) BA is the *hypotenuse*.

(v) BD is the *base*.

Therefore sin DBA, which is $\frac{\text{perpendicular}}{\text{hypotenuse}}$, $= \frac{DA}{BA}$,

cos DBA, which is $\frac{\text{base}}{\text{hypotenuse}}$, $= \frac{BD}{BA}$.

EXAMPLES. XII.

(1) Let ABC be any triangle and let AD be drawn perpendicular to BC. Write down the *perpendicular*, and the *base* when the following angles are referred to: (i) the angle ABD, (ii) the angle BAD, (iii) the angle ACD, (iv) the angle DAC.



(2) Write down the following ratios in the above figure; (i) sin *BAD*, (ii) cos *ACD*, (iii) tan *DAC*, (iv) sin *ABD*, (v) tan *BAD*, (vi) sin *DAC*, (vii) cos *DCA*, (viii) tan *DCA*, (ix) cos *ABD*, (x) sin *ACD*.

(3) Let *ACB* be any angle and let *ABC* and *BDC* be right angles; (see next figure). Write down *two* values for each of the following ratios; (i) sin *ACB*, (ii) cos *ACB*, (iii) tan *ACB*, (iv) sin *BAC*, (v) cos *BAC*, (vi) tan *BAC*.



(4) In the accompanying figure *BDC*, *CBA* and *EAC* are right angles. Write down (i) sin *DBA*, (ii) sin *BEA*, (iii) sin *CBD*, (iv) cos *BAE*, (v) cos *BAD*, (vi) cos *CBD*, (vii) tan *BCD*, (viii) tan *DBA*, (ix) tan *BEA*, (x) tan *CBD*, (xi) sin *DAB*, (xii) sin *BAE*.

(5) Let ABC be a right-angled triangle such that $AB=5$ ft., $BC=3$ ft., then AC will be 4 ft.



Find the sine, cosine and tangent of the angles at A and B respectively.

In the above triangle if A stand for the angle at A and B for the angle at B, show that $\sin^2 A+\cos^2 A=1$, and that $\sin^2 B+\cos^2 B=1$.

(6) If ABC be any right-angled triangle with a right angle at C, and let A, B, and C stand for the angles at A, B and C respectively, and let a, b and c be the measures of the sides opposite the angles A, B and C respectively.

Show that $\sin A=\frac{a}{c}$, $\cos A=\frac{b}{c}$, $\tan A=\frac{a}{b}$

Show also that $\sin^2 A+\cos^2 A=1$.

Show also that (i) $a=c.\sin A$, (ii) $b=c.\sin B$, (iii) $a=c.\cos B$, (iv) $b=c.\cos A$, (v) $\sin A=\cos B$, (vi) $\cos A=\sin B$, (vii) $\tan A=\cot B$.

(7) The sides of a right-angled triangle are in the ratio 5 : 12 : 13. Find the sine, cosine and tangent of each acute angle of the triangle.

(8) The sides of a right-angled triangle are in the ratio $1 : 2 : \sqrt{3}$. Find the sine, cosine and tangent of each acute angle of the triangle.

(9) Prove that if A be either of the angles of the above two triangles $\sin^2 A+\cos^2 A=1$.

CHAPTER VI.

On the Trigonometrical Ratios of Certain Angles.

87. The Trigonometrical Ratios of an angle are *numerical quantities simply*, as their name ratio implies. They are in nearly all cases incommensurable numbers.

Their practical value has been found for all angles between 0 and 90°, which differ by 1′; and a list of these values will be found in any volume of Mathematical Tables.

The student is recommended to get a copy of Chambers' Mathematical Tables for instruction and reference.

88. The finding the values of these Ratios has involved a large amount of labour; but, as the results have been published in Tables, the finding the Trigonometrical Ratios does not form any part of a student's work, except to exemplify the method employed.

89. The general method of finding Trigonometrical Ratios belongs to a more advanced part of the subject than the present, but there are certain angles whose Ratios can be found in a simple manner.

90. *To find the sine, cosine and tangent of an angle of* 45°.

If one angle of a right-angled triangle be 45°, that is, the half of a right angle, the third angle must also be 45°. Hence 45° is one angle of an isosceles right-angled triangle.



Let POM be an isosceles triangle such that PMO is a right angle, and $OM = MP$. Then $POM = OPM = 45°$.

Let the measures of OM and of MP each be m. Let the measure of OP be x.

Then $$x^2 = m^2 + m^2 = 2m^2;$$

$$\therefore\ x = \sqrt{2}\,.\,m.$$

Hence, $$\sin 45° = \sin POM = \frac{MP}{OP} = \frac{m}{\sqrt{2}\,.\,m} = \frac{1}{\sqrt{2}},$$

$$\cos 45° = \cos POM = \frac{OM}{OP} = \frac{m}{\sqrt{2}\,.\,m} = \frac{1}{\sqrt{2}},$$

$$\tan 45° = \tan POM = \frac{MP}{OM} = \frac{m}{m} = \frac{1}{1} = 1.$$

91. *To find the sine, cosine and tangent of* 60°.

In an equilateral triangle, each of the equal angles is 60°, because they are each one third of 180°. And if we draw a perpendicular from one of the angular points of the triangle to the opposite side, we get a right-angled triangle in which one angle is 60°.

Let OPQ be an equilateral triangle. Draw PM perpendicular to OQ. Then OQ is bisected in M.

Let the measure of OM be m; then that of OQ is $2m$ and therefore that of OP is $2m$.



Let the measure of MP be x.

Then $x^2 = (2m)^2 - m^2 = 4m^2 - m^2 = 3m^2$,

$\therefore\ x = \sqrt{3}\,.\,m.$

Hence, $\sin 60^\circ = \sin POM = \dfrac{MP}{OP} = \dfrac{\sqrt{3}\,.\,m}{2m} = \dfrac{\sqrt{3}}{2}$,

$\cos 60^\circ = \cos POM = \dfrac{OM}{OP} = \dfrac{m}{2m} = \dfrac{1}{2}$,

$\tan 60^\circ = \tan POM = \dfrac{MP}{OM} = \dfrac{\sqrt{3}\,.\,m}{m} = \dfrac{\sqrt{3}}{1} = \sqrt{3}.$

92. *To find the sine, cosine and tangent of* 30°.

With the same figure and construction as above, we have the angle $OPM = 30^\circ$, since it is a half of OPQ, i.e. of 60°.

Hence, $\sin 30^\circ = \sin OPM = \dfrac{MO}{PO} = \dfrac{m}{2m} = \dfrac{1}{2}$,

$\cos 30^\circ = \cos OPM = \dfrac{PM}{PO} = \dfrac{\sqrt{3}\,.\,m}{2m} = \dfrac{\sqrt{3}}{2}$,

$\tan 30^\circ = \tan OPM = \dfrac{MO}{PM} = \dfrac{m}{\sqrt{3}\,.\,m} = \dfrac{1}{\sqrt{3}}$.

**93. *To find the sine of* 18°.

In the 10th Prop. of Euclid IV. a triangle is described such that each of the angles at the base is double of the third angle. [See also Example (6) p. 140.]



Let POQ be such a triangle, and let the vertical angle POQ contain n degrees; then

$$n + 2n + 2n = 180,$$

$$n = 36.$$

Draw OM perpendicular to PQ, bisecting the angle POQ. Then since $QOP = 36°$, $MOP = 18°$. Also $PM = MQ$.

Let the measure of MP be m, and the measure of OP be x. From OP cut off $OR = PQ$. Then by Euclid IV. 10 $PO \,.\, PR = PQ^2$.

$$\therefore\ x(x - 2m) = (2m)^2,$$

$$\therefore\ x^2 - 2mx + m^2 = 4m^2 + m^2 = 5m^2,$$

$$\therefore\ x - m = \sqrt{5} \,.\, m,$$

$$\therefore\ x = \sqrt{5} \,.\, m + m,$$

$$\therefore \sin 18° = \sin MOP = \frac{MP}{OP} = \frac{m}{\sqrt{5} \,.\, m + m} = \frac{1}{\sqrt{5} + 1} = \frac{\sqrt{5} - 1}{4}.$$

94. *To find the sine, cosine and tangent of* 0°.

By this is meant,—*To find the values, if any, to which the Trigonometrical Ratios of a very small angle approach, as the angle is continually diminished.*

Let ROP be a small angle. Draw PM perpendicular to OR, and let OP be always of the same length, so that P lies on a circle whose centre is O.

Then if the angle ROP be diminished, we can see that MP is diminished also, and that consequently $\frac{MP}{OP}$, which is sin ROP, is diminished. And, by diminishing the angle ROP sufficiently, *we can make* MP *as small as we please,* and therefore *we can make* sin ROP *smaller than any assignable number however small that number may be.*

Thus we see that the value to which sin ROP approaches as the angle is diminished, is 0.

This is expressed by saying, sin $0° = 0$ ..... i.

Again, as the angle ROP diminishes, OM approaches OP in length; and cos ROP, which is $\frac{OM}{OP}$, approaches in value to $\frac{OP}{OP}$, i.e. to 1.

This is expressed by saying, cos $0° = 1$ ..... ii.

Also, tan ROP is $\frac{MP}{OM}$; and we have seen that MP approaches 0, while OM does not; $\therefore$ tan ROP approaches 0.

This is expressed by saying, tan $0° = 0$ ..... iii.

95. *To find the sine, cosine and tangent of* 90°.

By this is meant,—*To find the values, if any, to which the Trigonometrical Ratios of an angle approach, as the angle approaches a right angle.*

Let ROU be a right angle = 90°.



Draw ROP nearly a right angle; draw PM perpendicular to OR, and let OP be always of the same length, so that P lies on a circle whose centre is O.

Then, as the angle ROP approaches to ROU, we can see that MP approaches OP, while OM continually diminishes.

Hence when ROP approaches 90°, sin ROP, which is $\frac{MP}{OP}$, approaches in value to $\frac{OP}{OP}$, that is to $\frac{1}{1}$, i.e. to 1.

Hence we say, that sin 90° = 1 ..... i.

Again, when ROP approaches 90°, cos ROP, which is $\frac{OM}{OP}$, approaches in value to $\frac{0}{OP}$, that is to 0.

Hence we say, that cos 90° = 0 ..... ii.

Again, when ROP approaches 90^0, $\tan ROP$ which is $\frac{MP}{OM}$ approaches in value to $\frac{OP}{\text{a quantity which approaches } 0}$.

But in any fraction whose numerator does not diminish, the smaller the denominator the greater is the value of that fraction; and if the denominator continually diminishes the value of the fraction continually increases.

Hence, *tan ROP can be made larger than any assignable number by making the angle ROP approach* 90^0 *near enough.*

This is what we mean when we say, that

$\tan 90^0$ is **infinity,** or, $\tan 90^0 = \infty$ ..... iii.

96. The following table exhibits the results of this Chapter.

angle	0^0	18^0	30^0	45^0	60^0	90^0
sine	0	$\frac{\sqrt{5}-1}{4}$	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
cosine	1		$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
tangent	0		$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	∞

The student may notice that the sine increases with the angle, while the cosine diminishes as the angle increases.

Also that the squares of the sines of 0^0, 30^0, 45^0, 60^0 and 90^0 are respectively 0, $\frac{1}{4}$, $\frac{2}{4}$, $\frac{3}{4}$ and $\frac{4}{4}$, and that the squares of the cosines of the same angles are $\frac{4}{4}$, $\frac{3}{4}$, $\frac{2}{4}$, $\frac{1}{4}$, and 0.

EXAMPLES. XIII.

If $A=90^0$, $B=60^0$, $C=30^0$, $D=45^0$, $E=18^0$, prove the following :

(1) $2 . \sin D . \cos D = \sin A$. (2) $2 . \sin C . \cos C = \sin B$.

(3) $\cos^2 B - \sin^2 B = 1 - 2\sin^2 B$.

(4) $\sin B . \cos C + \sin C . \cos B = \sin A$.

(5) $\cos^2 D - \sin^2 D = \cos A$. (6) $4 . \sin^2 E + 2 . \sin E = 1$.

(7) $\sin^2 B + \cos^2 B = 1$. (8) $\cos^2 C + \sin^2 C = 1$.

(9) $\cos^2 D + \sin^2 D = 1$.

(10) $\sin B . \cos C - \sin C . \cos B = \sin C$.

(11) $2(\cos B . \cos D + \sin B . \sin D)^2 = 1 + \cos C$.

(12) $2(\sin D . \cos C - \sin C . \cos D)^2 = 1 - \cos C$.

(13) $\sin 30^0 = \cdot 5$. (14) $\sin 45^0 = \cdot 7071\ldots$.

(15) $\sin 60^0 = \cdot 8660\ldots$. (16) $\tan 60^0 = 1 \cdot 7320508\ldots$.

(17) $\tan 30^0 = \cdot 5773\ldots$. (18) $\sin 18^0 = \cdot 3090\ldots$.

97. The actual measurement of the *line* joining two points which are any considerable distance apart, is a very tedious and difficult operation, especially when great accuracy is required; while the accurate measurement of an *angle* can, with proper instruments, be made with comparative ease and quickness.

98. A **Sextant** is an instrument for measuring the angle between the two lines drawn from the observer's eye to each of two distant objects respectively.

A **Theodolite** is an instrument for measuring angles in a horizontal plane; also for measuring '*angles of elevation*' and '*angles of depression.*'

99. The angle made with the horizontal plane, by the line joining the observer's eye with a distant object, is called

(i) its **angle of elevation**, when the object is *above* the observer;

(ii) its **angle of depression**, when the object is *below* the observer. †

100. **Trigonometry** enables us by measuring certain *angles*, to deduce, from one known distance, the *lengths* of other distances: or, by the measurement of a *convenient* line, to deduce by the measurement of *angles* the lengths of lines whose actual measurement is difficult or impossible.

[In the Trigonometrical Survey of England, made by the Ordnance Department, the only distance actually measured was one of about seven miles on Salisbury Plain.]

101. For this purpose we require the numerical values of the Trigonometrical Ratios of the angles observed. Accordingly mathematical tables have been compiled, containing *lists* of the values of these Ratios. These Tables constitute a kind of numerical **Dictionary**, in which we can find the numerical value of the Trigonometrical Ratios of any required angle.

Example 1. At a point 100 feet from the foot of a tower the angle of elevation of the top of the tower is observed to be 60^0. Find the height above the point of observation of the top of the tower.



† In measuring the angle of *depression* the telescope is turned from a horizontal position *downwards*. See Ex. (3) p. 63.

Let O be the point of observation; let P be the top of the tower; let a horizontal line through O meet the foot of the tower at the point M. Then $OM=100$ feet, and the angle $MOP=60^0$. Let MP contain x feet.

Then $$\frac{MP}{OM}=\tan MOP=\tan 60^0=\sqrt{3}$$

$$\therefore \frac{x}{100}=\sqrt{3}.$$

$$\therefore x=100.\sqrt{3}=100\times 1{\cdot}7320 \text{ etc.}$$

$$=173{\cdot}2.$$

Therefore the required height is 173·2 feet.

Example 2. A flagstaff, 25 feet high, stands on the top of a cliff, and from a point on the seashore the angles of elevation of the highest and lowest points of the flagstaff are observed to be $47^0\ 12'$ and $45^0\ 13'$ respectively. Find the height of the cliff.



Let O be the point of observation, PQ the flagstaff.

Let a horizontal line through O meet the vertical line PQ produced in M.

Then $QP=25$ feet, $MOP=47^0\ 12'$, $MOQ=45^0\ 13'$.

Let $MQ=x$ feet; let $OM=y$ feet.

Then $\frac{MP}{OM} = \tan 47^0\ 12'$, $\therefore \frac{x+25}{y} = \tan 47^0\ 12'$,

and $\frac{MQ}{OM} = \tan 45^0\ 13'$, $\therefore \frac{x}{y} = \tan 45^0\ 13'$.

Hence, by division $\therefore \frac{x+25}{x} = \frac{\tan 47^0\ 12'}{\tan 45^0\ 13'}$.

In the Tables we find that

$$\tan 47^0\ 12' = 1{\cdot}0799018, \text{ and } \tan 45^0\ 13' = 1{\cdot}0075918,$$

$$\therefore 1 + \frac{25}{x} = \frac{1{\cdot}0799018}{1{\cdot}0075918} = 1 + \frac{{\cdot}0723100}{1{\cdot}0075918},$$

$$\therefore \frac{x}{25} = \frac{1{\cdot}0075918}{{\cdot}0723100} = \frac{100759}{7231}.$$

$$\therefore x = \frac{2518975}{7231} = 348 \text{ nearly.}$$

Therefore the cliff is 348 feet high.

EXAMPLES. XIV.

The answers are given correct to three significant figures.

(1) At a point 179 feet in a horizontal line from the foot of a column, the angle of elevation of the top of the column is observed to be 45^0. What is the height of the column?

(2) At a point 200 feet from, and on a level with the base of a tower, the angle of elevation of the top of the tower is observed to be 60^0: what is the height of the tower?

(3) From the top of a vertical cliff, the angle of depression of a point on the shore 150 feet from the base of the cliff, is observed to be 30^0: find the height of the cliff.

(4) From the top of a tower 117 feet high the angle of depression of the top of a house 37 feet high is observed to be 30^0: how far is the top of the house from the tower?

(5) A man 6 ft. high stands at a distance of 4 ft. 9 in. from a lamp-post, and it is observed that his shadow is 19 ft. long. Find the height of the lamp.

(6) The shadow of a tower in the sunlight is observed to be 100 ft. long, and at the same time the shadow of a lamp-post 9 ft. high is observed to be $3\sqrt{3}$ ft. long. Find the angle of elevation of the sun, and the height of the tower.

(7) From a point P on the bank of a river, just opposite a post Q on the other bank, a man walks at right angles to PQ to a point R so that PR is 100 yards; he then observes the angle PRQ to be $32^0\ 17'$: find the breadth of the river.

$$\tan 32^0\ 17' = \cdot 6317667.$$

(8) A flagstaff 25 feet high stands on the top of a house; from a point on the plain on which the house stands the angles of elevation of the top and bottom of the flagstaff are observed to be 60^0 and 45^0 respectively: find the height of the house above the point of observation.

(9) From the top of a cliff 100 feet high, the angles of depression of two ships at sea are observed to be 45^0 and 30^0 respectively; if the line joining the ships points directly to the foot of the cliff, find the distance between the ships.

(10) A tower 100 feet high stands on the top of a cliff; from a point on the sand at the foot of the cliff the angles of elevation of the top and bottom of the tower are observed to be 75^0 and 60^0 respectively; find the height of the cliff. (Tan $75^0 = 2 + \sqrt{3}$).

(11) A man walking along a straight road observes at one milestone a house in a direction making an angle 30^0 with the road, and that at the next milestone the angle is 60^0: how far is the house from the road?

(12) A man stands at a point A on the bank AB of a straight river and observes that the line joining A to a post C on the opposite bank makes with AB an angle of 30^0. He then goes 400 yards along the bank to B and finds that BC makes with BA an angle of 60^0; find the breadth of the river.

**(13) A building on a square base $ABCD$ has two of its sides, AB and CD, parallel to the bank of a river. An observer, standing at E on the other side of the river so that DAE is a straight line, finds that AB subtends at his eye an angle of 45^0. Having walked a yards parallel to the bank, he finds that DE subtends an angle whose tangent is $\sqrt{2}$. Show that $DB = a$ yards.

(14) From the top of a hill the angles of depression of the top and bottom of a flagstaff 25 feet high at the foot of the hill are observed to be $45^0\ 13'$ and $47^0\ 12'$ respectively; find the height of the hill.

$$\tan 45^0\ 13' = 1{\cdot}0075918,$$
$$\tan 47^0\ 12' = 1{\cdot}0799018.$$

(15) From each of two stations, East and West of each other, the altitude of a balloon is observed to be 45^0, and its bearings to be respectively N.W. and N.E.: if the stations be 1 mile apart, determine the height of the balloon.

(16) The angle of elevation of a balloon from a station due south of it is 60^0; and from another station due west of the former and distant a mile from it it is 45^0. Find the height of the balloon.

(17) An isosceles triangle of wood is placed on the ground in a vertical position facing the sun. If $2a$ be the base of the triangle, b its height, and 30^0 the altitude of the sun, find the tangent of half the angle at the apex of the shadow.

(18) The length of the shadow of a vertical stick is to the length of the stick as $\sqrt{3} : 1$. If the stick be turned about its lower extremity in a vertical plane, so that the shadow is always in the same direction, find what will be the angle of its inclination to the horizon when the length of the shadow is the same as before.

(19) What distance in space is travelled in an hour in consequence of the earth's rotation, by a person situated in latitude 60^0? (Earth's radius = 4000 miles.)

CHAPTER VII.

On the Relations between the Trigonometrical Ratios of the Same Angle.

102. The following relations are evident from the definitions:

$$\operatorname{cosec}\theta = \frac{1}{\sin\theta}, \quad \sec\theta = \frac{1}{\cos\theta}, \quad \cot\theta = \frac{1}{\tan\theta}.$$

103. To prove $\tan\theta = \dfrac{\sin\theta}{\cos\theta}$.



We have $\sin\theta = \dfrac{\text{perpendicular}}{\text{hypotenuse}}$,

and $\cos\theta = \dfrac{\text{base}}{\text{hypotenuse}}$;

$$\therefore \frac{\sin\theta}{\cos\theta} = \frac{\text{perpendicular}}{\text{base}} = \tan\theta.$$

104. We may prove similarly $\cot\theta = \dfrac{\cos\theta}{\sin\theta}$.

Or thus, $\cot\theta = \dfrac{1}{\tan\theta} = \dfrac{\cos\theta}{\sin\theta}$.

* 105. Euclid I. 47 gives us that in any right-angled triangle the square on the hypotenuse = the sum of the squares on the perpendicular and on the base,

or, $(\text{hypotenuse})^2 = (\text{perpendicular})^2 + (\text{base})^2$.

(i) Divide each side of this identity by $(\text{hypotenuse})^2$, and we get

$$\left(\frac{\text{hypotenuse}}{\text{hypotenuse}}\right)^2 = \left(\frac{\text{perpendicular}}{\text{hypotenuse}}\right)^2 + \left(\frac{\text{base}}{\text{hypotenuse}}\right)^2,$$

that is, $1 = \sin^2\theta + \cos^2\theta$.

(ii) Divide each side of the same identity by $(\text{base})^2$, and we get

$$\left(\frac{\text{hypotenuse}}{\text{base}}\right)^2 = \left(\frac{\text{perpendicular}}{\text{base}}\right)^2 + \left(\frac{\text{base}}{\text{base}}\right)^2,$$

that is, $\sec^2\theta = \tan^2\theta + 1$.

(iii) Divide each side of the same identity by $(\text{perpendicular})^2$, and we get

$$\left(\frac{\text{hypotenuse}}{\text{perpendicular}}\right)^2 = \left(\frac{\text{perpendicular}}{\text{perpendicular}}\right)^2 + \left(\frac{\text{base}}{\text{perpendicular}}\right)^2,$$

that is, $\operatorname{cosec}^2\theta = \cot^2\theta + 1$.

* 106. Thus the three results

$$\left.\begin{array}{ll} \text{(i)} & \cos^2\theta + \sin^2\theta = 1 \\ \text{(ii)} & 1 + \tan^2\theta = \sec^2\theta \\ \text{(iii)} & 1 + \cot^2\theta = \operatorname{cosec}^2\theta \end{array}\right\},$$

are each a statement in Trigonometrical language of Euc. I. 47.

107. We give the above proof in a different form.

To prove that $\cos^2\theta + \sin^2\theta = 1$.

Let ROE be any angle θ.



In OE take any point P, and draw PM perpendicular to OR. Then with respect to θ, MP is the perpendicular, OP is the hypotenuse, and OM is the base;

$$\therefore \sin^2\theta = \frac{MP^2}{OP^2}, \quad \cos^2\theta = \frac{OM^2}{OP^2}$$

We have to prove that $\sin^2\theta + \cos^2\theta = 1$.

Now

$$\cos^2\theta + \sin^2\theta = \frac{MP^2}{OP^2} + \frac{OM^2}{OP^2}$$

$$= \frac{MP^2 + OM^2}{OP^2} = \frac{OP^2}{OP^2} = 1,$$

since by Euclid I. 47, $MP^2 + OM^2 = OP^2$,

Therefore $\cos^2\theta + \sin^2\theta = 1$.

Similarly we may prove that

$$1 + \tan^2\theta = \sec^2\theta,$$

and that

$$1 + \cot^2\theta = \operatorname{cosec}^2\theta.$$

108. The following is a LIST OF FORMULÆ with which the student must make himself familiar:

$$\operatorname{cosec}\theta = \frac{1}{\sin\theta},$$

$$\sec\theta = \frac{1}{\cos\theta},$$

$$\cot\theta = \frac{1}{\tan\theta},$$

$$\tan\theta = \frac{\sin\theta}{\cos\theta},$$

$$\cot\theta = \frac{\cos\theta}{\sin\theta},$$

$$\sin^2\theta + \cos^2\theta = 1,$$

$$\tan^2\theta + 1 = \sec^2\theta,$$

$$\cot^2\theta + 1 = \operatorname{cosec}^2\theta.$$

109. In proving Trigonometrical identities it is often convenient to express the other Trigonometrical Ratios *in terms of the sine and cosine.*

Example. *Prove that* tan A + *cot* A = *sec* A . *cosec* A.

Since $$\tan A = \frac{\sin A}{\cos A}, \quad \cot A = \frac{\cos A}{\sin A},$$

$$\sec A = \frac{1}{\cos A} \text{ and } \operatorname{cosec} A = \frac{1}{\sin A},$$

we have to prove that

$$\frac{\sin A}{\cos A} + \frac{\cos A}{\sin A} = \frac{1}{\cos A} \cdot \frac{1}{\sin A},$$

or that

$$\frac{\sin^2 A + \cos^2 A}{\cos A \,.\, \sin A} = \frac{1}{\cos A \,.\, \sin A},$$

and this is true, because $\sin^2 A + \cos^2 A = 1$.

110. Sometimes it is more convenient to express all the other Trigonometrical Ratios in terms of the *sine* only, or in terms of the *cosine* only.

Example (i). *Prove that*

$$\sin^4\theta + 2\sin^2\theta\cos^2\theta = 1 - \cos^4\theta.$$

Here the right-hand expression does not contain $\sin\theta$; hence for $\sin^2\theta$ in the left-hand expression we substitute $(1-\cos^2\theta)$; thus

$$\begin{aligned}\sin^4\theta + 2\sin^2\theta\cos^2\theta &= (\sin^2\theta)^2 + 2(\sin^2\theta)\cos^2\theta \\ &= (1-\cos^2\theta)^2 + 2(1-\cos^2\theta)\cos^2\theta \\ &= 1 - 2\cos^2\theta + \cos^4\theta + 2\cos^2\theta - 2\cos^4\theta \\ &= 1 - \cos^4\theta. \quad \text{Q.E.D.}\end{aligned}$$

Example (ii). *Prove that*

$$\sin^6\theta + \cos^6\theta = 1 - 3\cos^2\theta + 3\cos^4\theta.$$

Here

$$\begin{aligned}\sin^6\theta + \cos^6\theta &= (\sin^2\theta + \cos^2\theta)(\sin^4\theta - \sin^2\theta\cos^2\theta + \cos^4\theta) \\ &= 1 \times (\sin^4\theta - \sin^2\theta\cos^2\theta + \cos^4\theta) \\ &= (\sin^2\theta)^2 - (\sin^2\theta)\cos^2\theta + \cos^4\theta \\ &= (1-\cos^2\theta)^2 - (1-\cos^2\theta)\cos^2\theta + \cos^4\theta \\ &= 1 - 2\cos^2\theta + \cos^4\theta - \cos^2\theta + \cos^4\theta + \cos^4\theta \\ &= 1 - 3\cos^2\theta + 3\cos^4\theta. \quad \text{Q.E.D.}\end{aligned}$$

NOTE. $(1-\cos\theta)$ is called the **versed sine** of θ; it is abbreviated thus $\operatorname{versin}\theta$.

EXAMPLES. XV.

Prove the following statements.

(1) $\cos A \,.\, \tan A = \sin A$, (2) $\cot A \,.\, \tan A = 1$.

(3) $\cos A = \sin A \,.\, \cot A$. (4) $\sec A \,.\, \cot A = \operatorname{cosec} A$.

(5) $\operatorname{cosec} A \,.\, \tan A = \sec A$.

(6) $(\tan A + \cot A) \sin A \,.\, \cos A = 1$.

(7) $(\tan A - \cot A) \sin A \,.\, \cos A = \sin^2 A - \cos^2 A$.

(8) $\cos^2 A - \sin^2 A = 2\cos^2 A - 1 = 1 - 2\sin^2 A$.

(9) $(\sin A + \cos A)^2 = 1 + 2 \sin A \,.\, \cos A$.

(10) $(\sin A - \cos A)^2 = 1 - 2 \sin A \,.\, \cos A$.

(11) $\cos^4 B - \sin^4 B = 2\cos^2 B - 1$.

(12) $(\sin^2 B + \cos^2 B)^2 = 1$.

(13) $(\sin^2 B - \cos^2 B)^2 = 1 - 4\cos^2 B + 4\cos^4 B$.

(14) $1 - \tan^4 B = 2\sec^2 B - \sec^4 B$.

(15) $(\sec B - \tan B)(\sec B + \tan B) = 1$.

(16) $(\operatorname{cosec} \theta - \cot \theta)(\operatorname{cosec} \theta + \cot \theta) = 1$.

(17) $\sin^3 \theta + \cos^3 \theta = (\sin \theta + \cos \theta)(1 - \sin \theta \cos \theta)$.

(18) $\cos^3 \theta - \sin^3 \theta = (\cos \theta - \sin \theta)(1 + \sin \theta \cos \theta)$.

(19) $\sin^6 \theta + \cos^6 \theta = 1 - 3\sin^2 \theta \,.\, \cos^2 \theta$.

(20) $(\sin^6 \theta - \cos^6 \theta) = (2\sin^2 \theta - 1)(1 - \sin^2 \theta + \sin^4 \theta)$.

(21) $\dfrac{\tan A + \tan B}{\cot A + \cot B} = \tan A \,.\, \tan B$.

(22) $\dfrac{\cot \alpha + \tan \beta}{\tan \alpha + \cot \beta} = \cot \alpha \,.\, \tan \beta$.

(23) $\dfrac{1 - \sin A}{1 + \sin A} = (\sec A - \tan A)^2$.

(24) $\dfrac{1 + \cos A}{1 - \cos A} = (\operatorname{cosec} A + \cot A)^2$.

(25) $2 \operatorname{versin} \theta - \operatorname{versin}^2 \theta = \sin^2 \theta$.

(26) $\operatorname{versin} \theta (1 + \cos \theta) = \sin^2 \theta$.

111. All the Trigonometrical Ratios of an angle can be expressed in terms of any one of them.

Example 1. *To express all the trigonometrical ratios of an angle in terms of the sine.*



Let ROE be any angle A.

We can take P anywhere in the line OE; so that we can make *one* of the lines, OP, OM, or MP any length we please.

Let us take OP so that its measure is 1, and let s be the measure of MP; so that $\sin A$, which is $\frac{MP}{OP}$, $=\frac{s}{1}$; or, $s=\sin A$.

Let x be the measure of OM.

Then since $$OM^2=OP^2-MP^2,$$

$$\therefore\ x^2=1-s^2,$$

$$\therefore\ x=\sqrt{1-s^2},$$

Hence $$\cos A=\frac{OM}{OP}=\frac{\sqrt{1-s^2}}{1}=\sqrt{1-\sin^2 A},$$

$$\tan A=\frac{MP}{OM}=\frac{s}{\sqrt{1-s^2}}=\frac{\sin A}{\sqrt{1-\sin^2 A}},$$

and so on.

NOTE. The solution of the equation $x^2=1-s^2$, gives

$$x=\pm\sqrt{1-s^2},$$

and therefore the ambiguity ($\pm$) must stand before each of the root symbols in the above. This ambiguity, as will be explained later on, is of great use when the magnitude of the angle is not limited. When we limit A to be less than a right angle we have no use for the negative sign.

Example 2. To express all the other trigonometrical ratios of an angle in terms of the tangent.



In this case $\tan\theta = \frac{MP}{OM}$.

Take P so that the measure of OM is 1, and let t be the measure of MP; so that $\tan\theta$, which is $\frac{MP}{OM}$, $= \frac{t}{1}$; or, $t = \tan\theta$.

Then we can show that the measure of OP is $\sqrt{1+t^2}$.

Hence, $$\sin\theta = \frac{MP}{OP} = \frac{t}{\sqrt{1+t^2}} = \frac{\tan\theta}{\sqrt{1+\tan^2\theta}}.$$

$$\cos\theta = \frac{OM}{OP} = \frac{1}{\sqrt{1+t^2}} = \frac{1}{\sqrt{1+\tan^2\theta}}.$$

and so on.

112. The same results may be obtained by the use of the formulæ on page 69.

Example $$\cos^2\theta + \sin^2\theta = 1,$$

$$\therefore \cos^2\theta = 1 - \sin^2\theta,$$

$$\therefore \cos\theta = \sqrt{1-\sin^2\theta}.$$

Again $$\tan\theta = \frac{\sin\theta}{\cos\theta} = \frac{\sin\theta}{\sqrt{1-\sin^2\theta}},$$

and so on.

EXAMPLES. XVI.

(1) Express all the other Trigonometrical Ratios of A in terms of $\cos A$.

(2) Express all the other Trigonometrical Ratios of A in terms of $\cot A$.

(3) Express all the other Trigonometrical Ratios of A in terms of $\sec A$.

(4) Express all the other Trigonometrical Ratios of A in terms of $\operatorname{cosec} A$.

(5) Use the formulæ of page 69 to express all the other Trigonometrical Ratios of A in terms of $\sin A$.

(6) Use the formulæ of page 69 to express all the other Trigonometrical Ratios of A in terms of the $\tan A$.

113. *Given one of the Trigonometrical Ratios of an angle less than a right angle, we can find all the others.*

Since all the Trigonometrical Ratios of an angle can be expressed in terms of any one of them, it is clear that if the numerical value of any one of them be given, the numerical value of all the rest can be found.

Example. Given $\sin\theta=\frac{3}{4}$, find the other Trigonometrical Ratios of θ.

Let ROE be the angle θ.

Take P on OE so that the measure of OP is 4. Draw PM perpendicular to OR.



Then since $\sin\theta=\frac{3}{4}$ $\left(\text{so that } \frac{MP}{OP}=\frac{3}{4}\right)$, and since the measure of OP is 4, therefore the measure of MP must be 3.

Let x be the measure of OM;

then $$OM^2=OP^2-MP^2,$$

$$\therefore\ x^2=4^2-3^2=16-9=7.$$

$$x=\sqrt{7}.$$

Therefore the measure of OM is $\sqrt{7}$.

Hence, $$\cos\theta=\frac{OM}{OP}=\frac{\sqrt{7}}{4},$$

$$\tan\theta=\frac{MP}{OM}=\frac{3}{\sqrt{7}}=\frac{3\sqrt{7}}{7}.$$

$$\cot\theta=\frac{\sqrt{7}}{3}.$$

EXAMPLES. XVII.

(1) Given that $\sin A=\frac{3}{5}$, find $\tan A$ and $\operatorname{cosec} A$.

(2) Given that $\cos B=\frac{1}{3}$, find $\sin B$ and $\cot B$.

(3) Given that $\tan A=\frac{4}{3}$, find $\sin A$ and $\sec A$.

(4) Given that $\sec\theta=4$, find $\cot\theta$ and $\sin\theta$.

(5) Given that $\tan\theta=\sqrt{3}$, find $\sin\theta$ and $\cos\theta$.

(6) Given that $\cot\theta=\frac{2}{\sqrt{5}}$, find $\sin\theta$ and $\sec\theta$.

(7) Given that $\sin\theta=\frac{b}{c}$, find $\tan\theta$.

(8) Given that $\tan\theta=\frac{a}{b}$, find $\sin\theta$ and $\cos\theta$.

(9) Given that $\cos\theta=\frac{1}{a}$, find $\sin\theta$ and $\cot\theta$.

(10) If $\sin\theta=a$, and $\tan\theta=b$, prove that $(1-a^2)(1+b^2)=1$.

(11) If $\cos\theta=h$, and $\tan\theta=k$, find the equation connecting h and k.

114. The following propositions are important.

PROP. I. *To trace the changes in the magnitude of* sin A *as* A *increases from* 0^0 *to* 90^0.



Take a line OR, of any length; and describe the quadrant RPU of the circle whose centre is O and radius OR.

Draw the right angle ROU, cutting the circle in U.

Let OP make any angle $ROP\ (=A)$ with OR; draw PM perpendicular to OR.

Then $$\sin A = \frac{MP}{OP}.$$

When the angle A is 0^0, MP is zero, and when A is 90^0, MP is equal to OP; and as A continuously increases from 0^0 to 90^0, MP increases continuously from zero to OP; also OP is always equal to OR.

Therefore, when $A=0^0$, the fraction $\frac{MP}{OP}$ is equal to $\frac{0}{OP}$, that is 0; when $A=90^0$ the fraction $\frac{MP}{OP}$ is equal to $\frac{OP}{OP}$, that is 1; and as A continuously increases from 0^0 to 90^0, the *numerator* of the fraction $\frac{MP}{OP}$ *continuously increases* from zero to OP, while the denominator is unchanged, and therefore the *fraction* $\frac{MP}{OP}$, which is sin A, *increases continuously* from 0 to 1.

PROP. II. *To trace the changes in the magnitude of* tan A *as* A *increases from* 0^0 *to* 90^0.

With the same construction and figure as in the last article, we have

$$\tan A = \frac{MP}{OM}.$$

When the angle A is 0^0, MP is zero; when A is 90^0, MP is equal to OP; and as the angle continuously increases from 0^0 to 90^0, MP increases continuously from zero to OP.

When the angle A is 0^0, OM is equal to OP; when A is 90^0, OM is zero; and as A continuously increases from 0^0 to 90^0, OM continuously decreases from OP to zero.

Hence, when A is 0^0, the fraction $\frac{MP}{OM}$ is equal to $\frac{0}{OM}$, that is 0; when A is 90^0, the fraction $\frac{MP}{OM}$ is equal to $\frac{OP}{0}$, that is 'infinity' (see art. 95); and as A continuously increases from 0^0 to 90^0, the numerator continuously increases from zero to OP, while the denominator continuously diminishes from OP to zero; so that the fraction $\frac{MP}{OM}$, which is $\tan A$, *continuously* increases from 0 until it is *greater than any assignable numerical quantity.*

EXAMPLES. XVIII.

(1) Show that as A continuously increases from 0^0 to 90^0, $\cos A$ continuously diminishes from 1 to 0.

(2) Trace the changes in the magnitude of $\sec\theta$ as θ increases from 0 to $\frac{\pi}{2}$.

(3) Trace the changes in the magnitude of $\sin A$ as A diminishes from 90^0 to 0^0.

(4) Trace the changes in the magnitude of $\cot\theta$ as θ increases from 0 to $\frac{\pi}{2}$.

115. Since the hypotenuse is the *greatest* side in a right-angled triangle, it is clear

(i) that *sin A*, which is $\frac{\text{perpendicular}}{\text{hypotenuse}}$, is never greater than unity,

(ii) that *cos A*, which is $\frac{\text{base}}{\text{hypotenuse}}$, is never greater than unity,

(iii) that *cosec A*, which is $\frac{\text{hypotenuse}}{\text{perpendicular}}$, is never *numerically* less than unity (we shall see later on that it may be *negative*),

(iv) that *sec A*, which is $\frac{\text{hypotenuse}}{\text{base}}$, is never numerically less than unity (it may be negative).

Hence such a question as '*Find an angle whose* **sine** *is* $\frac{4}{3}$' has no solution, since there is no angle whose sine is greater than unity.

And such a question as '*Find an angle whose* **secant** *is* $\frac{3}{4}$' has no solution, because there is no angle whose secant is numerically less than unity.

116. If A be small, the perpendicular is smaller than the base; and tan A, which is $\frac{\text{perpendicular}}{\text{base}}$, can be made as small as we please (see figure in Art. 114). Also if the angle A be nearly a right angle, the perpendicular is greater than the base; and tan A, which is $\frac{\text{perpendicular}}{\text{base}}$, can be made as large as we please.

So that an angle whose **tangent** is a can *always* be found if a be positive; that is, there is always an angle A between 0° and 90°, such that $\tan A = a$, if a be a positive number.

[The student will find as he proceeds that the equation $\tan x = a$ can always be solved if a be any *arithmetical* quantity].

Example 1. *To draw an angle* A *whose sine is* $\frac{3}{4}$.

Draw any line OR, and take R such that the measure of OR is unity; describe the quadrant RPU of the circle whose centre is O and radius OR.

In OU which is perpendicular to OR, take ON so that the measure of ON is $\frac{3}{4}$.



Draw NP parallel to OR cutting the quadrant in P. (The student should observe that if the measure of ON were *greater than unity*, N would be outside the circle altogether, and the line parallel to OR would not cut the circle at all.)

Join OP, and draw PM perpendicular to OR.

Then ROP is the angle required.

For $\sin ROP = \frac{MP}{OP} = \frac{ON}{OP} = \frac{3}{4} \div 1 = \frac{3}{4}$.

Therefore an angle POR has been drawn whose sine is $\frac{3}{4}$.

Example 2. *To draw an angle whose cosine is* b; where b is a *proper* fraction and positive.



Draw any line OR, and take OR so that its measure is unity.

Describe the quadrant RPU of the circle whose centre is O and radius OR.

Take M in OR so that the measure of OM is b. OM will be less than OR because b is a *proper* fraction.

Draw MP perpendicular to OR cutting the quadrant in P. (The student will observe that if b were an *improper* fraction, M would be outside the circle altogether.)

Then the angle ROP is the angle required.

For $\cos ROP = \frac{OM}{OP} = \frac{b}{1} = b$.

Therefore an angle ROP has been described such that the cos $ROP = b$.

Example 3. *To draw an angle* A *such that* $\tan A = a$, where a is any positive numerical quantity.



Draw any line OR, and take M in it so that the measure of OM is unity.

Draw MP perpendicular to OM, and take P so that the measure of MP is a. Join OP. Then the angle ROP is the angle required.

For $\tan ROP = \frac{MP}{OM} = \frac{a}{1} = a$.

Thus an angle ROP has been drawn so that $\tan ROP = a$.

117. Definition. One angle is called the **complement** of another, when the two angles added together make up a *right angle.*

Example 1. The complement of A is $(90^0 - A)$.

Example 2. The complement of 190^0 is $(90^0 - 190^0) = -100^0$.
For $190^0 + (90^0 - 190^0) = 90^0$.

Example 3. The complement of $\frac{5\pi}{4}$ is $\left(\frac{\pi}{2} - \frac{5\pi}{4}\right) = -\frac{3\pi}{4}$.

118. *To prove that the sine of an angle* A *is equal to the cosine of its complement* $(90^0 - \text{A})$.

[This is true whatever be the magnitude of A, as will be proved later on.]

If A be less than 90^0, let ROP be A (see last figure).

Draw PM perpendicular to OR.

Then since $PMO = 90^0$, therefore $POM + OPM = 90^0$ and therefore $OPM = (90^0 - A)$.

Now, $\sin A = \frac{MP}{OP} = \cos OPM = \cos (90^0 - A)$. Q.E.D.

EXAMPLES. XIX.

(1) Draw an angle whose sine is $\frac{1}{2}$.

(2) Draw an angle whose cosecant is 2.

(3) Draw an angle whose tangent is 2.

(4) Can an angle be drawn whose tangent is 431?

(5) Can an angle be drawn whose cosine is $\frac{4}{3}$?

(6) Can an angle be drawn whose secant is 5?

(7) Find the complements of

(i) 30^0. (ii) 190^0. (iii) 90^0. (iv) 350^0.

(v) -25^0. (vi) -320^0. (vii) $\frac{3\pi}{4}$. (viii) $-\frac{\pi}{6}$.

(8) Prove that $\sin 70^0 = \cos 20^0$.

(9) Prove that $\cos 47^0\ 16' = \sin 42^0\ 44'$.

(10) Prove that $\tan 79^0 = \cot 11^0$.

(11) Prove that $\sec 36^0 = \operatorname{cosec} 54^0$.

(12) If A be less than 90^0, prove

(i) $\cos A = \sin (90^0 - A)$.
(ii) $\tan A = \cot (90^0 - A)$.
(iii) $\sec A = \operatorname{cosec} (90^0 - A)$.
(iv) $\cot A = \tan (90^0 - A)$.

On the Solution of Trigonometrical Equations.

119. A Trigonometrical equation is an equation in which there is a letter, such as θ, which stands for an *angle* of unknown magnitude.

The *solution* of the equation is the process of finding an angle which, if it be substituted for θ, satisfies the equation.

Example 1. *Solve* $\cos\theta=\frac{1}{2}$.

This is a Trigonometrical equation. To solve it we must find some angle such that its cosine is $\frac{1}{2}$.

We know that $\cos 60^0=\frac{1}{2}$.

Therefore if 60^0 be put for θ the equation is satisfied.

$\therefore\ \theta=60^0$ is a solution of the equation.

Example 2. *Solve* $\sin\theta-\operatorname{cosec}\theta+\frac{3}{2}=0$.

The usual method of solution is to express all the Trigonometrical Ratios in terms of one of them.

Thus we put $\dfrac{1}{\sin\theta}$ for cosec θ, and we get

$$\sin\theta-\frac{1}{\sin\theta}+\frac{3}{2}=0.$$

This is an equation in which θ, and therefore *sin* θ is unknown. It will be convenient if we put x for *sin* θ, and then solve the equation for x as an ordinary algebraical equation. Thus we get

$$x-\frac{1}{x}+\frac{3}{2}=0,$$

or,
$$x^2+\frac{3x}{2}=1,$$

$$\therefore\ x^2+\frac{3x}{2}+\frac{9}{16}=1+\frac{9}{16}=\frac{25}{16}.$$

$$\therefore\ x+\frac{3}{4}=\pm\frac{5}{4}.$$

Whence
$$x=-2,\text{ or }x=\frac{1}{2}.$$

But x stands for $\sin\theta$.

Thus we get
$$\sin\theta=-2,\text{ or }\sin\theta=\frac{1}{2}.$$

The value -2 is inadmissible for $\sin\theta$, for there is no angle whose sine is numerically greater than 1 (Art. 115).

$$\therefore\ \sin\theta=\frac{1}{2}.$$

But
$$\sin 30^0=\frac{1}{2}.$$

$$\therefore\ \sin\theta=\sin 30^0.$$

Therefore *one* angle which satisfies this equation for θ is 30^0.

Example 3. To solve the equation $\operatorname{cosec}\theta - \cot^2\theta + 1 = 0$.

We have $1 + \cot^2\theta = \operatorname{cosec}^2\theta$. [Art. 108]

$\therefore \cot^2\theta = \operatorname{cosec}^2\theta - 1$.

and we get $\operatorname{cosec}\theta - \operatorname{cosec}^2\theta + 2 = 0$.

Let x stand for *cosec* θ. Then

$$x - x^2 + 2 = 0.$$

$$\therefore x^2 - x = +2.$$

or $$\therefore x^2 - x + \tfrac{1}{4} = +\tfrac{1}{4} + 2 = \tfrac{9}{4}.$$

$$\therefore x - \tfrac{1}{2} = \pm\tfrac{3}{2}.$$

Whence $$x = 2, \text{ or } x = -1,$$

$$\therefore \operatorname{cosec}\theta = 2,$$

but $$\operatorname{cosec} 30^0 = 2,$$

$$\therefore \operatorname{cosec}\theta = \operatorname{cosec} 30^0.$$

therefore 30^0 is *one* angle which satisfies the equation for θ.

EXAMPLES. XX.

Find one angle which satisfies each of the following equations.

(1) $\sin\theta = \dfrac{1}{\sqrt{2}}$.

(2) $4\sin\theta = \operatorname{cosec}\theta$.

(3) $2\cos\theta = \sec\theta$.

(4) $4\sin\theta - 3\operatorname{cosec}\theta = 0$.

(5) $4\cos\theta - 3\sec\theta = 0$.

(6) $3\tan\theta = \cot\theta$.

(7) $3\sin\theta - 2\cos^2\theta = 0$.

(8) $\sqrt{2}\sin\theta = \tan\theta$.

(9) $2\cos\theta = \sqrt{3}\cot\theta$.

(10) $\tan\theta = 3\cot\theta$.

(11) $\tan\theta + 3\cot\theta = 4$.

(12) $\tan\theta + \cot\theta = 2$.

(13) $2\sin^2\theta + \sqrt{2}\cos\theta = 2$.

(14) $4\sin^2\theta + 2\sin\theta = 1$.

(15) $3\tan^2\theta - 4\sin^2\theta = 1$.

(16) $2\sin^2\theta + \sqrt{2}\sin\theta = 2$.

(17) $\cos^2\theta - \sqrt{3}\cos\theta + \tfrac{3}{4} = 0$.

(18) $\cos^2\theta + 2\sin^2\theta - \tfrac{5}{2}\sin\theta = 0$.

* * MISCELLANEOUS EXAMPLES. XXI.

(1) Prove that $3 \sin 60^0 - 4 \sin^3 60^0 = 4 \cos^3 30^0 - 3 \cos 30^0$.

(2) Prove that $\tan 30^0 (1 + \cos 30^0 + \cos 60^0) = \sin 30^0 + \sin 60^0$.

(3) If $2 \cos^2 \theta - 7 \cos \theta + 3 = 0$, show there is only one value of $\cos \theta$.

(4) Find $\cos \theta$ from the equation $8 \cos^2 \theta - 8 \cos \theta + 1 = 0$.

(5) Find $\sin \theta$ from the equation $8 \sin^2 \theta - 10 \sin \theta + 3 = 0$, and prove that one value of θ is $\frac{\pi}{6}$.

(6) Find $\tan \theta$ from the equation $12 \tan^2 \theta - 13 \tan \theta + 3 = 0$.

(7) If $3 \cos^2 \theta + 2 . \sqrt{3} . \cos \theta = 5\frac{1}{4}$, show that there is only one value of $\cos \theta$, and that one value of θ is $\frac{\pi}{6}$.

(8) Prove that the value of $\sin^4 \theta + \cos^4 \theta + 2 . \sin^2 \theta . \cos^2 \theta$ is always the same.

(9) Simplify $\cos^4 A + 2 . \sin^2 A . \cos^2 A$.

(10) Express $\sin^6 A + \cos^6 A$ in terms of $\sin^2 A$ and powers of $\sin^2 A$.

(11) Express $1 + \tan^4 \theta$ in terms of $\cos \theta$ and its powers.

(12) Prove that $\frac{\cos A + \cos B}{\sin A - \sin B} + \frac{\sin A + \sin B}{\cos A - \cos B} = 0$.

(13) Express $(\sec A - \tan A)^2$ in terms of $\sin A$.

(14) Trace the changes in the magnitude of $\operatorname{cosec} \theta$ as θ increases from 0 to $\frac{\pi}{2}$.

(15) Trace the changes in the magnitude of $\cot \theta$ as θ decreases from $\frac{\pi}{2}$ to 0.

(16) Solve $\sin (\theta + \phi) = \frac{\sqrt{3}}{2}$, $\cos (\theta - \phi) = \frac{\sqrt{3}}{2}$.

CHAPTER VIII.

On the Use of the Signs + and −.

120. The student is probably aware that, in the application of Algebra to Problems concerning *distance*, we sometimes find that the solution of an equation gives the measure of a *distance* with the sign − before it.

Example. Let M, N, O be places in a straight line; let the distance from M to N be 3 miles, and the distance from N to O, 3 miles.

One man A starting from M, rides towards O at the rate of 10 miles an hour, while another man B starting simultaneously from N, walks towards O at the rate of 4 miles an hour;

If Q be the point at which they meet, *how far is Q beyond O?*

Let P be any point beyond O, and let x be the number of miles in OP. We wish to find x, i.e. the measure of OP, so that P may coincide with Q, the point at which A overtakes B.

When A arrives at P, he has ridden $6+x$ miles. The time occupied at the rate of 10 miles an hour is $\frac{6+x}{10}$ hours.

When B arrives at P, he has walked $3+x$ miles. The time occupied at the rate of 4 miles an hour is $\frac{3+x}{4}$ hours.

When P is the point at which they meet, these times are equal, so that

$$\frac{6+x}{10}=\frac{3+x}{4}; \quad \text{whence } x=-1.$$

Thus the required *number of miles* has the sign − before it; and we have failed to find a point *beyond* O at which A overtakes B.

121. Such a result can generally be *interpreted* by altering the statement of the problem, thus:

The line whose measure appears with the sign − before it, must be considered as drawn in the **direction opposite** to that in which it was drawn in the first statement of the problem.



Example. Taking the former example, let us alter the question as follows:

If Q be the point at which A overtakes B, how far is Q **to the left** of O?

Let P be any point *to the left* of O, and let x be the number of miles in OP.

We wish to find x (i.e. the measure of OP), so that P may coincide with Q, the point at which A overtakes B.

When A arrives at P, he has ridden $6-x$ miles.

When B arrives at P, he has walked $3-x$ miles.

Proceeding as before, we get

$$\frac{6-x}{10}=\frac{3-x}{4}; \quad \text{whence } x=+1.$$

Therefore if P is to coincide with Q (the point at which A overtakes B), OP must be one mile *to the left* of O.

122. The consideration of such examples as the above has suggested, that the sign − may be made use of, in the application of Algebra to Geometry, to represent a **direction** exactly opposite to that represented by the sign +.

Accordingly the following Rule, or Convention, has been made.

RULE. Any straight line AB being given, then

lines drawn parallel to AB in **one direction** shall be **positive**; that is, shall be *represented algebraically* by their *measures with the sign + before them:*

lines drawn parallel to BA in the **opposite direction** shall be **negative**; that is, shall be *represented algebraically* by their *measures with the sign − before them.*

123. We may choose for the positive direction in each case that direction which is most convenient.

Example. Let LR be a straight line parallel to the printed lines in the page,



and let lines drawn in the direction from L to R in the figure that is, from the left-hand towards the right, be considered *positive.* Then by the above rule, lines drawn in the direction from R to L, that is, from right to left, must be *negative.*

124. In naming a line by the letters at its extremities, we can indicate by the **order of the letters**, the direction in which the line is supposed to be drawn.

Example. Let O and P be two points in the line LR as in the figure, and let the measure of the distance between them be a.

Then OP, i.e. the line drawn from O to P, which is in the *positive* direction, is *represented algebraically* by $+a$.

While PO, i.e. the line drawn from P to O, which is in the *negative* direction, is *represented algebraically* by $-a$.

125. Hence in using the two letters at its extremities to represent a line, the student will find it advantageous always to pay careful attention to the **order of the letters.**

Example. Let LR be a straight line parallel to the printed lines in the page.

Let A, B, C, D, E be points in LR, such that the measures of AB, BC, CD, DE, are 1, 2, 3, 4 respectively.

Find the algebraical representation of

(i) $AC+CB$ (ii) $AD+DC-BC$.

(i) The algebraical representation of AC is $+3$,

the algebraical representation of CB is -2.

Hence that of $AC+CB$ is $+3-2$; that is, $+1$*.

(ii) The algebraical representation of AD is $+6$, that of DC is -3, and that of BC is $+2$.

Therefore that of $AD+DC-BC=6-3-2=+1$.

This is equivalent to that of AB.

EXAMPLES. XXII.

In the above figure, find the algebraical representation of

(1) $AB+BC+CD$. (2) $AB+BC+CA$.

(3) $BC+CD+DE+EC$. (4) $AD-CD$.

(5) $AD+DB+BE$. (6) $BC-AC+AD-BD$.

(7) $CD+DB+BE$. (8) $CD-BD+BA+AC+CE$.

* By $AC+CB$ (attention being paid to *direction*), we mean 'Go from A to C and from C to B.' The result is equivalent to starting from A and stopping at B, *i.e.* equivalent to AB.

CHAPTER IX.

On the Use of the Signs + and − in Trigonometry.

126. In **Trigonometry** in order conveniently to treat of angles of *any* magnitude, we proceed as follows.

We take a fixed point O, called the **origin**; and a fixed straight line OR, called the **initial line.**

The angle of which we wish to treat is described by a line OP, called the **revolving line.** This line OP *starts from* the initial line OR, and turns about O through an angle ROP of any proposed magnitude into the position OP.



127. We have already said in Art. 41

(i) that, when an angle ROP is described by OP turning about O in the direction *contrary to that of the hands of a watch,* the angle ROP is said to be **positive**; that is, *is represented algebraically* by its *measure with the sign + before it.*

(ii) that, when an angle ROP is described by OP turning about O in the *same* direction as the hands of a watch, the angle is said to be **negative**; that is, is *represented algebraically* by its *measure with the sign − before it.*

Example. $(180^0 - A)$ indicates

(i) the angle described by OP turning about O from the position OR in the positive direction until it has described an angle of $(180 - A)$ degrees.



Or, (ii) the angle described by OP turning about O, from the position OR, in the positive direction until it has described an angle of 180^0 (when it has turned into the position OL), and then turning back from OL in the negative direction through the angle $-A$ into the position OP.

Or, (iii) the angle described by OP turning about O from the position OR, in the negative direction through the angle $-A$, and then turning back in the positive direction through the angle 180^0, into the position OP.

The student should observe that in each of these three ways of regarding the angle $(180^0 - A)$, the *resulting* angle ROP is the same.

EXAMPLES. XXIII.

Draw a figure giving the position of the revolving line after it has turned through each of the following angles.

(1) 270^0. (2) 370^0. (3) 425^0. (4) 590^0.

(5) -30^0. (6) -330^0. (7) -480^0. (8) -750^0.

(9) $\frac{27\pi}{4}$. (10) $2n\pi + \frac{\pi}{6}$. (11) $(2n+1)\pi + \frac{\pi}{3}$.

(12) $(2n+1)\pi - \frac{\pi}{4}$. (13) $2n\pi - \frac{\pi}{2}$. (14) $(2n+1)\pi - \frac{\pi}{2}$.

Note. $n\pi$ always stands for a whole number of two right angles.

128. It is often convenient to keep the revolving line of the same length.



In this case the point P lies always on the circumference of a circle whose centre is O.

Let this circle cut the lines LOR, UOD in the points L, R, U, D respectively.

The circle $RULD$ is thus divided at the points R, U, L, D into four **Quadrants**, of which

RU is called the **first Quadrant.**

UL is called the **second Quadrant.**

LD is called the **third Quadrant.**

DR is called the **fourth Quadrant.**

Hence, in the figure,

ROP_1 is an angle of the first Quadrant.

ROP_2 „ „ second Quadrant.

ROP_3 „ „ third Quadrant.

ROP_4 „ „ fourth Quadrant.

129. When we are told that an angle is of some particular Quadrant, say the third, we know that the position in which the revolving line *stops* is in the third Quadrant. But there is an unlimited number of *angles* having this same final position of OP.

Example. 25^0; 385^0 i.e. 360^0+25^0; 745^0 i.e. $2\times360^0+25^0$; -335 i.e. -360^0+25^0 are each an angle of the first Quadrant, and are all represented *geometrically* by the *same* final position of OP.

130. Let A be an angle between 0^0 and 90^0, and let n be any *whole* number, positive or negative.

Then

(i) $2n\times180^0+A$ represents algebraically an angle of the *first* Quadrant.

(ii) $2n\times180^0-A$ represents algebraically an angle of the *fourth* Quadrant.

[For $2n\times180^0$ represents some number n of *complete* revolutions of OP; so that after describing $n\times360^0$, OP is again in the position OR.]

(iii) $(2n+1)\times180^0-A$ represents algebraically an angle of the *second* Quadrant.

(iv) $(2n+1)\times180^0+A$ represents algebraically an angle of the *third* Quadrant.

[For after describing $(2n+1)\times180^0$, OP is in the position OL.]

The corresponding expressions in circular measure are

(i) $2n\,\pi+\theta$; (ii) $2n\,\pi-\theta$; (iii) $(2n+1)\,\pi-\theta$;

(iv) $(2n+1)\,\pi+\theta$.

EXAMPLES. XXIV.

State in which Quadrant the revolving line will be after describing the following angles :

(1) 120^0. (2) 340^0. (3) 490^0.

(4) -100^0. (5) -380^0. (6) -1000^0.

(7) $\frac{2\pi}{3}$. (8) $10\pi+\frac{\pi}{4}$. (9) $9\pi-\frac{3\pi}{4}$.

(10) $2n\pi-\frac{\pi}{4}$. (11) $(2n+1)\pi+\frac{2\pi}{3}$. (12) $n\pi+\frac{\pi}{6}$.

131. The principal **directions of lines** with which we are concerned in **Trigonometry** are as follows ;

i. that parallel to the initial line OR (OR is usually drawn from O towards the right hand, parallel to the printed lines in the page ; and RO is produced to L.)

ii. that parallel to the line DOU, which is drawn through O at right angles to LOR ;

iii. that parallel to the revolving line OP.



Accordingly we make the following **rules:**

I. Any line drawn parallel to LR in the direction **from left to right** is to be **positive**; and *consequently* (Art. 122) any line drawn parallel to RL in the *opposite* direction, i.e. from **right to left**, is to be **negative.**

II. Any line drawn parallel to DU in the direction from D to U, **upwards**, is to be **positive**; and consequently any line drawn parallel to UD in the *opposite* direction, i.e. **downwards**, is to be **negative.**

III. Any line drawn parallel to the revolving line in the direction from O to P is to be **positive**, and consequently any line drawn in the direction from P to O is to be **negative.**

Note. The student must notice that the revolving line OP *carries its positive direction round with it*, so that the line 'OP' is *always positive*. The revolving line, as it turns about O, nowhere undergoes any *sudden* change of direction such as would be indicated by a change of sign.

132. We said, in Art. 81, that **the definitions of the Trigonometrical Ratios** (on pp. 46, 47), **apply to angles of any magnitude.** We have only to remark that it is generally convenient to take P on the revolving line; that PM is drawn perpendicular to the other line *produced if necessary;* and that the **order of the letters** in MP, OP, OM is an essential part of the definition.

The **order of the letters** P, M, O in the expressions $\frac{MP}{OP}$, etc., is therefore of great importance.

Minus to left; Plus to the right
Negative depth; Positive height

133. We proceed to show that the Trigonometrical Ratios of an angle vary in **Sign** according to the **Quadrant** in which the revolving line of the angle happens to be.

From the definition we have, with the usual letters,

$$\sin ROP = \frac{MP}{OP}, \cos ROP = \frac{OM}{OP}, \tan ROP = \frac{MP}{OM}.$$

Fig.1.



Fig.11.



Fig.111.



Fig.1V.



I. When OP is in the **first** Quadrant (Fig. I.).

MP is *positive* because from M to P is *upwards*

(Rule II. p. 95.)

OM is *positive* because from O to M is *towards the right*

(Rule I.)

OP is *positive*

(Rule III.)

Hence, if A be any angle of the *first* Quadrant,

$$\sin A, \text{ which is } \frac{MP}{OP}, \text{ is } \textit{positive};$$

$$\cos A, \text{ which is } \frac{OM}{OP}, \text{ is } \textit{positive};$$

$$\tan A, \text{ which is } \frac{MP}{OM}, \text{ is } \textit{positive}.$$

II. When OP is in the **second** Quadrant, (Fig. II.).

MP is *positive*, because from M to P is *upwards*,

OM is *negative*, because from O to M is *towards the left*.

OP is *positive*.

Hence, if A be any angle of the *second* Quadrant,

$$\sin A, \text{ which is } \frac{MP}{OP}, \text{ is } \textit{positive};$$

$$\cos A, \text{ which is } \frac{OM}{OP}, \text{ is } \textit{negative};$$

$$\tan A, \text{ which is } \frac{MP}{OM}, \text{ is } \textit{negative}.$$

III. When OP is in the **third** Quadrant (Fig. III.)

MP is *negative*, OM is *negative*, OP is *positive*.

So that, if A be any angle of the *third* Quadrant,

$\sin A$ is *negative*, $\cos A$ is *negative*, $\tan A$ is *positive*.

IV. When OP is in the **fourth** Quadrant (Fig. IV.)

MP is *negative*, OM is *positive*, OP is *positive*.

So that, if A be any angle of the *fourth* Quadrant,

$\sin A$ is *negative*, $\cos A$ is *positive*, $\tan A$ is *negative*.

134. The table given below exhibits the results of the last article.

Quadrant ...	I.	II.	III.	IV.
Sine	+	+	−	−
Cosine	+	−	−	+
Tangent ...	+	−	+	−

The student should notice that in any particular Quadrant the three *signs* of sine, cosine, and tangent are unlike their signs in any other Quadrant.

EXAMPLES. XXV.

State the *sign* of the sine, cosine, and tangent of each of the following angles :

(1) 60^0. (2) 135^0. (3) 265^0.

(4) 275^0. (5) -10^0. (6) -91^0.

(7) -193^0. (8) -350^0. (9) -1000^0.

(10) $2n\pi+\frac{\pi}{4}$. (11) $2n\pi+\frac{3\pi}{4}$. (12) $2n\pi-\frac{\pi}{6}$.

135. The NUMERICAL VALUES through which the Trigonometrical Ratios of the angle ROP pass, as OP moves through the *first* Quadrant, are **repeated** as OP moves through each of the other Quadrants.

Thus as OP moves through the second Quadrant from U to L, Fig. II, (OP being always of the same length) MP and OM pass through the same succession of numerical values through

which they pass, as P moves through the first Quadrant in the *opposite* direction from U to R.

Example 1. *Find the sine, cosine and tangent of* 120^0.

120^0 is an angle of the second Quadrant.

Let the angle ROP be 120^0 (Fig. II. p. 96.)

Then the angle $POL = 180^0 - 120^0 = 60^0$.

Hence, $\sin 120^0 = \frac{MP}{OP} = \sin 60^0$ *numerically*, and in the *second* Quadrant the sine is *positive*.

Therefore $\sin 120^0 = \frac{\sqrt{3}}{2}$ ..... (i).

Again, $\cos 120^0 = \frac{OM}{OP} = \cos 60^0$, *numerically*, and in the *second* Quadrant the cosine is *negative*.

Therefore $\cos 120^0 = -\frac{1}{2}$ ..... (ii).

Similarly, $\tan 120^0 = -\sqrt{3}$ ..... (iii).

Example 2. *Find the sine, cosine and tangent of* 225^0.

225^0 is an angle in the third Quadrant

Let the angle ROP be 225^0 (Fig. III.)

Here the angle $POL = 225^0 - 180^0 = 45^0$.

Therefore the Trigonometrical Ratios of 225^0 = those of 45^0 *numerically ;* and in the *third* Quadrant the sine and cosine are each *negative* and the tangent is *positive*.

Hence, $\sin 225^0 = -\frac{1}{\sqrt{2}}$; $\cos 225^0 = -\frac{1}{\sqrt{2}}$; $\tan 225^0 = 1$.

136. The cosecant, secant and cotangent of an angle A have the same *sign* as the sine, cosine, and tangent of A respectively.

EXAMPLES. XXVI.

Find the algebraical value of the sine, cosine and tangent of the following angles :

(1) 150^0.	(2) 135^0.	(3) -240^0.	(4) 330^0.
(5) -45^0.	(6) -300^0.	(7) 225^0.	(8) -135^0.
(9) 390^0.	(10) 750^0.	(11) -840^0.	(12) 1020^0.

(13) $2n\pi+\frac{\pi}{4}$. (14) $(2n+1)\pi-\frac{\pi}{3}$. (15) $(2n-1)\pi+\frac{\pi}{6}$.

* *Example. To trace the changes in the magnitude and sign of sin* A, *as* A *increases from* 0^0 *to* 360^0.

Take the figure and construction of page 96.

As A increases from 0^0 to 90^0, MP increases from zero to OP, and is positive.

Therefore sin A increases from 0 to 1 and is positive.

As A increases from 90^0 to 180^0, MP decreases from OP to zero, and is positive.

Therefore sin A decreases from 1 to 0 and is positive.

As A increases from 180^0 to 270^0, MP increases from zero to OP, and is negative.

Therefore sin A increases numerically from 0 to 1 and is negative.

As A increases from 270^0 to 360^0, MP decreases from OP to zero, and is negative.

Therefore sin A decreases numerically from 1 to 0 and is negative.

* EXAMPLES. XXVII.

Trace the changes in sign and magnitude as A increases from 0^0 to 360^0 of

(1) cos A.	(2) tan A.	(3) cot A.	(4) sec A.
(5) cosec A.	(6) $1-\sin A$.	(7) $\sin^2 A$.	(8) $\sin A \cdot \cos A$.
(9) $\sin A+\cos A$.	(10) $\tan A+\cot A$.	(11) $\sin A-\cos A$.	

CHAPTER X**.

On Angles Unlimited in Magnitude.

137. Just as the definitions of the Trigonometrical Ratios apply to angles of any magnitude whatever, so **every general Formula** involving these Ratios **is true for angles of any magnitude whatever.**

It is most important that the student should examine for himself into the truth of this statement.

138. The formulæ

$$\operatorname{cosec} A = \frac{1}{\sin A}, \quad \sec A = \frac{1}{\cos A}, \quad \cot A = \frac{1}{\tan A},$$

are really definitions; and since the definitions apply, therefore these formulæ are true, whatever be the magnitude of A.

The formulæ $\tan A = \dfrac{\sin A}{\cos A}$, $\quad \cot A = \dfrac{\cos A}{\sin A}$,

are deduced immediately from the definitions, and therefore they are true whatever be the magnitude of A.

139. The formulæ $\sin^2 A + \cos^2 A = 1$,

$$1 + \tan^2 A = \sec^2 A,$$

$$1 + \cot^2 A = \operatorname{cosec}^2 A,$$

are each a trigonometrical statement of Euc. I. 47, and depend only on the fact that MP, OM, and OP are the sides of a *right-angled* triangle. That this is the case, whatever be the magnitude of the angle A, is evident from the figures on page 96.

** To be omitted on a first reading, except pp. 104, 105.

140. In Art. 118 we proved that the sine of an angle is equal to the cosine of its complement, provided the angle lies between 0^0 and 90^0. We now give some examples of a method of proving the truth of this and other like formulæ, **whatever** be the magnitude of the angle concerned.

Example 1. *To prove that the sine of an angle = the cosine of its complement.*

That is, *to prove* $\sin A = \cos(90^0 - A)$

and $\cos A = \sin(90^0 - A)$.

We take **two revolving lines** OP and OP'. OP starting from OR is to describe the angle A; OP' starting from OR is to describe $(90^0 - A)$.

As usual, PM, $P'M'$ are perpendiculars on OR and $P'N'$ is a perpendicular on OU.

In describing $(90^0 - A)$ we shall consider that OP' starting from OR, turns first through 90^0 into the position OU, and then turns back from OU through the angle $UOP' = (-A)$.



So that ROP, the angle which OP describes from OR, is always equal to UOP', the angle which OP' describes from OU in the *opposite* direction.

Hence, $N'P'$, that is OM', is *always* equal to MP in *magnitude.*

Also it will be seen that when P is *above* LOR, P' is to the *right* of UOD; when P is *below* LOR, P' is to the *left* of UOD.

Hence, OM' and MP have *always* the *same sign.*

Therefore $$\frac{MP}{OP}=\frac{OM'}{OP'} \text{ always,}$$

or, $$\sin A=\cos(90^0-A), \text{ for all values of } A.$$

Again, ON', that is $M'P'$, is *always* equal to OM in *magnitude.*

And P' is above or below LOR according as P is to the right or to the left of UOD.

So that $M'P'$ and OM have *always* the *same sign.*

Therefore $$\frac{OM}{OP}=\frac{M'P'}{OP'} \text{ always,}$$

or, $$\cos A=\sin(90^0-A) \text{ for all values of } A.$$

EXAMPLES. XXVIII.

Prove, drawing a separate figure for each example, that

(1) $\sin 30^0=\cos 60^0$. (2) $\sin 65^0=\cos 25^0$.

(3) $\sin 195^0=\cos(-105^0)$. (4) $\cos 275=\sin(-185^0)$.

(5) $\cos(-27^0)=\sin 117^0$. (6) $\cos 300^0=\sin(-210^0)$.

If A, B, C be the angles of a triangle, so that $A+B+C=180^0$, prove

(7) $\cos\frac{A}{2}=\sin\frac{B+C}{2}$. (8) $\cos\frac{B}{2}=\sin\frac{A+C}{2}$.

(9) $\sin\frac{C}{2}=\cos\frac{A+B}{2}$. (10) $\sin\frac{A}{2}=\cos\frac{B+C}{2}$.

141. **Def.** Two angles are said to be the **Supplements** the one of the other when their sum is two right angles.

Thus $(180^0 - A)$ is the **supplement** of A

If A, B, C be the angles of a triangle, $(A+B+C)=180^0$, so that $(B+C)$ is the **supplement** of A.

Example 2. *To prove that the sine of an angle=the sine of its supplement; and that the cosine of an angle= −(the cosine of its supplement).*

That is, *to prove* $\sin A = \sin(180^0 - A)$

and $$\cos A = -\cos(180^0 - A).$$

We take **two revolving lines** OP and OP'. OP starting from OR describes the angle A; OP' starting from OR describes the angle $(180^0 - A)$.

In describing $(180^0 - A)$ we consider that OP' starting from OR turns first through 180^0 into the position OL, and then back from OL through the angle $LOP' = (-A)$.



So that ROP, the angle which OP describes from OR, is always equal to LOP', the angle which OP' describes from OL in the *opposite* direction.

Hence, MP and $M'P'$ are *always* equal in *magnitude.*

Also, P and P' are always *both* above, or *both* below LOR.

So that MP and $M'P'$ are *always* of the *same sign.*

Therefore $$\frac{MP}{OP} = \frac{M'P'}{OP'} \text{ always,}$$

or, $$\sin A = \sin (180^0 - A), \text{ for all values of } A.$$

Again, OM and OM' are *always* equal in *magnitude.*

Also it will be seen that when P is on the *right* of UOD, P' is on the *left* of UOD; when P is on the *left* of UOD, P' is on *right* of UOD.

So that OM and OM' are *always* of *opposite sign.*

Therefore $$\frac{OM}{OP} = -\frac{OM'}{OP'} \text{ always,}$$

or, $$\cos A = -\cos (180^0 - A), \text{ for all values of } A.$$

EXAMPLES. XXIX.

Prove, drawing a separate figure in each case, that

(1) $\sin 60^0 = \sin 120^0$. (2) $\sin 340^0 = \sin (-160^0)$.

(3) $\sin (-40^0) = \sin 220^0$. (4) $\cos 320^0 = -\cos (-140^0)$

(5) $\cos (-380^0) = -\cos 560^0$. (6) $\cos 195^0 = -\cos (-15^0)$.

If A, B, C be the angles of a triangle, prove

(1) $\sin A = \sin (B + C)$. (2) $\sin C = \sin (A + B)$.

(3) $\cos B = -\cos (A + C)$ (4) $\cos A = -\cos (C + B)$.

Example 3. *To prove* $\sin A = -\sin(180^0 + A)$,

and $\cos A = -\cos(180^0 + A)$.

As before, we take **two revolving lines** OP and OP'. OP starting from OR describes the angle A; OP' starting from OR describes the angle $(180^0 + A)$.

In describing $(180^0 + A)$ we consider that OP' starting from OR turns first through 180^0 into the position OL, and then on from OL through the angle A.



So that ROP, the angle which OP describes from OR, is always equal to LOP', the angle which OP' describes from OL in the *same* direction.

Hence, $M'P'$ *always* $= MP$ in *magnitude*.

Also it will be seen that when P is *above* LOR, P' is *below* LOR; and *vice versâ**.

So that MP and $M'P'$ are *always* of *opposite sign*

Hence, $$\frac{MP}{OP} = -\frac{M'P'}{OP'} \text{ always,}$$

or, $$\sin A = -\sin(180^0 + A), \text{ for all values of } A.$$

Similarly, OM *always* $= OM'$ in *magnitude*.

And P' is to the left or to the right of UOD according as P is to the right or to the left of UOD.

* This will be more clear if the student observes that POP' is always a straight line.

Hence, $$\frac{OM}{OP} = -\frac{OM'}{OP'} \text{ always,}$$

or, $$\cos A = -\cos(180^0 + A), \text{ for all values of } A.$$

Example 4. *To prove* $\sin A = -\cos(90^0 + A)$,

and $$\cos A = \sin(90^0 + A).$$

As before, we take **two revolving lines** OP and OP'. OP starting from OR describes the angle A; OP' starting from OR describes the angle $90^0 + A$.

In describing $(90^0 + A)$ we consider that OP' starting from OR turns first through 90^0 into the position OU, and then on from OU through the angle A.



So that ROP, the angle which OP describes from OR, is always equal to UOP', the angle which OP' describes from OU in the *same* direction.

Hence, $N'P'$, that is OM', *always* $= MP$ in *magnitude*.

Also, P' is to the left or to the right of UOD according as P is above or below LOR.

So that MP and OM' are always of *opposite sign*.

Hence, $$\frac{MP}{OP} = -\frac{OM'}{OP'} \text{ always,}$$

or. $$\sin A = -\cos(90^0 + A), \text{ for all values of } A.$$

Similarly, OM *always* $= M'P'$ in *magnitude.*

And P' is above or below LOR according as P is to the right or to the left of UOD.

So that OM and $M'P'$ are *always* of the *same sign.*

Hence, $$\frac{OM}{OP} = \frac{M'P'}{OP} \text{ always,}$$

or, $$\cos A = \sin (90^0 + A), \text{ for all values of } A.$$

MISCELLANEOUS EXAMPLES. XXX.

Prove, drawing a separate figure in each case, that

(1) $\sin 60^0 = -\sin 240^0$.

(2) $\sin 170^0 = -\sin 350^0$.

(3) $\sin(-20^0) = -\sin 160^0$.

(4) $\cos 380^0 = -\cos 560^0$.

(5) $\cos(-225) = -\cos(-45^0)$.

(6) $\cos 1005^0 = -\cos 1185^0$

(7) $\sin 60^0 = -\cos 150^0$.

(8) $\cos 60^0 = \sin 150^0$

(9) $\sin 225 = -\cos 315^0$.

(10) $\cos(-60^0) = \sin 30^0$.

If $A+B+C$ be the angles of a triangle, prove that

(11) $\sin A = -\sin(2A+B+C)$.

(12) $\sin A = -\cos \dfrac{3A+B+C}{2}$.

(13) $\cos B = \sin \dfrac{A+3B+C}{2}$.

(14) $\cos C = -\cos(A+B+2C)$.

(15) $\cos \dfrac{B-C}{2} = \sin \dfrac{A+2B}{2}$.

(16) $\sin \dfrac{C-A}{2} = -\cos \dfrac{B+2C}{2}$.

Prove the following statements for all values of A and of a.

(17) $\sin A = -\sin(-A)$. (18) $\cos A = \cos(-A)$.

(19) $\sin A = \cos(A - 90^0)$. (20) $\cos A = -\sin(A - 90^0)$.

(21) $\sin a = \cos\left(\frac{3\pi}{2} + a\right)$.

(22) $\cos a = -\sin\left(\frac{3\pi}{2} + a\right)$

(23) $\sin a = -\cos\left(\frac{3\pi}{2} - a\right)$.

(24) $\cos a = -\sin\left(\frac{3\pi}{2} - a\right)$.

(25) $\sin\left(\frac{\pi}{2} - a\right) = \sin\left(\frac{\pi}{2} + a\right)$.

(26) $\cos(\pi + a) = \cos(\pi - a)$.

(27) $\tan(90^0 - A) = \cot A$. (28) $\tan A = -\tan(-A)$.

(29) $\tan(90^0 + A) = -\cot(A)$. (30) $\tan a = -\tan(\pi - a)$.

(31) $\tan A = \tan(180^0 + A)$. (32) $\cot\left(\frac{\pi}{2} - a\right) = \tan a$.

142. We have seen (Art. 135) that there are many angles of different magnitude which have the same *sine*.

If two such angles are in the *same Quadrant* they are represented geometrically by the same position of OP, so that they differ by some multiple of four right angles.

143. If we are given the value of the sine of an angle it is important to be able to find, Geometrically and Algebraically, all angles which have that value for their sine.

144. *To find the complete Geometrical Solution of the equation* $\sin\theta = a$.

With the usual construction, let the radius of the circle $RULD$ be the unit of length; then the measure of OP is 1.





From O draw on OU a line ON so that its measure is a.

[ON will be drawn upwards (Fig. I.) or downwards (Fig. II.) from O according as a is positive or negative.]

Through N draw P_1P_2 parallel to LOR to cut the circle in P_1 and P_2*. Join OP_1, OP_2. Draw P_1M_1, P_2M_2 perpendicular to LOR.

Then $M_1P_1 = ON = M_2P_2$, and $\dfrac{M_1P_1}{OP_1} = \dfrac{a}{1} = \dfrac{M_2P_2}{OP_2}$.

Hence any angle described by the revolving line OP, when OP, starting from OR, stops either in the position OP_1, or in the position OP_2, is an angle whose **sine** is a.

And no other angle has its sine equal to a, for there is no other possible position of N.

145. *To find the complete Algebraical Solution of the equation* $\sin\theta = a$.

* If a were *greater than unity*, ON would be greater than OU, and the construction would fail.

With the figure and construction of the last article; let α be the least angle whose sine $= a$.

Then in the figure, $ROP_1 = \alpha$ and $P_2OL = \alpha$.

Every angle whose sine $= a$ is included among those described by the revolving line OP when, starting from OR, OP stops in one or the other of the positions OP_1 or OP_2.

When OP, starting from OR, stops in the position OP_1, the angle may be described by OP turning through a certain number of complete revolutions to OR, and then through ROP_1; that is, the angle is one of those included in

$$2m\pi + \alpha,$$

where m is some integer, positive or negative. [Art. 130.]

When OP, starting from OR, stops in the position OP_2, the angle may be described by OP turning through a certain odd number of half-revolutions to OL, and then *back* through LOP_2; that is, the angle is one of those included in

$$(2r+1)\,\pi - \alpha,$$

where r is some integer, positive or negative.

Both of these expressions are included in

$$n\pi + (-1)^n \alpha,$$

where n is some integer, positive or negative †.

Thus, the solution of the equation $\sin\theta = \sin\alpha$ is

$$\theta = n\pi + (-1)^n \alpha.$$

Example. Find six angles between -4 right angles and $+8$ right angles which satisfy the equation $\sin A = \sin 18^0$.

We have $\theta = n\pi + (-1)^n \frac{\pi}{10}$, or $A = n \times 180^0 + (-1)^n 18^0$.

Put for n the values -2, -1, 0, 1, 2, 3, 4 successively and we get the six angles

$$-360^0 + 18^0,\ -180^0 - 18^0,\ 18^0,\ 180^0 - 18^0,\ 360^0 + 18^0,\ 540^0 - 18^0,$$

i.e. -342^0, -198^0, 18^0, 162^0, 378^0, 522^0.

† For if n be *even*, let it be $2m$, when $(-1)^{2m} = +1$; if n be *odd*, let it be $2r+1$, when $(-1)^{2r+1} = -1$.

The student is recommended to draw a figure in the above example. Also to draw a figure in each example of this kind which he works for exercise.

EXAMPLES. XXXI.

(1) Find the four smallest angles which satisfy the equations

(i) $\sin A = \frac{1}{2}$. (ii) $\sin A = \frac{1}{\sqrt{2}}$. (iii) $\sin A = \frac{\sqrt{3}}{2}$. (iv) $\sin A = -\frac{1}{2}$.

(2) Find four angles between zero and $+8$ right angles which satisfy the equations

(i) $\sin A = \sin 20^0$. (ii) $\sin \theta = -\frac{1}{\sqrt{2}}$. (iii) $\sin \theta = -\sin \frac{\pi}{7}$.

(3) Find the complete algebraical solution of

(i) $\sin \theta = -\frac{1}{2}$. (ii) $2\sin^2\theta + 3\sin\theta = 2$. (iii) $\sin^2\theta = \cos^2\theta$.

(4) Prove that 30^0, 150^0, -330^0, 390^0, -210^0 have all the same sine.

146. *To find the complete Geometrical Solution of the equation* $\cos\theta = a$.

With the usual construction, let the radius of the circle *RULD* be the unit of length, so that the measure of *OP* is 1.



From O draw on OR a line OM_1 such that its measure is a.

OM_1 will be drawn towards the right or towards the left according as a is positive or negative.

Through M_1 draw $P_1M_1P_2$ perpendicular to OR to cut the circle in P_1, P_2. Join OP_1, OP_2.

Then $$\frac{OM_1}{OP_1} = \frac{a}{1} = \frac{OM_1}{OP_2}.$$

Hence any angle described when OP, starting from OR, stops in the position OP_1, or in the position OP_2, is an angle whose cosine is a.

And no other angle has its cosine $= a$, for there is no other possible position of M_1.

147. *To find the complete Algebraical Solution of the equation* $\cos\theta = a$.

With the figure and construction of the last article; let α be the least angle whose cosine $= a$.

Then in the figure $ROP_1 = \alpha$ and $P_2OR = \alpha$.

Every angle whose cosine $= a$ is included among those described when OP, starting from OR, stops in one or the other of the positions OP_1 or OP_2.

If OP starting from OR stop in the position OP_1, the angle described is one of those included in the expression

$$2m\pi + \alpha. \qquad \text{[Art. 130.]}$$

If OP starting from OR stop in the position OP_2, the angle described is one of those included in the expression

$$2r\pi - \alpha.$$

Both of these expressions are included in

$$2n\pi \pm \alpha.$$

Thus the solution of the equation $\cos\theta = \cos\alpha$ is

$$\theta = 2n\pi \pm \alpha.$$

148. *To find the complete Geometrical Solution of the equation* $\tan\theta = a$.

From O draw on the line OR two lines OM_1, OM_2, whose measures are $+1$ and -1 respectively.



And draw perpendicular to LOR from M_1 a line M_1P_1 whose measure is a, and from M_2 a line M_2P_2 whose measure is $-a$. Join OP_1, OP_2.

Then $$\frac{M_1P_1}{OM_1} = \frac{a}{1} = a, \text{ and } \frac{M_2P_2}{OM_2} = \frac{-a}{-1} = a.$$

Hence any angle described when OP, starting from OR, stops in one or the other of the positions OP_1 or OP_2, is an angle whose **tangent** is a.

And no other angle has its tangent $= a$.

149. *To find the complete Algebraical Solution of the equation* $\tan\theta = a$.

With the figure and construction of the last article; let α be the least angle whose tangent is a.

Then in the figure $ROP_1 = \alpha$ and $LOP_2 = \alpha$.

Every angle whose tangent is a is included among those described by OP, starting from OR and stopping in one or the other of the positions OP_1 or OP_2.

If OP starting from OR stop in the position OP_1, the angle described is one of those included in the expression

$$2m\pi + \alpha. \qquad \text{[Art. 130.]}$$

If OP starting from OR stop in the position OP_2, the angle described is one of those included in the expression

$$(2r+1)\,\pi + \alpha.$$

Both of these expressions are included in

$$n\pi + \alpha^*.$$

Thus the solution of the equation $\tan\theta = \tan\alpha$ is

$$\theta = n\pi + \alpha.$$

EXAMPLES. XXXII.

(1) Write down the complete Algebraical Solution of each of the following equations:

(i) $\cos\theta = \frac{1}{2}$. (ii) $\tan\theta = 1$. (iii) $\tan\theta = -1$.

(iv) $\tan\theta = -\sqrt{3}$. (v) $\cos\theta = \cos\frac{4\pi}{5}$. (vi) $\tan\theta = \tan\frac{3\pi}{4}$.

(2) Show that each of the following angles has the same cosine:

$$-120^0,\quad 240^0,\quad 480^0,\quad -480^0.$$

(3) The angles 60^0 and -120^0 have one of the Trigonometrical Ratios the same for both; which of the ratios is it?

(4) Can the following angles have any one of their Trigonometrical Ratios the same for all?

$$-23^0,\quad -157^0 \text{ and } 157^0.$$

(5) Find four angles which satisfy each of the equations in (1).

* For if n be even, this is the first formula; if n be odd it is the second.

150. We can now point out the use of the ambiguous sign $\pm$ in the formula $\cos\theta = \pm\sqrt{1-\sin^2\theta}$.

If we know the numerical value of the sine of an angle θ, without knowing the magnitude of the angle, we cannot from the identity, $\cos^2\theta = 1-\sin^2\theta$, *completely* determine $\cos\theta$, for we get $\cos\theta = \pm\sqrt{1-\sin^2\theta}$.

This is a **general formula**, and we shall find that it represents an important Geometrical truth.

151. Given $\sin\theta = a$, we can say that θ is *one of the angles* represented by one or the other of the positions OP_1, OP_2 of the revolving line in Fig. I. on page 110.

If we attempt to find the cosine of these angles we get two different values for the cosine; for $\frac{OM_1}{OP_1}$ and $\frac{OM_2}{OP_2}$ although equal in magnitude, are *opposite in sign*. Hence, if α be the least angle whose sine is equal to a, we have

$$\cos\theta = \pm\cos\alpha = \pm\sqrt{1-\sin^2\alpha}.$$

152. The same result may be obtained from the formula $\theta = n\pi + (-1)^n\alpha$. For $\cos\{n\pi + (-1)^n\alpha\}$ is of different sign according as n is even or odd.

EXAMPLES. XXXIII.

(1) If θ be found from the equation $\cos\theta = a$, show geometrically that there are two values of $\sin\theta$ and of $\tan\theta$.

(2) If θ be found from the equation $\tan\theta = a$, show geometrically that there are two values of $\sin\theta$ and of $\cos\theta$.

(3) If A be the *least* angle without regard to sign such that $\sin A = a$, show that $\cos A = +\sqrt{1-\sin^2 A}$.

(4) If A be the least *positive* angle such that $\cos A = a$, prove that $\sin A = +\sqrt{1-\cos^2 A}$.

CHAPTER XI.

On the Trigonometrical Ratios of Two Angles.

153. We proceed to establish the following fundamental formulæ:

$$\left.\begin{aligned} \sin(A+B) &= \sin A \,.\, \cos B + \cos A \,.\, \sin B \\ \cos(A+B) &= \cos A \,.\, \cos B - \sin A \,.\, \sin B \\ \sin(A-B) &= \sin A \,.\, \cos B - \cos A \,.\, \sin B \\ \cos(A-B) &= \cos A \,.\, \cos B + \sin A \,.\, \sin B \end{aligned}\right\} \ldots \text{(i)}.$$

Here, A and B are angles; so that $(A+B)$ and $(A-B)$ are also angles.

Hence, $\sin(A+B)$ is the sine of an angle, and must not be confounded with $\sin A + \sin B$.

$\sin(A+B)$ is a single fraction.

$\sin A + \sin B$ is the sum of two fractions.

154. The *proofs* given in the next two pages are perfectly general, as will be explained below (cf. Art. 169); but the *figures* are drawn for the simplest case in each.

The student should notice that the words of the two proofs are very nearly the same.

To prove that $\sin(A+B)=\sin A\,.\,\cos B+\cos A\,.\,\sin B$,
and that $\cos(A+B)=\cos A\,.\,\cos B-\sin A\,.\,\sin B$.



Let ROE be the angle A, and EOF the angle B. Then in the figure, ROF is the angle $(A+B)$.

In OF, *the line which bounds the compound angle* $(A+B)$, take any point P, and from P draw PM, PN at right angles to OR and OE respectively. Draw NH, NK at right angles to MP and OR respectively. Then the angle

$$NPH=90^0-HNP=HNO=ROE=A^*.$$

$$\text{Now } \sin(A+B)=\sin ROF=\frac{MP}{OP}=\frac{MH+HP}{OP}=\frac{KN}{OP}+\frac{HP}{OP}$$

$$=\frac{KN\,.\,\text{ON}}{\text{ON}\,.\,OP}+\frac{HP\,.\,\text{NP}}{\text{NP}\,.\,OP}=\frac{KN}{ON}\cdot\frac{ON}{OP}+\frac{HP}{NP}\cdot\frac{NP}{OP}$$

$$=\sin ROE\,.\,\cos EOF+\cos HPN\,.\,\sin EOF$$

$$=\sin A\,.\,\cos B+\cos A\,.\,\sin B.$$

$$\text{Also } \cos(A+B)=\cos ROF=\frac{OM}{OP}=\frac{OK-MK}{OP}=\frac{OK}{OP}-\frac{HN}{OP}$$

$$=\frac{OK\,.\,\text{ON}}{\text{ON}\,.\,OP}-\frac{HN\,.\,\text{NP}}{\text{NP}\,.\,OP}=\frac{OK}{ON}\cdot\frac{ON}{OP}-\frac{HN}{NP}\cdot\frac{NP}{OP}$$

$$=\cos ROE\,.\,\cos EOF-\sin HPN\,.\,\sin EOF$$

$$=\cos A\,.\,\cos B-\sin A\,.\,\sin B.$$

* Or thus. A circle goes round $OMNP$, because the angles OMP and ONP are right angles; therefore MPN and MON are angles in the same segment; so that $MPN=MON=A$.

To prove that $\sin(A-B)=\sin A\,.\cos B-\cos A\,.\sin B$,
and that $\cos(A-B)=\cos A\,.\cos B+\sin A\,.\sin B$.



Let ROE be the angle A, and FOE the angle B. Then in the figure, ROF is the angle $(A-B)$.

In OF, *the line which bounds the compound angle* $(A-B)$, take any point P, and from P draw PM, PN at right angles to OR and OE respectively. Draw NH, NK at right angles to MP and OR respectively. Then the angle

$$NPH=90^0-HNP=HNE=ROE=A^*.$$

$$\text{Now } \sin(A-B)=\sin ROF=\frac{MP}{OP}=\frac{MH-PH}{OP}=\frac{KN}{OP}-\frac{PH}{OP}$$

$$=\frac{KN.\,\mathrm{ON}}{\mathrm{ON}.\,OP}-\frac{PH.\,\mathrm{NP}}{\mathrm{NP}.\,OP}=\frac{KN}{ON}\cdot\frac{ON}{OP}-\frac{PH}{NP}\cdot\frac{NP}{OP}$$

$$=\sin ROE\,.\cos FOE-\cos HPN\,.\sin FOE$$

$$=\sin A\,.\cos B-\cos A\,.\sin B.$$

$$\text{Also } \cos(A-B)=\cos ROF=\frac{OM}{OP}=\frac{OK+KM}{OP}=\frac{OK}{OP}+\frac{NH}{OP}$$

$$=\frac{OK.\,\mathrm{ON}}{\mathrm{ON}.\,OP}+\frac{NH.\,\mathrm{NP}}{\mathrm{NP}.\,OP}=\frac{OK}{ON}\cdot\frac{ON}{OP}+\frac{NH}{NP}\cdot\frac{NP}{OP}$$

$$=\cos ROE\,.\cos FOE+\sin HPN\,.\sin FOE$$

$$=\cos A\,.\cos B+\sin A\,.\sin B.$$

* Or thus. A circle goes round $OMPN$, because the angles OMP and ONP are right angles; therefore MPN and MON together make up two right angles; so that $HPN=MON=A$.

Example. Find the value of $\sin 75^0$.

$$\sin 75^0 = \sin(45^0 + 30^0)$$
$$= \sin 45^0 . \cos 30^0 + \cos 45^0 . \sin 30^0$$
$$= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} \cdot \frac{1}{2}$$
$$= \frac{\sqrt{3}+1}{2\sqrt{2}} = \frac{\sqrt{2}(\sqrt{3}+1)}{4}.$$

EXAMPLES. XXXIV.

(1) Show that $\cos 75^0 = \frac{\sqrt{3}-1}{2\sqrt{2}}$.

(2) Show that $\sin 15^0 = \frac{\sqrt{3}-1}{2\sqrt{2}}$.

(3) Show that $\cos 15^0 = \frac{\sqrt{3}+1}{2\sqrt{2}}$.

(4) Show that $\tan 75^0 = 2 + \sqrt{3}$.

(5) If $\sin A = \frac{4}{5}$ and $\sin B = \frac{3}{5}$, find a value for $\sin(A+B)$ and for $\cos(A-B)$.

(6) If $\sin A = \cdot 6$ and $\sin B = \frac{5}{13}$, find a value for $\sin(A-B)$ and for $\cos(A+B)$.

(7) If $\sin A = \frac{1}{\sqrt{5}}$ and $\sin B = \frac{1}{\sqrt{10}}$, show that one value of $(A+B)$ is 45^0.

(8) Prove that $\sin 75^0 = \cdot 9659\ldots$

(9) Prove that $\sin 15^0 = \cdot 2588\ldots$

(10) Prove that $\tan 15^0 = \cdot 2679\ldots$

155. It is important that the student should become thoroughly familiar with the formulæ proved on the last two pages, and that he should be able to work examples involving their use.

EXAMPLES. XXXV.

Prove the following statements.

(1) $\sin(A+B)+\sin(A-B)=2\sin A\,.\cos B.$

(2) $\sin(A+B)-\sin(A-B)=2\cos A\,.\sin B.$

(3) $\cos(A+B)+\cos(A-B)=2\cos A\,.\cos B.$

(4) $\cos(A-B)-\cos(A+B)=2\sin A\,.\sin B.$

(5) $\dfrac{\sin(A+B)+\sin(A-B)}{\cos(A+B)+\cos(A-B)}=\tan A.$

(6) $\tan\alpha+\tan\beta=\dfrac{\sin(\alpha+\beta)}{\cos\alpha\,.\cos\beta}.$

(7) $\tan\alpha-\tan\beta=\dfrac{\sin(\alpha-\beta)}{\cos\alpha\,.\cos\beta}.$

(8) $\cot\alpha+\tan\beta=\dfrac{\cos(\alpha-\beta)}{\sin\alpha\,.\cos\beta}.$

(9) $\cot\alpha-\tan\beta=\dfrac{\cos(\alpha+\beta)}{\sin\alpha\,.\cos\beta}.$

(10) $\tan\alpha+\cot\beta=\dfrac{\cos(\alpha-\beta)}{\cos\alpha\,.\sin\beta}.$

(11) $\dfrac{\tan\theta+\tan\phi}{\tan\theta-\tan\phi}=\dfrac{\sin(\theta+\phi)}{\sin(\theta-\phi)}.$

(12) $\dfrac{\tan\theta\,.\tan\phi+1}{1-\tan\theta\,.\tan\phi}=\dfrac{\cos(\theta-\phi)}{\cos(\theta+\phi)}.$

(13) $\dfrac{\tan\theta+\cot\phi}{\cot\phi-\tan\theta}=\cos(\theta-\phi)\,.\sec(\theta+\phi).$

(14) $\dfrac{\cot\theta+\cot\phi}{\cot\theta-\cot\phi}=\dfrac{\sin(\theta+\phi)}{\sin(\theta-\phi)}.$

(15) $\dfrac{\tan\theta\,.\cot\phi+1}{\tan\theta\,.\cot\phi-1}=\dfrac{\sin(\theta+\phi)}{\sin(\theta-\phi)}.$

(16) $\dfrac{1+\cot\gamma\,.\tan\delta}{\cot\gamma-\tan\delta}=\tan(\gamma+\delta).$

(17) $\dfrac{1-\cot\gamma\,.\tan\delta}{\cot\gamma+\tan\delta}=\tan(\gamma-\delta).$

(18) $\dfrac{\tan\gamma\,,\cot\delta-1}{\tan\gamma+\cot\delta}=\tan(\gamma-\delta).$

(19) $\dfrac{\tan\gamma\,.\cot\delta+1}{\cot\delta-\tan\gamma}=\tan(\gamma+\delta).$

(20) $\dfrac{\cot\delta-\cot\gamma}{\cot\gamma\,.\cot\delta+1}=\tan(\gamma-\delta).$

(21) $\tan^2\alpha-\tan^2\beta=\dfrac{\sin(\alpha+\beta)\,.\sin(\alpha-\beta)}{\cos^2\alpha\,.\cos^2\beta}.$

(22) $\cot^2\alpha-\tan^2\beta=\dfrac{\cos(\alpha+\beta)\,.\cos(\alpha-\beta)}{\sin^2\alpha\,.\cos^2\beta}.$

(23) $\dfrac{\tan^2\alpha-\tan^2\beta}{1-\tan^2\alpha\,.\tan^2\beta}=\tan(\alpha+\beta)\,.\tan(\alpha-\beta).$

(24) $\sin(\alpha+\beta)\,.\sin(\alpha-\beta)=\sin^2\alpha-\sin^2\beta=\cos^2\beta-\cos^2\alpha.$

(25) $\cos(\alpha+\beta)\,.\cos(\alpha-\beta)=\cos^2\alpha-\sin^2\beta=\cos^2\beta-\sin^2\alpha.$

(26) $\sin(A-45^0)=\dfrac{\sin A-\cos A}{\sqrt{2}}.$

(27) $\sqrt{2}\,.\sin(A+45^0)=\sin A+\cos A.$

(28) $\cos A-\sin A=\sqrt{2}\,.\cos(A+45^0).$

(29) $\cos(A+45^0)+\sin(A-45^0)=0.$

(30) $\cos(A-45^0)=\sin(A+45^0).$

(31) $\sin(\theta+\phi)\,.\cos\theta-\cos(\theta+\phi)\,.\sin\theta=\sin\phi.$

(32) $\sin(\theta-\phi)\,.\cos\phi+\cos(\theta-\phi)\,.\sin\phi=\sin\theta.$

(33) $\cos(\theta+\phi)\,.\cos\theta+\sin(\theta+\phi)\,.\sin\theta=\cos\phi.$

(34) $\dfrac{\tan(\theta-\phi)+\tan\phi}{1-\tan(\theta-\phi)\,.\tan\phi}=\tan\theta.$

(35) $\dfrac{\tan(\theta+\phi)-\tan\theta}{1+\tan(\theta+\phi)\,.\tan\theta}=\tan\phi.$

(36) $2\sin\left(\alpha+\dfrac{\pi}{4}\right).\cos\left(\beta-\dfrac{\pi}{4}\right)=\cos(\alpha-\beta)+\sin(\alpha+\beta).$

(37) $2\sin\left(\dfrac{\pi}{4}-\alpha\right).\cos\left(\dfrac{\pi}{4}+\beta\right)=\cos(\alpha-\beta)-\sin(\alpha+\beta).$

(38) $\cos(\alpha+\beta)+\sin(\alpha-\beta)=2\sin\left(\dfrac{\pi}{4}+\alpha\right).\cos\left(\dfrac{\pi}{4}+\beta\right).$

(39) $\cos(\alpha+\beta)-\sin(\alpha-\beta)=2\sin\left(\dfrac{\pi}{4}-\alpha\right).\cos\left(\dfrac{\pi}{4}-\beta\right).$

(40) $\sin nA \,.\, \cos A + \cos nA \,.\, \sin A = \sin (n+1)\, A.$

(41) $\cos (n-1)\, A \,.\, \cos A - \sin (n-1)\, A \,.\, \sin A = \cos nA.$

(42) $\sin nA \,.\, \cos (n-1)\, A - \cos nA \,.\, \sin (n-1)\, A = \sin A.$

(43) $\cos (n-1)\, A \,.\, \cos (n+1)\, A - \sin (n-1)\, A \,.\, \sin (n+1)\, A = \cos 2nA.$

156. The following formulæ are important:

$$\left.\begin{aligned} \tan (A+B) &= \frac{\tan A + \tan B}{1 - \tan A \,.\, \tan B} \\ \tan (A-B) &= \frac{\tan A - \tan B}{1 + \tan A \,.\, \tan B} \end{aligned}\right\} \quad \ldots\ldots\ldots \text{(ii)}.$$

$$\tan (A+45^\circ) = \frac{\tan A + 1}{1 - \tan A},$$

$$\tan (A-45^\circ) = \frac{\tan A - 1}{1 + \tan A}.$$

The proof of the first is given below. The student should prove the second in a similar manner. The third and fourth follow at once from the first two by putting 45° for B.

Example. To prove $\tan (A+B) = \dfrac{\tan A + \tan B}{1 - \tan A \,.\, \tan B}$.

(i) By using the results of Art. 154, we have

$$\tan (A+B) = \frac{\sin (A+B)}{\cos (A+B)} = \frac{\sin A \,.\, \cos B + \cos A \,.\, \sin B}{\cos A \,.\, \cos B - \sin A \,.\, \sin B}.$$

Divide the numerator and the denominator of this fraction each by $\cos A \,.\, \cos B$, and we get

$$\tan (A+B) = \frac{\dfrac{\sin A \,.\, \cos B}{\cos A \,.\, \cos B} + \dfrac{\cos A \,.\, \sin B}{\cos A \,.\, \cos B}}{\dfrac{\cos A \,.\, \cos B}{\cos A \,.\, \cos B} - \dfrac{\sin A \,.\, \sin B}{\cos A \,.\, \cos B}}$$

$$= \frac{\tan A + \tan B}{1 - \tan A \,.\, \tan B}. \quad \text{Q.E.D.}$$

(ii) * * By Geometry. Make the construction of page 118 ;



Then $\tan(A+B)=\tan ROF=\dfrac{MP}{OM}=\dfrac{MH+HP}{OK-MK}$

$$=\frac{\dfrac{KN}{OK}+\dfrac{HP}{OK}}{\dfrac{OK}{OK}-\dfrac{HN}{OK}}=\frac{\dfrac{KN}{OK}+\dfrac{HP}{OK}}{1-\dfrac{HN\,.\,HP}{HP\,.\,OK}}=\frac{\tan A+\dfrac{HP}{OK}}{1-\tan A\,.\,\dfrac{HP}{OK}}.$$

Now the triangles NOK and NPH are similar,

$$\therefore \frac{HP}{OK}=\frac{NP}{ON}=\tan B,$$

$$\therefore \tan(A+B)=\frac{\tan A+\tan B}{1-\tan A\,.\,\tan B}.$$ Q.E.D.

EXAMPLES. XXXVI.

(1) If $\tan A=\frac{1}{2}$ and $\tan B=\frac{1}{4}$, prove that $\tan(A+B)=\frac{6}{7}$, and $\tan(A-B)=\frac{2}{9}$.

(2) If $\tan A=1$ and $\tan B=\dfrac{1}{\sqrt{3}}$, prove that

$$\tan(A+B)=2+\sqrt{3}.$$

(3) Prove that $\tan 15^0=2-\sqrt{3}$.

(4) If $\tan A=\frac{5}{6}$ and $\tan B=\frac{1}{11}$, prove that $\tan(A+B)=1$. What is $(A+B)$ in this case?

(5) If $\tan A = m$ and $\tan B = \frac{1}{m}$, prove that $\tan (A+B) = \infty$. What is $(A+B)$ in this case?

Prove the following statements:

(6) $\cot (A+B) = \dfrac{\cot A \,.\, \cot B - 1}{\cot A + \cot B}$.

(7) $\cot (A-B) = \dfrac{\cot A \,.\, \cot B + 1}{\cot B - \cot A}$.

(8) $\cot \left(\theta - \frac{\pi}{4}\right) = \dfrac{\cot \theta + 1}{1 - \cot \theta}$.

(9) $\dfrac{\cot \theta - 1}{\cot \theta + 1} = \cot \left(\theta + \frac{\pi}{4}\right)$.

(10) $\tan \left(\theta - \frac{\pi}{4}\right) + \cot \left(\theta + \frac{\pi}{4}\right) = 0$.

(11) $\cot \left(\theta - \frac{\pi}{4}\right) + \tan \left(\theta + \frac{\pi}{4}\right) = 0$.

(12) If $\tan \alpha = \dfrac{m}{m+1}$ and $\tan \beta = \dfrac{1}{2m+1}$, prove that

$$\tan (\alpha + \beta) = 1.$$

(13) $\dfrac{\tan (n+1)\phi - \tan n\phi}{1 + \tan (n+1)\phi \,.\, \tan n\phi} = \tan \phi$.

(14) $\dfrac{\tan (n+1)\phi + \tan (1-n)\phi}{1 - \tan (n+1)\phi \,.\, \tan (1-n)\phi} = \tan 2\phi$.

(15) If $\tan \alpha = m$ and $\tan \beta = n$, prove that

$$\cos (\alpha + \beta) = \frac{1 - mn}{\sqrt{(1+m^2)(1+n^2)}}.$$

(16) If $\tan \alpha = (a+1)$ and $\tan \beta = (a-1)$, then $2 \cot (\alpha - \beta) = a^2$.

(17) If $\alpha + \beta + \gamma = 90^0$, then

$$\tan \gamma = \frac{1 - \tan \alpha \tan \beta}{\tan \alpha + \tan \beta}.$$

157. From pages 118 and 119 we have

$$\left.\begin{aligned}
\sin(A+B) &= \sin A \,.\, \cos B + \cos A \,.\, \sin B \\
\sin(A-B) &= \sin A \,.\, \cos B - \cos A \,.\, \sin B \\
\cos(A+B) &= \cos A \,.\, \cos B - \sin A \,.\, \sin B \\
\cos(A-B) &= \cos A \,.\, \cos B + \sin A \,.\, \sin B
\end{aligned}\right\} \ldots \text{(i).}$$

From these by addition and subtraction we get

$$\left.\begin{aligned}
\sin(A+B) + \sin(A-B) &= 2 \sin A \,.\, \cos B \\
\sin(A+B) - \sin(A-B) &= 2 \cos A \,.\, \sin B \\
\cos(A+B) + \cos(A-B) &= 2 \cos A \,.\, \cos B \\
\cos(A-B) - \cos(A+B) &= 2 \sin A \,.\, \sin B
\end{aligned}\right\}$$

Now put S for $(A+B)$,

and put T for $(A-B)$,

Then $S+T=2A$, and $S-T=2B$,

so that $A=\dfrac{S+T}{2}$, and $B=\dfrac{S-T}{2}$.

Hence the above results may be written

$$\left.\begin{aligned}
\sin S + \sin T &= 2 \sin \frac{S+T}{2} \,.\, \cos \frac{S-T}{2} \\
\sin S - \sin T &= 2 \cos \frac{S+T}{2} \,.\, \sin \frac{S-T}{2} \\
\cos S + \cos T &= 2 \cos \frac{S+T}{2} \,.\, \cos \frac{S-T}{2} \\
*\cos T - \cos S &= 2 \sin \frac{S+T}{2} \,.\, \sin \frac{S-T}{2}
\end{aligned}\right\} \ldots \text{(iii).}$$

* If A and B are each less than 90^0, then S, which is their *sum*, is greater than T, their *difference*. Therefore if S be less than 90^0, $\cos S$ is less than $\cos T$; so that $\cos T - \cos S$ is *positive*.

158. The formulæ (iii) are most important, and the student is recommended to get thoroughly familiar with them *in words*, thus:

(1) *The sum of the sines of two angles is equal to twice the sine of half their sum multiplied by the cosine of half their difference.*

(2) *The sine of the greater of two angles minus the sine of the lesser is equal to twice the cosine of half their sum multiplied by the sine of half their difference.*

(3) *The sum of the cosines of two angles is equal to twice the cosine of half their sum multiplied by the cosine of half their difference.*

(4) *The cosine of the lesser of two angles minus the cosine of the greater is equal to twice the sine of half their sum multiplied by the sine of half their difference.*

159. It will be convenient to refer to the formulæ (i) as the 'A,B' formulæ, and to the formulæ (iii) as the 'S,T' formulæ.

160. The 'S,T' formulæ can be proved directly from a figure.

We give the proof for the case in which S and T are each less than 90°.

On first reading the subject however the student should omit the next two pages, and all other alternative proofs.

**(1) *To prove that*

$$\sin S + \sin T = 2 \sin \frac{S+T}{2} \cdot \cos \frac{S-T}{2}.$$



In the figure, let ROE be the angle S, ROF the angle T; then FOE is the angle $S-T$. Let OG bisect the angle EOF.

Then the angle $FOG = \frac{FOE}{2} = \frac{S-T}{2}$.

Also the angle $ROG = ROF + FOG = T + \frac{S-T}{2} = \frac{S+T}{2}$.

In OE take any point P, and from OF cut off OQ equal to OP. Join PQ, cutting OG in K. Then OG which bisects the vertical angle of the isosceles triangle POQ, bisects PQ at right angles. Draw PM, KL, QN at right angles to OR, and draw QVW and KH parallel to NLM.

Then by parallels, since $PK = KQ$, therefore $PH = HW$ and $WV = VQ$. Hence,

$$\sin S + \sin T = \frac{MP}{OP} + \frac{NQ}{OQ} = \frac{MP + NQ}{OQ}$$

$$= \frac{(LK + HP) + (LK - WH)}{OQ} = \frac{2LK}{OQ}$$

$$= \frac{2LK \cdot OK}{OK \cdot OQ} = 2 \sin ROK \cdot \cos QOK$$

$$= 2 \sin \frac{S+T}{2} \cdot \cos \frac{S-T}{2}. \quad \text{Q. E. D.}$$

(2) *To prove*

$$\cos S+\cos T=2\cos\frac{S+T}{2}\,.\,\cos\frac{S-T}{2}.$$

With the same figure and construction, we have

$$\cos S+\cos T=\frac{OM}{OP}+\frac{ON}{OQ}=\frac{OM+ON}{OQ},$$

$$=\frac{(OL-ML)+(OL+LN)}{OQ}=\frac{2OL}{OQ},$$

$$=\frac{2OL\,.\,\mathrm{OK}}{\mathrm{OK}\,.\,OQ}=2\cos ROK\,.\,\cos QOK,$$

$$=2\cos\frac{S+T}{2}\,.\,\cos\frac{S-T}{2}.\quad \text{Q.E.D.}$$

(3) *To prove*

$$\sin S-\sin T=2\sin\frac{S-T}{2}\,.\,\cos\frac{S+T}{2}.$$

In the above figure the angles VKQ, ROK are each the complement of LKO $\quad\therefore\ VKQ=ROK=\frac{S+T}{2}$. Hence

$$\sin S-\sin T=\frac{MP}{OP}-\frac{NQ}{OQ}=\frac{MP-NQ}{OQ},$$

$$=\frac{(LK+HP)-(LK-WH)}{OQ}=\frac{2VK}{OQ},$$

$$=\frac{2VK\,.\,\mathrm{KQ}}{\mathrm{KQ}\,.\,OQ}=2\cos VKQ\,.\,\sin QOK,$$

$$=2\sin\frac{S-T}{2}\,.\,\cos\frac{S+T}{2}.\quad \text{Q.E.D.}$$

(4) *To prove*

$$\cos T-\cos S=2\sin\frac{S+T}{2}\,.\,\sin\frac{S-T}{2}.$$

With the same figure and construction, we have

$$\cos T-\cos S=\frac{ON}{OQ}-\frac{OM}{OP}=\frac{ON-OM}{OQ},$$

$$=\frac{(OL+LN)-(OL-ML)}{OQ}=\frac{2LN}{OQ},$$

$$=\frac{2VQ\,.\,\mathrm{KQ}}{\mathrm{KQ}\,.\,OQ}=2\sin VKQ\,.\,\sin QOK,$$

$$=2\sin\frac{S+T}{2}\,.\,\sin\frac{S-T}{2}.\quad \text{Q.E.D.}$$

EXAMPLES. XXXVII.

Prove the following statements :

(1) $\sin 60^0 + \sin 30^0 = 2 \sin 45^0 . \cos 15^0$.

(2) $\sin 60^0 + \sin 20^0 = 2 \sin 40^0 . \cos 20^0$.

(3) $\sin 40^0 - \sin 10^0 = 2 \cos 25^0 . \sin 15^0$.

(4) $\cos \frac{\pi}{3} + \cos \frac{\pi}{2} = 2 \cos \frac{5\pi}{12} . \cos \frac{\pi}{12}$.

(5) $\cos \frac{\pi}{3} - \cos \frac{\pi}{2} = 2 \sin \frac{5\pi}{12} . \sin \frac{\pi}{12}$.

(6) $\sin 3A + \sin 5A = 2 \sin 4A . \cos A$.

(7) $\sin 7A - \sin 5A = 2 \cos 6A . \sin A$.

(8) $\cos 5A + \cos 9A = 2 \cos 7A . \cos 2A$.

(9) $\cos 5A - \cos 4A = -2 \sin \frac{9A}{2} . \sin \frac{A}{2}$.

(10) $\cos A - \cos 2A = 2 \sin \frac{3A}{2} . \sin \frac{A}{2}$.

(11) $\frac{\sin 2\theta + \sin \theta}{\cos \theta + \cos 2\theta} = \tan \frac{3\theta}{2}$.

(12) $\frac{\sin 2\theta - \sin \theta}{\cos \theta - \cos 2\theta} = \cot \frac{3\theta}{2}$.

(13) $\frac{\sin 3\theta + \sin 2\theta}{\cos 2\theta - \cos 3\theta} = \cot \frac{\theta}{2}$.

(14) $\frac{\sin \theta + \sin \phi}{\cos \theta - \cos \phi} = \frac{\cos \theta + \cos \phi}{\sin \phi - \sin \theta}$.

(15) $\cos (60^0 + A) + \cos (60^0 - A) = \cos A$.

(16) $\cos (45^0 + A) + \cos (45^0 - A) = \sqrt{2} . \cos A$.

(17) $\sin (45^0 + A) - \sin (45^0 - A) = \sqrt{2} . \sin A$.

(18) $\cos (30^0 - A) - \cos (30^0 + A) = \sin A$.

(19) $\frac{\sin \theta - \sin \phi}{\cos \phi - \cos \theta} = \cot \frac{\theta + \phi}{2}$.

(20) $\frac{\sin \theta - \sin \phi}{\sin \theta + \sin \phi} = \cot \left(\frac{\theta + \phi}{2} \right) . \tan \left(\frac{\theta - \phi}{2} \right)$.

161. It is important that the student should be thoroughly familiar with the second set of formulæ on p. 126.

Written as follows, they may be regarded as the inverse of the '*S*, *T*' formulæ.

$$\left.\begin{aligned} 2\sin A\,.\cos B &= \sin(A+B)+\sin(A-B),\\ 2\cos A\,.\sin B &= \sin(A+B)-\sin(A-B),\\ 2\cos A\,.\cos B &= \cos(A+B)+\cos(A-B),\\ 2\sin A\,.\sin B &= \cos(A-B)-\cos(A+B). \end{aligned}\right\}\ \text{iv.}$$

EXAMPLES. XXXVIII.

Express as the sum or as the difference of two trigonometrical ratios the ten following expressions :

(1) $2\sin\theta\,.\cos\phi$.

(2) $2\cos\alpha\,.\cos\beta$.

(3) $2\sin 2\alpha\,.\cos 3\beta$.

(4) $2\cos(\alpha+\beta)\,.\cos(\alpha-\beta)$.

(5) $2\sin 3\theta\,.\cos 5\theta$.

(6) $2\cos\frac{3\theta}{2}\,.\cos\frac{\theta}{2}$.

(7) $\sin 4\theta\,.\sin\theta$.

(8) $\cos\frac{5\theta}{2}\,.\sin\frac{3\theta}{2}$.

(9) $2\cos 10^0\,.\sin 50^0$.

(10) $\cos 45^0\,.\sin 15^0$.

(11) Simplify $2\cos 2\theta\,.\cos\theta-2\sin 4\theta\,.\sin\theta$.

(12) Simplify $\sin\frac{5\theta}{2}\,.\cos\frac{\theta}{2}-\sin\frac{9\theta}{2}\,.\cos\frac{3\theta}{2}$

(13) Simplify $\sin 3\theta+\sin 2\theta+2\sin\frac{3\theta}{2}\,.\cos\frac{\theta}{2}$.

(14) Prove that $\sin\frac{11\theta}{4}\,.\sin\frac{\theta}{4}+\sin\frac{7\theta}{4}\,.\sin\frac{3\theta}{4}=\sin 2\theta\,.\sin\theta$.

** MISCELLANEOUS EXAMPLES. XXXIX.

(1) If $\tan a=\frac{1}{2}$ and $\tan \beta=\frac{1}{3}$, prove that $\tan (a+\beta)=1$.

(2) If $\tan a=\frac{3}{4}$ and $\tan \beta=\frac{1}{7}$, prove that one of the values of $a+\beta$ is $\frac{\pi}{4}$.

(3) If $\tan a=\frac{m-1}{m}$ and $\tan \beta=\frac{1}{2m-1}$, shew that one value of $(a+\beta)$ is $\frac{\pi}{4}$.

(4) Simplify $\frac{\cos a-\cos 5a}{\sin a+\sin 5a}$.

(5) Simplify $\frac{\sin 5x-\sin 3x}{\cos 5x+\cos 3x}$.

(6) Prove that $\frac{\cos A+\cos 3A}{\cos 3A+\cos 5A}=\frac{\cos 2A}{\cos 4A}$.

(7) Simplify $\frac{\sin 3x-\sin x}{\cos 3x+\cos x}+\frac{\sin 3x+\sin x}{\cos 3x-\cos x}$.

(8) Simplify $\frac{(\sin 4A-\sin 2A)(\cos A-\cos 3A)}{(\cos 4A+\cos 2A)(\sin A+\sin 3A)}$.

(9) Prove that

$$2 \sin 2a \,.\, \cos a+2 \cos 4a \,.\, \sin a=\sin 5a+\sin a.$$

(10) Prove that

$$\cos 2a \,.\, \cos a-\sin 4a \,.\, \sin a=\cos 3a \,.\, \cos 2a.$$

(11) $\tan 2A \,.\, \tan 3A \,.\, \tan 5A=\tan 5A-\tan 3A-\tan 2A$.

(12) Solve $\left.\begin{aligned} 4 \sin (\theta+\phi) \,.\, \cos (\theta-\phi)&=3 \\ 4 \cos (\theta+\phi) \,.\, \sin (\theta-\phi)&=1 \end{aligned}\right\}$.

(13) Prove that

$$\frac{\sin A \,.\, \sin 2A+\sin 2A \,.\, \sin 5A+\sin 3A \,.\, \sin 10A}{\cos A \,.\, \sin 2A+\sin 2A \,.\, \cos 5A-\cos 3A \,.\, \sin 10A}=-\tan 7A.$$

(14) $\tan \frac{A+B}{2}-\tan \frac{A-B}{2}=\frac{2 \sin B}{\cos A+\cos B}$.

CHAPTER XII.

On the Trigonometrical Ratios of Multiple Angles.

162. To express the Trigonometrical Ratios of the angle $2A$ in terms of those of the angle A.

Since $\sin(A+B) = \sin A \,.\, \cos B + \cos A \,.\, \sin B$;

$\therefore \sin(A+A) = \sin A \,.\, \cos A + \cos A \,.\, \sin A$;

$\therefore \sin 2A = 2 \sin A \,.\, \cos A$ ..... (1).

Also, since $\cos(A+B) = \cos A \,.\, \cos B - \sin A \,.\, \sin B$;

$\therefore \cos(A+A) = \cos A \,.\, \cos A - \sin A \,.\, \sin A$;

$\therefore \cos 2A = \cos^2 A - \sin^2 A$ ..... (2).

But $1 = \cos^2 A + \sin^2 A$;

$\therefore 1 + \cos 2A = 2 \cos^2 A$,

and $1 - \cos 2A = 2 \sin^2 A$.

The last two results are usually written

$$\cos 2A = 2\cos^2 A - 1 \quad \ldots\ldots\ldots\ldots \quad (3),$$

$$\text{and } \cos 2A = 1 - 2\sin^2 A \quad \ldots\ldots\ldots\ldots \quad (4).$$

$$\text{Again, } \tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \,.\, \tan B};$$

$$\therefore \tan(A+A) = \frac{\tan A + \tan A}{1 - \tan A \,.\, \tan A};$$

$$\therefore \tan 2A = \frac{2\tan A}{1 - \tan^2 A} \quad \ldots\ldots\ldots\ldots \quad (5).$$

163.** *To prove the '2A' formulæ geometrically.*



Let ROP be the angle $2A$. With centre O and any radius describe the semicircle RPL. Draw PM perpendicular to OR. Join RP . PL.

Then the angle RPL in a semicircle is a right angle. The angle $ROP = OLP + OPL = 2\,OLP$ [since $OL = OP$]. $\therefore$ $OLP =$ a half of $ROP = A$. Also MPR and OLP are each the complement of MPL. $\therefore$ $MPR = OLP = A$.

Hence

(1) $$\sin 2A = \frac{MP}{OP} = \frac{2MP}{2OR} = \frac{2MP}{LR} = \frac{2MP \,.\, \mathrm{PR}}{\mathrm{PR} \,.\, LR}$$
$$= 2\cos MPR \,.\, \sin PLR = 2\sin A \,.\, \cos A.$$

(3) $$\cos 2A = \frac{OM}{OP} = \frac{LM - LO}{OP} = \frac{2LM}{2OP} - \frac{LO}{OP}$$
$$= \frac{2LM \,.\, \mathrm{LP}}{\mathrm{LP} \,.\, LR} - \frac{OP}{OP} = 2\cos MLP \,.\, \cos PLR - 1$$
$$= 2\cos^2 A - 1.$$

(2) Let $OM' = OM$.

Then $2OM = M'M = LM - LM' = LM - MR$. Hence,

$$\cos 2A = \frac{2OM}{2OP} = \frac{LM - MR}{LR} = \frac{LM}{LR} - \frac{MR}{LR},$$
$$= \frac{LM \,.\, \mathrm{LP}}{\mathrm{LP} \,.\, LR} - \frac{MR \,.\, \mathrm{PR}}{\mathrm{PR} \,.\, LR} = \cos^2 A - \sin^2 A.$$

(5) $$\tan 2A = \frac{2MP}{2OM} = \frac{2MP}{LM - MR},$$
$$= \frac{\dfrac{2MP}{LM}}{\dfrac{LM}{LM} - \dfrac{MR}{LM}} = \frac{2\tan A}{1 - \dfrac{MR \,.\, MP}{MP \,.\, LM}} = \frac{2\tan A}{1 - \tan^2 A}.$$

164. These five formulæ are very important,

$$\left.\begin{aligned} &\sin 2A = 2\sin A \,.\, \cos A \quad \ldots\ldots\ldots\ldots (1),\\ &\cos 2A = \cos^2 A - \sin^2 A \quad \ldots\ldots\ldots\ldots (2),\\ &\cos 2A = 2\cos^2 A - 1 \quad \ldots\ldots\ldots\ldots (3),\\ &\cos 2A = 1 - 2\sin^2 A \quad \ldots\ldots\ldots\ldots (4),\\ &\tan 2A = \frac{2\tan A}{1-\tan^2 A} \quad \ldots\ldots\ldots\ldots (5), \end{aligned}\right\} \text{v.}$$

165. The following result is important,

$$\frac{\sin 2A}{1+\cos 2A} = \frac{2\sin A \,.\, \cos A}{2\cos^2 A} = \tan A.$$

166. The student must notice that A is *any* angle, and therefore these formulæ will be true whatever we put for A.

Example. Write $\frac{A}{2}$ instead of A, and we get

$$\sin A = 2\sin\frac{A}{2} \,.\, \cos\frac{A}{2} \quad \ldots\ldots\ldots\ldots (1),$$

$$\cos A = \cos^2\frac{A}{2} - \sin^2\frac{A}{2} \quad \ldots\ldots\ldots\ldots (2),$$

and so on.

EXAMPLES. XL.

Prove the following statements :

(1) $2\operatorname{cosec} 2A = \sec A \,.\, \operatorname{cosec} A.$

(2) $\dfrac{\operatorname{cosec}^2 A}{\operatorname{cosec}^2 A - 2} = \sec 2A.$

(3) $\dfrac{2 - \sec^2 A}{\sec^2 A} = \cos 2A.$

(4) $\cos^2 A\,(1 - \tan^2 A) = \cos 2A$

(5) $\cot 2A = \dfrac{\cot^2 A - 1}{2\cot A}.$

(6) $\dfrac{2\tan B}{1+\tan^2 B}=\sin 2B.$

(7) $\tan B+\cot B=2\operatorname{cosec} 2B.$

(8) $\dfrac{1-\tan^2 B}{1+\tan^2 B}=\cos 2B.$

(9) $\cot B-\tan B=2\cot 2B.$

(10) $\dfrac{\cot^2 B+1}{\cot^2 B-1}=\sec 2B.$

(11) $\left(\sin\dfrac{\theta}{2}+\cos\dfrac{\theta}{2}\right)^2=1+\sin\theta.$

(12) $\left(\sin\dfrac{\theta}{2}-\cos\dfrac{\theta}{2}\right)^2=1-\sin\theta.$

(13) $\cos^2\dfrac{\theta}{2}\left(1+\tan\dfrac{\theta}{2}\right)^2=1+\sin\theta.$

(14) $\sin^2\dfrac{\theta}{2}\left(\cot\dfrac{\theta}{2}-1\right)^2=1-\sin\theta.$

(15) $\left(\dfrac{\tan\dfrac{\theta}{2}+1}{\tan\dfrac{\theta}{2}-1}\right)^2=\dfrac{1+\sin\theta}{1-\sin\theta}.$

(16) $\dfrac{\sin\beta}{1+\cos\beta}=\tan\dfrac{\beta}{2}.$

(17) $\dfrac{\sin\beta}{1-\cos\beta}=\cot\dfrac{\beta}{2}.$

(18) $\dfrac{1-\cos\beta}{1+\cos\beta}=\tan^2\dfrac{\beta}{2}.$

(19) $\dfrac{1+\sec\beta}{\sec\beta}=2\cos^2\dfrac{\beta}{2}.$

(20) $\operatorname{cosec}\beta-\cot\beta=\tan\dfrac{\beta}{2}.$

(21) $\dfrac{\cos 2x}{1+\sin 2x}=\dfrac{1-\tan x}{1+\tan x}.$

(22) $\dfrac{\cos x}{1-\sin x}=\dfrac{1+\tan\dfrac{x}{2}}{1-\tan\dfrac{x}{2}}.$

(23) $\dfrac{\cos x}{1+\sin x}=\dfrac{\cot\frac{x}{2}-1}{\cot\frac{x}{2}+1}$.

(24) $\dfrac{\cos x}{1-\sin x}=\dfrac{\cot\frac{x}{2}+1}{\cot\frac{x}{2}-1}$.

(25) $\dfrac{1+\sin x+\cos x}{1+\sin x-\cos x}=\cot\dfrac{x}{2}$.

(26) $\dfrac{\cos^3 a+\sin^3 a}{\cos a+\sin a}=\dfrac{2-\sin 2a}{2}$.

(27) $\dfrac{\cos^3 a-\sin^3 a}{\cos a-\sin a}=\dfrac{2+\sin 2a}{2}$.

(28) $\cos^4 a-\sin^4 a=\cos 2a$.

(29) $\cos^6 a+\sin^6 a=\dfrac{1+3\cos^2 2a}{4}$.

(30) $\cos^6 a-\sin^6 a=\dfrac{(3+\cos^2 2a)\cos 2a}{4}$.

(31) $\dfrac{\sin 3\beta}{\sin\beta}-\dfrac{\cos 3\beta}{\cos\beta}=2$.

(32) $\dfrac{\cos 3\beta}{\sin\beta}+\dfrac{\sin 3\beta}{\cos\beta}=2\cot 2\beta$.

(33) $\dfrac{\sin 4\beta}{\sin 2\beta}=2\cos 2\beta$.

(34) $\dfrac{\sin 5\beta}{\sin\beta}-\dfrac{\cos 5\beta}{\cos\beta}=4\cos 2\beta$.

(35) $\dfrac{\sin\frac{5\pi}{12}}{\sin\frac{\pi}{12}}-\dfrac{\cos\frac{5\pi}{12}}{\cos\frac{\pi}{12}}=2\sqrt{3}$.

(36) $\tan(45^0+A)-\tan(45^0-A)=2\tan 2A$.

(37) $\tan(45^0-A)+\cot(45^0-A)=2\sec 2A$.

(38) $\dfrac{\tan^2(45^0+A)-1}{\tan^2(45^0+A)+1}=\sin 2A.$

(39) $\dfrac{\sec A+\tan A}{\sec A-\tan A}=\tan\left(45^0+\dfrac{A}{2}\right)\,.\,\cot\left(45^0-\dfrac{A}{2}\right).$

(40) $\dfrac{\cos(A+45^0)}{\cos(A-45^0)}=\sec 2A-\tan 2A.$

(41) $\tan B=\dfrac{\sin B+\sin 2B}{1+\cos B+\cos 2B}.$

(42) $\tan B=\dfrac{\sin 2B-\sin B}{1-\cos B+\cos 2B}.$

167. The following two formulæ should be remembered:

$$\left.\begin{aligned}\sin 3A&=3\sin A-4\sin^3 A\\ \cos 3A&=4\cos^3 A-3\cos A\end{aligned}\right\}\ldots\ldots\ldots\ldots\text{(vi)}.$$

NOTE. The similarity of these two results is apt to cause confusion. This may be avoided by observing that the second formula must be true when $A=0^0$; and then $\cos 3A=\cos 0^0=1$. In which case the formula gives $\cos 0^0=4\cos 0^0-3\cos 0^0$, or $1=4-3$, which is true.

The first formula may be proved thus:

$$\begin{aligned}\sin 3A&=\sin(2A+A)=\sin 2A\,.\,\cos A+\cos 2A\,.\,\sin A\\ &=(2\sin A\,.\,\cos A)\cos A+(1-2\sin^2 A)\sin A\\ &=2\sin A\,.\cos^2 A+\sin A-2\sin^3 A\\ &=2\sin A(1-\sin^2 A)+\sin A-2\sin^3 A\\ &=2\sin A-2\sin^3 A+\sin A-2\sin^3 A\\ &=3\sin A-4\sin^3 A.\end{aligned}$$

The second formula may be proved in a similar manner.

Example. *Prove that* $\tan 3A = \dfrac{3\tan A - \tan^3 A}{1 - 3\tan^2 A}$.

$$\tan 3A = \tan(2A + A) = \frac{\tan 2A + \tan A}{1 - \tan 2A \tan A}$$

$$= \frac{\dfrac{2\tan A}{1 - \tan^2 A} + \tan A}{1 - \dfrac{2\tan A}{1 - \tan^2 A} \times \tan A}$$

$$= \frac{2\tan A + \tan A(1 - \tan^2 A)}{1 - \tan^2 A - 2\tan^2 A}$$

$$= \frac{3\tan A - \tan^3 A}{1 - 3\tan^2 A}.$$

EXAMPLES. XLI.

Prove the following statements :

(1) $\dfrac{\sin 3A}{\sin A} = 2\cos 2A + 1.$

(2) $\dfrac{\cos 3A}{\cos A} = 2\cos 2A - 1.$

(3) $\dfrac{3\sin A - \sin 3A}{\cos 3A + 3\cos A} = \tan^3 A.$

(4) $\cot 3A = \dfrac{\cot^3 A - 3\cot A}{3\cot^2 A - 1}.$

(5) $\dfrac{\sin 3A - \sin A}{\cos 3A + \cos A} = \tan A.$

(6) $\dfrac{\sin 3A - \cos 3A}{\sin A + \cos A} = 2\sin 2A - 1.$

(7) $\dfrac{\sin 3A + \cos 3A}{\cos A - \sin A} = 2\sin 2A + 1.$

(8) $\dfrac{1}{\tan 3A - \tan A} + \dfrac{1}{\cot A - \cot 3A} = \cot 2A.$

(9) $\left(\dfrac{3\sin A - \sin 3A}{3\cos A + \cos 3A}\right)^2 = \left(\dfrac{\sec 2A - 1}{\sec 2A + 1}\right)^3.$

(10) $\dfrac{1 - \cos 3A}{1 - \cos A} = (1 + 2\cos A)^2.$

**MISCELLANEOUS EXAMPLES. XLII.

Prove the following statements :

(1) $\dfrac{\sin A+\cos A}{\cos A-\sin A}=\tan 2A+\sec 2A.$

(2) $\dfrac{\tan\frac{A}{2}+1}{1-\tan\frac{A}{2}}=\tan A+\sec A.$

(3) $\sin(n+1)a.\cos(n-1)a-\sin 2a=\sin(n-1)a.\cos(n+1)a.$

(4) $\dfrac{\sin a+\sin\beta}{\cos a+\cos\beta}=\tan\dfrac{a+\beta}{2}.$

(5) $\dfrac{\cos 2a+\cos 12a}{\cos 6a+\cos 8a}+\dfrac{\cos 7a-\cos 3a}{\cos a-\cos 3a}+2\dfrac{\sin 4a}{\sin 2a}=0.$

(6) If $A=18^0$, prove that $\sin 2A=\cos 3A$; hence prove that

$$\sin 18^0=\frac{\sqrt{5}-1}{4}.$$

(7) $\dfrac{\sin a+\sin\beta+\sin(a+\beta)}{\sin a+\sin\beta-\sin(a+\beta)}=\cot\dfrac{a}{2}.\cot\dfrac{\beta}{2}.$

(8) $\sin 2A.\sin 2B=\sin^2(A+B)-\sin^2(A-B).$

(9) $\cos 4A=8\cos^4 A-8\cos^2 A+1.$

(10) $\tan 50^0+\cot 50^0=2\sec 10^0.$

(11) $\sin 3A=4\sin A.\sin(60^0+A)\sin(60^0-A).$

(12) $\left(\cot\dfrac{A}{2}-\tan\dfrac{A}{2}\right)^2(\cot A-2\cot 2A)=4\cot A.$

(13) $\dfrac{\cos 3a-\sin\beta.\sin 5a-\cos 7a}{\sin 3a+\sin\beta.\cos 5a-\sin 7a}$ is independent of β.

(14) $(\cos x+\cos y)^2+(\sin x+\sin y)^2=4\cos^2\dfrac{x-y}{2}.$

(15) $2\cos^2 A.\cos^2 B+2\sin^2 A.\sin^2 B=1+\cos 2A.\cos 2B.$

(16) $\cot\dfrac{\pi}{8}-\tan\dfrac{\pi}{8}=2.$

(17) $\tan 4\theta=\dfrac{4\tan\theta(1-\tan^2\theta)}{1-6\tan^2\theta+\tan^4\theta}.$

(18) $2\cos\frac{\pi}{8}=\sqrt{2+\sqrt{2}}$.

(19) $2\cos 11^0\,15'=\sqrt{2+\sqrt{2+\sqrt{2}}}$.

(20) $\dfrac{\sin A\,.\,\sin 2A+\sin A\,.\,\sin 4A+\sin 2A\,.\,\sin 7A}{\sin A\,.\,\cos 2A+\sin 2A\,.\,\cos 5A+\sin A\,.\,\cos 8A}=\tan 5A$.

(21) $\dfrac{\sin\theta+\sin(\theta+\phi)+\sin(\theta+2\phi)}{\cos\theta+\cos(\theta+\phi)+\cos(\theta+2\phi)}=\tan(\theta+\phi)$.

(22) $2\cos^8 A-2\sin^8 A=\cos 2A\,(1+\cos^2 2A)$.

(23) $(3\sin A-4\sin^3 A)^2+(4\cos^3 A-3\cos A)^2=1$.

(24) $\dfrac{\sin 2a\,.\,\cos a}{(1+\cos 2a)\,(1+\cos a)}=\tan\frac{a}{2}$.

(25) $2\,\dfrac{\cot(n-2)a\,.\,\cot na+1}{\cot(n-2)a-\cot na}=\cot a-\tan a$.

(26) If $\tan a=\frac{1}{7}$ and $\tan\beta=\frac{2}{11}$, prove $\tan(2a+\beta)=\frac{1}{2}$.

(27) Prove that $\tan\frac{A}{2}$ and $\cot\frac{A}{2}$ are the roots of the equation

$$x^2-2x\,.\,\operatorname{cosec} A+1=0.$$

(28) If $\tan B=\dfrac{b}{a}$, prove that $\sqrt{\dfrac{a+b}{a-b}}+\sqrt{\dfrac{a-b}{a+b}}=\dfrac{2\cos B}{\sqrt{\cos 2B}}$.

*168. The following examples are symmetrical, and each involve more than two angles:

Example 1. Prove that

$$\sin(a+\beta+\gamma)=\sin a\,.\,\cos\beta\,.\,\cos\gamma+\sin\beta\,.\,\cos\gamma\,.\,\cos a$$
$$+\sin\gamma\,.\,\cos a\,.\,\cos\beta-\sin a\,.\,\sin\beta\,.\,\sin\gamma.$$

$$\sin(a+\beta+\gamma)=\sin(a+\beta)\,.\,\cos\gamma+\cos(a+\beta)\sin\gamma$$
$$=\sin a\,.\,\cos\beta\,.\,\cos\gamma+\cos a\,.\,\sin\beta\,.\,\cos\gamma$$
$$+\cos a\,.\,\cos\beta\,.\,\sin\gamma-\sin a\,.\,\sin\beta\,.\,\sin\gamma$$
$$=\sin a\,.\,\cos\beta\,.\,\cos\gamma+\sin\beta\,.\,\cos\gamma\,.\,\cos a$$
$$+\sin\gamma\,.\,\cos a\,.\,\cos\beta-\sin a\,.\,\sin\beta\,.\,\sin\gamma.$$

Q.E.D.

Example 2. Prove that

$$\sin\alpha+\sin\beta+\sin\gamma-\sin(\alpha+\beta+\gamma)$$
$$=4\,.\,\sin\frac{\beta+\gamma}{2}\,.\,\sin\frac{\gamma+\alpha}{2}\,.\,\sin\frac{\alpha+\beta}{2}\,.$$

Now $\sin\alpha-\sin(\alpha+\beta+\gamma)=-2\cos\,.\frac{2\alpha+\beta+\gamma}{2}\,.\,\sin\frac{\beta+\gamma}{2}\,.$

And $\sin\beta+\sin\gamma=2\sin\frac{\beta+\gamma}{2}\,.\,\cos\frac{\beta-\gamma}{2}$, [Art. 158]

$\therefore\ \sin\alpha+\sin\beta+\sin\gamma-\sin(\alpha+\beta+\gamma)$

$$=2\sin\frac{\beta+\gamma}{2}\,.\,\cos\frac{\beta-\gamma}{2}-2\cos\frac{2\alpha+\beta+\gamma}{2}\,.\,\sin\frac{\beta+\gamma}{2}$$

$$=2\sin\frac{\beta+\gamma}{2}\left\{\cos\frac{\beta-\gamma}{2}-\cos\frac{2\alpha+\beta+\gamma}{2}\right\}$$

$$=2\sin\frac{\beta+\gamma}{2}\,.\,2\sin\frac{\gamma+\alpha}{2}\,.\,\sin\frac{\alpha+\beta}{2}$$ [Art. 158]

$$=4\sin\frac{\beta+\gamma}{2}\,.\,\sin\frac{\gamma+\alpha}{2}\,.\,\sin\frac{\alpha+\beta}{2}\,.$$ Q.E.D.

* EXAMPLES. XLIII.

Prove the following statements:

(1) $\cos(\alpha+\beta+\gamma)=\cos\alpha\,.\,\cos\beta\,.\,\cos\gamma-\cos\alpha\,.\,\sin\beta\,.\,\sin\gamma$
$-\cos\beta\,.\,\sin\gamma\,.\,\sin\alpha-\cos\gamma\,.\,\sin\alpha\,.\,\sin\beta.$

(2) $\sin(\alpha+\beta-\gamma)=\sin\alpha\,.\,\cos\beta\,.\,\cos\gamma+\sin\beta\,.\,\cos\gamma\,.\,\cos\alpha$
$-\sin\gamma\,.\,\cos\alpha\,.\,\cos\beta+\sin\alpha\sin\beta\sin\gamma.$

(3) $\cos(\alpha-\beta+\gamma)=\cos\alpha\,.\,\cos\beta\,.\,\cos\gamma+\cos\alpha\,.\,\sin\beta\,.\,\sin\gamma$
$-\cos\beta\,.\,\sin\alpha\,.\,\sin\gamma+\cos\gamma\,.\,\sin\beta\,.\,\sin\alpha.$

(4) $\sin\alpha+\sin\beta-\sin\gamma-\sin(\alpha+\beta-\gamma)$
$$=4\sin\frac{\alpha-\gamma}{2}\,.\,\sin\frac{\beta-\gamma}{2}\,.\,\sin\frac{\alpha+\beta}{2}\,.$$

(5) $\sin(\alpha-\beta-\gamma)-\sin\alpha+\sin\beta+\sin\gamma$
$$=4\sin\frac{\alpha-\beta}{2}\,.\,\sin\frac{\alpha-\gamma}{2}\,.\,\sin\frac{\beta+\gamma}{2}\,.$$

(6) $\sin2\alpha+\sin2\beta+\sin2\gamma-\sin2(\alpha+\beta+\gamma)$
$=4\sin(\beta+\gamma)\,.\,\sin(\gamma+\alpha)\,.\,\sin(\alpha+\beta).$

(7) $\sin(\beta-\gamma)+\sin(\gamma-\alpha)+\sin(\alpha-\beta)$
$$+4\sin\frac{(\beta-\gamma)}{2}.\sin\frac{(\gamma-\alpha)}{2}.\sin\frac{(\alpha-\beta)}{2}=0.$$

(8) $\sin(\beta+\gamma-\alpha)+\sin(\gamma+\alpha-\beta)+\sin(\alpha+\beta-\gamma)$
$$-\sin(\alpha+\beta+\gamma)=4\sin\alpha.\sin\beta.\sin\gamma.$$

(9) $\sin(\alpha+\beta+\gamma)+\sin(\beta+\gamma-\alpha)+\sin(\gamma+\alpha-\beta)$
$$-\sin(\alpha+\beta-\gamma)=4\cos\alpha.\cos\beta.\sin\gamma.$$

(10) $\cos x+\cos y+\cos z+\cos(x+y+z)$
$$=4\cos\frac{y+z}{2}.\cos\frac{z+x}{2}.\cos\frac{x+y}{2}.$$

(11) $\cos 2x+\cos 2y+\cos 2z+\cos 2(x+y+z)$
$$=4\cos(y+z).\cos(z+x).\cos(x+y).$$

(12) $\cos(y+z-x)+\cos(z+x-y)+\cos(x+y-z)$
$$+\cos(x+y+z)=4\cos x.\cos y.\cos z.$$

(13) $\cos^2 x+\cos^2 y+\cos^2 z+\cos^2(x+y+z)$
$$=2\{1+\cos(y+z).\cos(z+x).\cos(x+y)\}.$$

(14) $\sin^2 x+\sin^2 y+\sin^2 z+\sin^2(x+y+z)$
$$=2\{1-\cos(y+z).\cos(z+x).\cos(x+y)\}.$$

(15) $\cos^2 x+\cos^2 y+\cos^2 z+\cos^2(x+y-z)$
$$=2\{1+\cos(x-z).\cos(y-z).\cos(x+y)\}.$$

(16) $\cos\alpha.\sin(\beta-\gamma)+\cos\beta.\sin(\gamma-\alpha)+\cos\gamma.\sin(\alpha-\beta)=0.$

(17) $\sin\alpha.\sin(\beta-\gamma)+\sin\beta.\sin(\gamma-\alpha)+\sin\gamma.\sin(\alpha-\beta)=0.$

(18) $\cos(\alpha+\beta).\cos(\alpha-\beta)+\sin(\beta+\gamma)\sin(\beta-\gamma)$
$$-\cos(\alpha+\gamma).\cos(\alpha-\gamma)=0.$$

(19) $\cos(\delta-\alpha).\sin(\beta-\gamma)+\cos(\delta-\beta).\sin(\gamma-\alpha)$
$$-\cos(\delta-\gamma).\sin(\beta-\alpha)=0.$$

(20) $8\cos\frac{\theta+\phi+\chi}{2}.\cos\frac{\phi+\chi-\theta}{2}.\cos\frac{\chi+\theta-\phi}{2}.\cos\frac{\theta+\phi-\chi}{2}$
$$=\cos 2\theta+\cos 2\phi+\cos 2\chi+4\cos\theta.\cos\phi.\cos\chi+1.$$

CHAPTER XIII.**

On Angles Unlimited in Magnitude. II.

169. The words of the proofs (on pages 118, 119) of the 'A, B' formulæ apply to angles of any magnitude. The figures will be different for angles of different magnitude.

170. The figure for the '$A - B$' formulæ on page 119 is the same for all cases in which A and B are each less than 90°.

The figure given below is for the proof of the '$A + B$' formulæ, when, A and B being each less than 90°, their sum is greater than 90°.



The words of the proof are precisely those of page 118.

We may notice however that

$$\cos(A+B) = \frac{OM}{OP}\left[= \frac{-MO}{OP}\right] = \frac{-MK+OK}{OP} = \frac{OK}{OP} - \frac{MK}{OP},$$

and the rest follows as on page 118.

171. Thus we have proved that the 'A, B' formulæ are true provided A and B each lie between 0^0 and 90^0.

The student can prove them for any other values by drawing the proper figure.

The 'A, B' formulæ are therefore true for any values whatever of the angles A and B.

172. By the aid of the 'A, B' formulæ we can prove the formulæ of Art. 140.

Example. *Prove that* $\sin(90^0+A)=\cos A$.

$$\sin(90^0+A)=\sin 90^0\cos A+\cos 90^0\sin A,$$
$$=1\times\cos A+0\times\sin A,$$
$$=\cos A. \quad \text{Q. E. D.}$$

EXAMPLES. XLIV.

Draw the figures for the first four of the following examples.

(1) For the $(A+B)$ formulæ, when A is greater than 90^0 and $(A+B)$ less than 180^0.

(2) For the $(A-B)$ formulæ, when A and B each lie between 90^0 and 180^0.

(3) For the $(A+B)$ formulæ, when A lies between 90^0 and 180^0, and $(A+B)$ lies between 180^0 and 270^0.

(4) For the $(A-B)$ formulæ, when A lies between 180^0 and 270^0, and $(A-B)$ lies between 180^0 and A.

Deduce the six following formulæ from the 'A, B' formulæ.

(5) $\cos(90^0+A)=-\sin A$. (6) $\sin(90^0-A)=\cos A$.

(7) $\cos(90^0-A)=\sin A$. (8) $\sin(180^0-A)=\sin A$.

(9) $\cos(180^0-A)=-\cos A$. (10) $\sin(180^0+A)=-\sin A$.

(11) Assuming that the formula $\sin(A+B)=\sin A\,.\cos B+\cos A\,.\sin B$ is true for all values of A and B, deduce the rest of the 'A, B' formulæ by the aid of the results on p. 107.

173. We may also conversely prove that the 'A, B' formulæ are true for angles of any magnitude by the aid of the result of page 107. (This is a very convenient method of proving the 'A, B' formulæ to be true for all values of the angles.)

For, assuming that the 'A, B' formulæ are true for certain values of the angles A and B, we can show that they are true if either of the angles A or B be *increased* by 90°.

Example.

$$\begin{aligned}\sin(90^\circ+A+B) &= \cos(A+B) && \text{[p. 107.]}\\ &= \cos A \,.\, \cos B - \sin A \,.\, \sin B,\\ &= \sin(90^\circ+A) \,.\, \cos B - \{-\cos(90^\circ+A)\} \sin B,\\ &= \sin(90^\circ+A) \,.\, \cos B + \cos(90^\circ+A) \,.\, \sin B.\end{aligned}$$

Or, writing A' for $90^\circ+A$, we have

$$\sin(A'+B) = \sin A' \,.\, \cos B + \cos A' \,.\, \sin B.$$

We have proved (Art. 170) that the 'A, B' formulæ are true for all values of A and B between 0° and 90°. And therefore, by what we have said above, they are true for all values of A or B between 0° and 180°. And so on.

Therefore the 'A, B' formulæ are true for any values whatever of the angles A and B.

174. It follows that all formulæ deduced from the 'A, B' formulæ are true for angles of any magnitude whatever.

Thus the 'S, T' formulæ (page 126) are true for angles of any magnitude. Also the formulæ of the last Chapter for multiple angles, and all general formulæ in the Examples, are true for angles of any magnitude.

EXAMPLES. XLV.

(1) Deduce the values of $\sin 180^0$ and $\cos 180^0$ from those of the sine and cosine of 90^0.

(2) The angle A is greater than 180^0 and less than 270^0 and $\tan A = \frac{1}{2}$: find $\sin 2A$ and $\sin 3A$.

(3) The angle θ lies in the fourth quadrant and $\cos\theta = \frac{1}{4}$, find $\sin 2\theta$ and $\sin 3\theta$. Find also $\cos 3\theta$, and hence determine in which quadrant 3θ lies.

(4) Prove that the different values of θ which satisfy the equation $\cos p\theta + \cos q\theta = 0$, form two series in *A. P.* with common differences $\frac{2\pi}{p+q}$ and $\frac{2\pi}{p \sim q}$ respectively.

On Submultiple Angles.

175. We have now proved all the formulæ of the last two Chapters to be universally true.

We may expect therefore that any result, which can be obtained from these formulæ by algebraical transformation, will have a complete geometrical interpretation. [See p. 116.]

176. Since, $\cos A = 1 - 2\sin^2\frac{A}{2}$, [Art. 166.]

and, $\cos A = 2\cos^2\frac{A}{2} - 1$;

we have, $\sin^2\frac{A}{2} = \frac{1-\cos A}{2}$,

and, $\cos^2\frac{A}{2} = \frac{1+\cos A}{2}$.

Or,
$$\left.\begin{aligned}\sin\frac{A}{2} &= \pm\sqrt{\frac{1-\cos A}{2}}\\ \cos\frac{A}{2} &= \pm\sqrt{\frac{1+\cos A}{2}}\end{aligned}\right\}.$$

Thus, given the value of $\cos A$ (nothing else being known about the angle A), we get **two** values for $\sin\frac{A}{2}$, one positive and one negative, and **two** like values for $\cos\frac{A}{2}$.

177. *To prove geometrically that, given the value of* $\cos A$ (nothing else being known about the angle A), *there are* **two** *values each of* $\sin\frac{A}{2}$ *and of* $\cos\frac{A}{2}$.



Let α be the least positive angle which has the given cosine, and let ROP_1 and P_2OR in the figure each $= \alpha$. Then A is one of the angles described by the revolving line OP when, starting from OR, OP stops either in the position OP or in the position OP_2; i.e. any one of the angles

$$2n\pi \pm \alpha. \qquad \text{[Art. 147.]}$$

The angle described is (some multiple of *four* right angles $\pm\alpha$); a half of this is (some multiple of *two* right angles $\pm\frac{1}{2}\alpha$); so that a half of the angle whose cosine is equal to that of α, may be any one of the angles indicated by OR and the four dotted lines OP_3 and OP_5, or OP_4 and OP_6 in the figure.

And it will be seen that

$$\sin ROP_3 = \sin ROP_4 = -\sin ROP_5 = -\sin ROP_6.$$

Also, $\cos ROP_3 = -\cos ROP_4 = -\cos ROP_5 = \cos ROP_6$.

From these it is clear that there are **two** values of $\sin\frac{A}{2}$, equal in magnitude and opposite in sign. Also, that there are **two** like values for $\cos\frac{A}{2}$.

EXAMPLES. XLVI.

(1) When A lies between -180^0 and 180^0, prove that

$$\cos\frac{A}{2} = +\sqrt{\frac{1+\cos A}{2}}.$$

(2) When A lies between 180^0 and 540^0, prove that

$$\cos\frac{A}{2} = -\sqrt{\frac{1+\cos A}{2}}.$$

(3) Find $\sin\frac{A}{2}$ in terms of $\cos A$, when A lies between 180^0 and 360^0.

(4) Prove that when A lies between $(4n+1)\pi$ and $(4n+3)\pi$, n being a positive integer, $\cos\frac{A}{2} = -\sqrt{\frac{1+\cos A}{2}}$.

(5) Find $\sin\frac{A}{2}$ in terms of $\cos A$, when A lies between $4n\pi$ and $(4n+2)\pi$, where n is a positive integer.

178. Since, $2\sin\frac{A}{2}.\cos\frac{A}{2}=\sin A$, [Art. 166.]

and, $\sin^2\frac{A}{2}+\cos^2\frac{A}{2}=1$,

we obtain by addition and subtraction

$$\sin^2\frac{A}{2}+\cos^2\frac{A}{2}+2\sin\frac{A}{2}.\cos\frac{A}{2}=1+\sin A,$$

$$\sin^2\frac{A}{2}+\cos^2\frac{A}{2}-2\sin\frac{A}{2}.\cos\frac{A}{2}=1-\sin A.$$

Therefore $\left(\sin\frac{A}{2}+\cos\frac{A}{2}\right)^2=1+\sin A$,

and $\left(\sin\frac{A}{2}-\cos\frac{A}{2}\right)^2=1-\sin A$.

Whence,
$$\left.\begin{aligned}\sin\frac{A}{2}+\cos\frac{A}{2}&=\pm\sqrt{1+\sin A}\\ \sin\frac{A}{2}-\cos\frac{A}{2}&=\pm\sqrt{1-\sin A}\end{aligned}\right\}\quad\begin{aligned}&\text{.....(i),}\\&\text{.....(ii).}\end{aligned}$$

Adding we get, $2\sin\frac{A}{2}=\pm\sqrt{1+\sin A}\pm\sqrt{1-\sin A}$ (iii),

Subtracting we get, $2\cos\frac{A}{2}=\pm\sqrt{1+\sin A}\mp\sqrt{1-\sin A}$ (iv).

Thus if we are given the value of $\sin A$, (nothing else being known about the angle A), we have **four** values for $\sin\frac{A}{2}$ and **four** values for $\cos\frac{A}{2}$.

179. *To prove Geometrically that, given the value of* $\sin A$ (nothing else being known about the angle A), *there are* **four** *values for* $\sin\frac{A}{2}$, *and* **four** *values for* $\cos\frac{A}{2}$.

Let α be the least positive angle which has the given sine, and let ROP_1, P_2OL in the figure each $=\alpha$.



Then A is one of the angles described by the revolving line OP, when, starting from OR, OP stops either in the position OP_1 or in the position OP_2.

The angle described is either (an *even* multiple of *two* right angles $+ \alpha$) or (an *odd* multiple of *two* right angles $- \alpha$); a half of this is either (an *even* multiple of one right angle $+ \frac{1}{2}\alpha$) or (an *odd* multiple of one right angle $- \frac{1}{2}\alpha$); so that a half of the angle whose sine is equal to that of α may be any one of the angles indicated by OR and the dotted lines OP_3 and OP_5, or OP_4 and OP_6 in the figure.

In the figure, $ROP_3 = P_4OU = LOP_5 = P_6OD$.

And $\sin\frac{A}{2}$ may have any one of the values $\sin ROP_3$, $\sin ROP_4$, $\sin ROP_5$, $\sin ROP_6$. These values are all different, and are those given by the solution (iii).

Hence, there are **four** values for $\sin\frac{A}{2}$, of the nature indicated by the solution (iii); also **four** values for $\cos\frac{A}{2}$ (iv).

180. If we know the magnitude of A, we can decide which sign to take in the formulæ

$$\sin\frac{A}{2}+\cos\frac{A}{2}=\pm\sqrt{1+\sin A}\ldots\ldots\ldots\ldots\text{(i)},$$

$$\sin\frac{A}{2}-\cos\frac{A}{2}=\pm\sqrt{1-\sin A}\ldots\ldots\ldots\ldots\text{(ii)}.$$

Example. When $\frac{A}{2}$ lies between -45^0 and $+45^0$,

$\cos\frac{A}{2}$ is *greater* than $\sin\frac{A}{2}$ and is *positive.*

So that $\left(\sin\frac{A}{2}+\cos\frac{A}{2}\right)$ is *positive*, and $=+\sqrt{1+\sin A}$,

and $\left(\sin\frac{A}{2}-\cos\frac{A}{2}\right)$ is *negative*, and $=-\sqrt{1-\sin A}$.

When $\frac{A}{2}$ lies between $+45^0$ and $+135^0$,

$\sin\frac{A}{2}$ is greater than $\cos\frac{A}{2}$, and is positive.

So that $\left(\sin\frac{A}{2}+\cos\frac{A}{2}\right)$ and $\left(\sin\frac{A}{2}-\cos\frac{A}{2}\right)$ are both positive.

And so on.

The following diagram completes the above results.



181. Since $$\tan A = \frac{2\tan\frac{A}{2}}{1-\tan^2\frac{A}{2}},$$

$$\therefore \tan^2\frac{A}{2} + \frac{2\tan\frac{A}{2}}{\tan A} - 1 = 0,$$

whence $$\tan\frac{A}{2} = \frac{-1 \pm \sqrt{1+\tan^2 A}}{\tan A}.$$

Thus, given $\tan A$ we find two unequal values for $\tan\frac{A}{2}$ one positive and one negative.

182. The student will be able by the aid of the following figure to verify this result geometrically.



183. We may remark that in this figure P_3OP_5 and P_4OP_6 are straight lines at right angles to each other. So that $\tan P_3OR = -\cot P_4OR$; or, $\tan P_3OR \,.\, \tan P_4OR = -1$.

Hence, one value of $\tan\frac{A}{2}$ is the reciprocal of the other, and of opposite sign. So that there is always one *positive* value of $\tan\frac{A}{2}$, and one *negative;* one numerically *greater* than unity and the other numerically *less* than unity.

Example. *If* $A = 190^0$, *prove that*

$$\tan\frac{A}{2} = \frac{-1-\sqrt{1+\tan^2 A}}{\tan A}.$$

Now $\tan A = \tan 190^0$, which is positive; $\tan\frac{A}{2} = \tan 95^0$, which is negative. Also $\tan\frac{A}{2} = \frac{-1\pm\sqrt{1+\tan^2 A}}{\tan A}$; the negative value of which is $\frac{-1-\sqrt{1+\tan^2 A}}{\tan A}$.

EXAMPLES. XLVII.

(1) State the signs of $\left(\sin\frac{A}{2}+\cos\frac{A}{2}\right)$ and $\left(\sin\frac{A}{2}-\cos\frac{A}{2}\right)$ when $\frac{A}{2}$ has the following values:

(i) 22^0, (ii) 191^0, (iii) 290^0, (iv) 345^0,
(v) -22^0, (vi) -275^0, (vii) -470^0, (viii) 1000^0.

(2) Prove that the formulæ which give the values of $\sin\frac{A}{2}$ and of $\cos\frac{A}{2}$ in terms of $\sin A$, are unaltered when A has the values

(i) 92^0, 268^0, 900^0, $4n\pi+\frac{3}{4}\pi$, or $(4n+2)\pi-\frac{3}{4}\pi$.

(ii) 88^0, -88^0, 770^0, -770^0, or $4n\pi\pm\frac{\pi}{8}$.

(3) Find the values of (i) $\sin 9^0$, (ii) $\cos 9^0$, (iii) $\sin 81^0$, (iv) $\cos 189^0$, (v) $\tan 202\frac{1}{2}^0$, (vi) $\tan 97\frac{1}{2}^0$.

(4) If $A = 200^0$, prove that

(i) $2\sin\frac{A}{2} = +\sqrt{1+\sin A}+\sqrt{1-\sin A}$.

(ii) $\tan\frac{A}{2} = \frac{-(1+\sqrt{1+\tan^2 A})}{\tan A}$.

(5) If A lie between 270^0 and 360^0, prove that

(i) $2\sin\frac{A}{2} = \sqrt{1-\sin A}-\sqrt{1+\sin A}$.

(ii) $\tan\frac{A}{2} = -\cot A+\operatorname{cosec} A$.

(6) If A lie between 450^0 and 630^0 prove that

$$2\sin\frac{A}{2} = -\sqrt{1+\sin A}-\sqrt{1-\sin A}.$$

(7) Find the limits between which $\frac{A}{2}$ must lie when

$$2 \sin \frac{A}{2} = \sqrt{1+\sin A} - \sqrt{1-\sin A},$$

(8) Given that A lies between 450^0 and 630^0, prove that

$$2 \cos \frac{A}{2} = -\sqrt{1+\sin A} + \sqrt{1-\sin A}.$$

(9) If A lie between $n \times 360^0 - 90^0$ and $n \times 360^0 + 90^0$ where n is a positive integer, prove that

$$\tan \frac{A}{2} = \frac{-1+\sqrt{1+\tan^2 A}}{\tan A};$$

and that when A lies between $n \times 360^0 + 90^0$ and $n \times 360^0 + 270^0$, then

$$\tan \frac{A}{2} = \frac{-1-\sqrt{1+\tan^2 A}}{\tan A}.$$

(10) Prove geometrically that if we are given the value of $\sin A$, there are *three* different values for $\sin \frac{A}{3}$, and *six* different values for $\cos \frac{A}{3}$.

(11) Prove geometrically that, if we are given the value of $\cos A$, there are three different values for $\cos \frac{A}{3}$, and six different values for $\sin \frac{A}{3}$.

(12) Prove that if we are given the value of $\tan A$ there are three different values for $\tan \frac{A}{3}$.

(13) Given the value of $\tan A$, prove that there are *four* values each for $\sin \frac{A}{2}$ and $\cos \frac{A}{2}$.

(14) Given the value of $\sin A$, prove that there are *two* values for $\tan \frac{A}{2}$

184. It is important if possible, in solving Trigonometrical equations, to avoid squaring both sides of the equation.

Example. *Solve* $\cos\theta = k\sin\theta$.

If we square both sides of the equation we get

$$\cos^2\theta = k^2\sin^2\theta = k^2(1-\cos^2\theta).$$

$$\therefore\ \cos^2\theta = \frac{k^2}{1+k^2},\ \text{or}\ \cos\theta = \pm\frac{k}{\sqrt{1+k^2}}.$$

Now if a be the least angle such that $\cos\theta = \dfrac{k}{\sqrt{1+k^2}}$,

then the above gives us $\theta = n\pi \pm a$ .....(i).

But the equation may be written $\cot\theta = k$,

whence $\theta = n\pi + a$ .....(ii).

(ii) is the complete solution of the proposed equation, while (i) is in fact the solution of both $\cos\theta = k\sin\theta$ and also of $\cos\theta = -k\sin\theta$. So that by squaring both sides of the equation we obtain solutions which do not belong to the given equation.

185. We can often avoid squaring by the use of a **Subsidiary** Angle.

Example. *Solve* $a\cos\theta + b\sin\theta = 1$.

That is $$a\left(\cos\theta + \frac{b}{a}\sin\theta\right) = 1.$$

Find in the tables the angle whose tangent is $\dfrac{b}{a}$; let it be a.

Then $\dfrac{b}{a} = \tan a$; and the equation becomes

$$a(\cos\theta + \tan a\,.\,\sin\theta) = 1,$$

or $$a\left(\frac{\cos\theta\,.\,\cos a + \sin\theta\,.\,\sin a}{\cos a}\right) = 1,$$

or $$\cos(\theta - a) = \frac{\cos a}{a}.$$

Find from the Tables the value of $\cos a$. Next find from the Tables the magnitude of the angle β whose cosine $= \frac{\cos a}{a}$, and we get $\cos(\theta - a) = \cos\beta$;

$$\therefore \theta - a = 2n\pi \pm \beta,$$

or $$\theta = a + 2n\pi \pm \beta.$$

186. DEF. When an angle a is introduced to facilitate calculation it is called a **Subsidiary Angle.**

EXAMPLES. XLVIII.

Solve the following equations.

(1) $2\sin\theta + 2\cos\theta = \sqrt{2}$. (2) $\sin\theta + \sqrt{3}\cdot\cos\theta = 1$.

(3) $\sqrt{2}\sin\theta + \sqrt{2}\cos\theta = \sqrt{3}$. (4) $\sin\theta - \cos\theta = 1$.

(5) $\sin\theta + \cos\theta = 1$. (6) $\sqrt{3}\sin\theta - \cos\theta - \sqrt{2} = 0$.

(7) $2\sin x + 5\cos x = 2$. [$2{\cdot}5 = \tan 68^0\,12'$]

(8) $3\cos x - 8\sin x = 3$. [$2{\cdot}6 = \tan 69^0\,26'\,30''$]

(9) $4\sin x - 15\cos x = 4$. [$3{\cdot}75 = \tan 75^0\,4'$]

(10) $\cos(a + x) = \sin(a + x) + \sqrt{2}\cos\beta$.

*ON THE INVERSE NOTATION.

187. The equation $\sin\theta = a$ means that θ is an angle whose sine is a.

$\theta = \sin^{-1} a$ is a convenient way of writing the same equation.

Thus $\sin^{-1}\alpha$ (is an angle, and) is an abbreviation for an angle whose sine is α.

Example 1. *Show that* 30^0 *is* one *value of* $\sin^{-1}\frac{1}{2}$.

We know that $\sin 30^0 = \frac{1}{2}$. Therefore 30^0 is an angle whose sine is $\frac{1}{2}$, or, $30^0 = \sin^{-1}\frac{1}{2}$.

Example 2. *Prove that* 45^0 *is one value of* $\tan^{-1}\frac{1}{2}+\tan^{-1}\frac{1}{3}$

$\tan^{-1}\frac{1}{2}$ is one of the angles whose tangent is $\frac{1}{2}$.

Let $\alpha=\tan^{-1}\frac{1}{2}$, so that, $\tan\alpha=\frac{1}{2}$.

Let $\beta=\tan^{-1}\frac{1}{3}$, so that, $\tan\beta=\frac{1}{3}$.

We have to prove that 45^0 is one value of $\alpha+\beta$.

$$\tan(\alpha+\beta)=\frac{\tan\alpha+\tan\beta}{1-\tan\alpha\,.\,\tan\beta}=\frac{\frac{1}{2}+\frac{1}{3}}{1-\frac{1}{2}\,.\,\frac{1}{3}}=1.$$

But $\tan 45^0=1$.

Therefore 45^0 is one value of $\alpha+\beta$, i.e. of $\tan^{-1}\frac{1}{2}+\tan^{-1}\frac{1}{3}$.

* EXAMPLES. XLIX.

Prove that the following statements are true when we take for $\sin^{-1}\alpha$, etc. their least *positive* value.

(1) $\sin^{-1}\frac{3}{5}=\cos^{-1}\frac{4}{5}=\tan^{-1}\frac{3}{4}$.

(2) $\sin^{-1}\frac{1}{2}=\cos^{-1}\frac{\sqrt{3}}{2}=\cot^{-1}\sqrt{3}$.

(3) $\sin^{-1}\alpha=\cos^{-1}\sqrt{1-\alpha^2}=\tan^{-1}\frac{\alpha}{\sqrt{1-\alpha^2}}$.

(4) If $\alpha=\sin^{-1}\frac{3}{5}$ and $\beta=\cos^{-1}\frac{3}{5}$, then $\alpha+\beta=\frac{\pi}{2}$.

(5) If $A=\sin^{-1}\alpha$ and $B=\cos^{-1}\alpha$, then $A+B=90^0$.

(6) $\tan^{-1}\frac{5}{7}+\tan^{-1}\frac{1}{3}=\frac{\pi}{4}$. (7) $\tan^{-1}\frac{2}{11}+2\tan^{-1}\frac{1}{7}=\tan^{-1}\frac{1}{2}$.

(8) $\tan^{-1}m_1+\tan^{-1}m_2=\tan^{-1}\frac{m_1+m_2}{1-m_1m_2}$.

(9) $\sin(2\sin^{-1}\alpha)=2\alpha\sqrt{1-\alpha^2}$.

(10) $2\cos^{-1}\alpha=\cos^{-1}(2\alpha^2-1)$.

(11) $\cos^{-1}\frac{1}{2}+2\sin^{-1}\frac{1}{2}=120^0$. (12) $2\sin^{-1}\frac{4}{5}-\sin^{-1}2\frac{4}{5}=2\cos^{-1}2\frac{4}{5}$.

(13) $2\tan^{-1}(\cos 2\alpha)=\tan^{-1}\left(\frac{\cot^2\alpha-\tan^2\alpha}{2}\right)$.

(14) $\tan^{-1}x+\tan^{-1}y+\tan^{-1}\frac{1-x-y-xy}{1+x+y-xy}=\frac{\pi}{4}$.

(15) $4\tan^{-1}\frac{1}{5}-\tan^{-1}2\frac{1}{49}=\frac{\pi}{4}$.

(16) $\sin^{-1}\frac{4}{5}+\sin^{-1}\frac{8}{17}+\sin^{-1}\frac{13}{85}=\frac{\pi}{2}$.

(17) $\tan^{-1}\sqrt{5}\,(2-\sqrt{3})-\cot^{-1}\sqrt{5}\,(2+\sqrt{3})=\cot^{-1}\sqrt{5}$.

(18) If $\sin^{-1}m+\sin^{-1}n=\frac{\pi}{2}$,

prove that $m\sqrt{1-n^2}+n\sqrt{1-m^2}=1$.

188. The student must notice carefully that such a statement as $sin^{-1}\frac{1}{2}=cos^{-1}\frac{\sqrt{3}}{2}$ is not an identity.

For $\sin^{-1}\frac{1}{2}$ is one of the values of $n\pi+(-1)^n 30°$,

and $\cos^{-1}\frac{\sqrt{3}}{2}$ is one of the values of $2n\pi\pm 30°$.

Thus $150°=\sin^{-1}\frac{1}{2}$, but $150°$ is not $=\cos^{-1}\frac{\sqrt{3}}{2}$.

**MISCELLANEOUS EXAMPLES. L.

(1) Prove that $\tan^{-1}\frac{1}{1+a}+\tan^{-1}\frac{1}{1-a}+\tan^{-1}\frac{2}{a^2}=n\pi$.

(2) Prove that $\tan^{-1}\frac{a-1}{a}+\tan^{-1}\frac{1}{2a-1}=n\pi+\frac{\pi}{4}$.

(3) Prove that
$\sin^{-1}x-\sin^{-1}y=\cos^{-1}\{xy\pm\sqrt{1-x^2-y^2+x^2y^2}\}$.

(4) If $\tan^{-1}\frac{x+1}{x+2}+\tan^{-1}\frac{x-1}{x-2}=\frac{\pi}{4}$, show that $x^2=\frac{1}{2}$.

(5) Prove that $\tan^{-1}a+\cot^{-1}a=(2n+1)\frac{\pi}{2}$.

(6) If $\tan^{-1}\alpha+\tan^{-1}\beta+\tan^{-1}\gamma=\pi$,

prove that $\alpha+\beta+\gamma=\alpha\beta\gamma$.

(7) Solve the equation $\tan^{-1}\frac{1}{x-1}-\tan^{-1}\frac{1}{x+1}=\frac{\pi}{12}$.

(8) Solve the equation

$$\tan^{-1}(x+1)-\tan^{-1}(x-1)=\cot^{-1}(x^2-1).$$

(9) Solve the equation $\sin^{-1}\frac{2x}{1+x^2}+\tan^{-1}\frac{2x}{1-x^2}=\frac{\pi}{2}$.

(10) Prove that

$$\tan^{-1}\frac{a-\sqrt{a^2-4}}{2\sqrt{a+1}}+\tan^{-1}\frac{1}{\sqrt{a+1}}+\tan^{-1}\frac{a+\sqrt{a^2-4}}{2\sqrt{a+1}}=n\pi+\frac{\pi}{2}$$

(11) If a be positive and less than unity, and if α be the least value of $\sin^{-1}a$, then

$$\sin^{-1}a+\cos^{-1}a=n\pi+(-1)^n\alpha\pm\left(\frac{\pi}{2}-\alpha\right).$$

(12) Prove that $\tan A$ and $\sin 2A$ have always the same sign.

Solve the six following equations.

(13) $\cos A+\cos 3A+\cos 5A=0.$

(14) $\sin 5\theta+\sin 3\theta+\sin\theta=3-4\sin^2\theta.$

(15) $2\sin^2 3A+\sin^2 6A=2.$

(16) $a(\cos 2x-1)+2b(\cos x+1)=0.$

(17) $\sin(m+n)\theta+\sin 2m\theta+\sin(m-n)\theta=0.$

(18)
$$\left.\begin{aligned}&\sin\{\pi x(x+y)\}+\sin\{\pi y(x+y)\}=0,\\&\sin\pi x^2+\sin\pi y^2=0.\end{aligned}\right\}$$

(19) Trace the changes in the sign and magnitude of the following expressions, as θ changes from 0 to π.

(i) $2\sin\theta\,.\cos\theta.$ (ii) $\cos^2\theta-\sin^2\theta.$

(iii) $\sin 3\theta.$ (iv) $\cot 2A.$

(v) $\sin(\theta+\alpha).$ (vi) $\cos(2\theta-\alpha).$

(20) Explain why the equations

$$\theta=n\pi+(-1)^n\alpha \quad\text{and}\quad \frac{\pi}{2}-\theta=2n\pi\pm\left(\frac{\pi}{2}-\alpha\right)$$

have exactly the same series of solutions.

(21) Explain why exactly the same series of angles are given by the two equations $\theta+\frac{\pi}{4}=n\pi+(-1)^n\frac{\pi}{6}$, and $\theta-\frac{\pi}{4}=2n\pi\pm\frac{\pi}{3}$.

CHAPTER XIV.

On Logarithms.

189. In Algebra it is explained

(i) that the *multiplication* of different powers of the same quantity is effected by *adding* the *indices* of those powers;

(ii) that *division* is effected by *subtracting* the *indices;*

(iii) that *involution* and *evolution* are respectively effected by the *multiplication* and *division* of the *indices.*

Example 1. If $m=a^h$, $n=a^k$,

then
$$m \times n = a^h \times a^k = a^{h+k} \quad \text{.....(i)},$$
$$m \div n = a^h \div a^k = a^{h-k} \quad \text{.....(ii)},$$
$$\left.\begin{aligned} m^3 &= (a^h)^3 = a^{3h}, \\ \sqrt[4]{m} &= m^{\frac{1}{4}} = (a^h)^{\frac{1}{4}} = a^{\frac{h}{4}} \end{aligned}\right\} \quad \text{.....(iii).}$$

Example 2. If $347 = 10^{2\cdot5403295}$ * and $461 = 10^{2\cdot6637009}$, prove that
$$347 \times 461 = 10^{5\cdot2040304}.$$

We have
$$\begin{aligned} 347 \times 461 &= 10^{2\cdot5403295} \times 10^{2\cdot6637009} \\ &= 10^{2\cdot5403295 + 2\cdot6637009} \\ &= 10^{5\cdot2040304}. \end{aligned}$$
Q. E. D.

* The number 347 lies between 100 and 1000, i. e. between 10^2 and 10^3. Hence, if there is a power of 10 which is equal to 347, its index must be greater than 2 and less than 3, i. e. equal to $2+$a fraction.

EXAMPLES. LI.

(1) If $m=a^h$, $n=a^k$, express in terms of a, h and k,

(i) $m^2 \times n^3$. (ii) $m^4 \div n^5$. (iii) $\sqrt[3]{m^4 \times n^5}$. (iv) $\{\sqrt[4]{m^5 \times n^3}\}^2$.

(2) If $453=10^{2\cdot6560982}$ and $650=10^{2\cdot8129134}$, find the indices of the powers of 10 which are equal to

(i) 453×650. (ii) $(453)^4$. (iii) $650^3 \times 453^2$. (iv) $\sqrt[3]{453}$.

(v) $\sqrt{453} \times \sqrt[6]{650}$. (vi) $\sqrt[5]{453} \times (650)^3$. (vii) $\sqrt{453 \times 650}$.

(3) Express in powers of 2 the numbers, 8, 32, $\frac{1}{2}$, $\frac{1}{16}$, ·125, 128.

(4) Express in powers of 3 the numbers, 9, 81, $\frac{1}{3}$, $\frac{1}{27}$, $\cdot\dot{1}$, $\frac{1}{81}$.

190. Suppose that some convenient number (such as 10) having been chosen, we are given a **list** of the **indices** of the powers of that number, which are equivalent to every whole number from 1 up to 100000

Such a list could be used to shorten Arithmetical calculations.

Example 1. Multiply 3759 by 4781 and divide the result by 2690

Looking in our *list* we should find $3759=10^{3\cdot5750723}$, $4781=10^{3\cdot6795187}$, $2690=10^{3\cdot4297523}$.

Therefore $3759 \times 4781 \div 2690=10^{3\cdot5750723} \times 10^{3\cdot6795187} \div 10^{3\cdot4297523}$
$=10^{3\cdot5750723+3\cdot6795187-3\cdot4297523}=10^{3\cdot8248387}$.

The list will give us that $10^{3\cdot8248387}=6680\cdot9$.

Therefore the answer correct to five significant figures is 6680·9.

Example 2. Simplify $3^6 \times 2^{10} \div \sqrt[3]{17601}$.

The *list* gives $2=10^{\cdot3010300}$, $3=10^{\cdot4771213}$ and $17601=10^{4\cdot2455373}$.

Thus $3^6 \times 2^{10} \div \sqrt[3]{17601} = (10^{\cdot 4771213})^6 \times (10^{\cdot 3010300})^{10} \div (10^{4 \cdot 2455373})^{\frac{1}{3}}$

$= 10^{2 \cdot 8627278} \times 10^{3 \cdot 0103000} \div 10^{1 \cdot 4151791}$

$= 10^{2 \cdot 8627278 + 3 \cdot 0103000 - 1 \cdot 4151791}$

$= 10^{4 \cdot 4578487}$.

And from our list we find $10^{4 \cdot 4578487} = 28697$, nearly.

EXAMPLES. LII.

Given that $2 = 10^{\cdot 3010300}$, $3 = 10^{\cdot 4771213}$ and $7 = 10^{\cdot 8450980}$, find the indices of the powers of 10 equivalent to the quantities in the first 6 examples.

(1) 2^2, 3^2, 2^3, 2×3, 2^4, 7^2. (2) 14, 16, 18, 24, 27, 42.

(3) 10, 5, 15, 25, 30, 35. (4) 36, 40, 48, 50, 200, 1000.

(5) $3^{10} \times 7^{10} \div 2^{20}$, $2^{12} \times 3^{20} \div 7^{11}$.

(6) $\sqrt[3]{21} \times \sqrt[4]{18}$, $\sqrt[3]{\cdot 49 \times 4^5} \times \sqrt[3]{3^4 \times 2^{10}}$.

(7) Find approximately the numerical value of $\sqrt[19]{42}$ having given that $10^{\cdot 1623249} = 1 \cdot 4532$ nearly.

(8) Find approximately the numerical value of $\sqrt[3]{(42)^4} \times \sqrt[4]{(42)^3}$ having given that $10^{3 \cdot 38177} = 2408 \cdot 6$.

(9) Find the value (i) of $\sqrt[3]{6} \times \sqrt[4]{7} \times \sqrt[5]{9}$, (ii) of $\sqrt[10]{2} \times 3^{-\frac{5}{4}} \times 7^{\frac{7}{11}}$ having given that $10^{\cdot 6615067} = 4 \cdot 5868$ and $10^{-\cdot 0285094} = \cdot 93646$.

(10) Find the value of $(67 \cdot 21)^{\frac{3}{5}} \times (49 \cdot 62)^{\frac{1}{5}} \times (3 \cdot 971)^{-\frac{7}{5}}$ having given that $67 \cdot 21 = 10^{1 \cdot 8274339}$, $49 \cdot 62 = 10^{1 \cdot 6956568}$, $3 \cdot 971 = 10^{\cdot 5988999}$ and $10^{\cdot 5971310} = 3 \cdot 9549$.

(11) Find the area of a square field whose side is 640·12 feet; having given that $640 \cdot 12 = 10^{2 \cdot 8062614}$ and that $10^{5 \cdot 6125228} = 40975 \cdot 3$.

(12) Find the edge of a solid cube which contains 42601 cubic inches; having given $42601 = 10^{4 \cdot 6294198}$ and $10^{1 \cdot 5431399} = 34 \cdot 925$.

(13) Find the edge of a solid cube which contains 34·701 cubic inches; having given that $34 \cdot 701 = 10^{1 \cdot 5403420}$, and $10^{\cdot 5134473} = 3 \cdot 2617$.

(14) Find the volume of the cube the length of one of whose edges is 47·931 yards; having given that $47 \cdot 931 = 10^{1 \cdot 6806165}$ and that $10^{5 \cdot 0418495} = 110115$.

191. The powers of any other number than 10 might be used in the manner explained above, but 10 is the most *convenient* number, as will presently appear.

192. This method, in which the *indices* of the powers of a certain fixed number (such as 10) are made use of, is called the *Method of Logarithms.*

Indices thus used are called **logarithms.**

The *fixed number* whose powers are used is called the **base.** Hence we have the following definition:

DEF. The **logarithm** of a number to a given *base,* is the **index** of that power of the base, which is equal to the given number.

Thus, if l be the logarithm of the number n to the base a, then $a^l = n$.

193. The **notation** used is $\log_a n = l$.

Here, $\log_a n$ is an abbreviation for the words 'the logarithm of the number n to the base a.' And this means, as we have explained above, 'the **index** of that power of a which is equal to the number n.'

Example 1. What is the logarithm of $a^{\frac{3}{2}}$ to the base a?

That is, what is the index of the power of a which is $a^{\frac{3}{2}}$?

The index is $\frac{3}{2}$; therefore $\frac{3}{2}$ is the required logarithm, or $\log_a a^{\frac{3}{2}} = \frac{3}{2}$.

Example 2. What is the logarithm of 32 to the base 2?

That is, what is the **index** of the power of 2 which is equal to 32?

Now $32 = 2^5$. $\therefore$ the required index is 5; or $\log_2 32 = 5$.

The use of Logarithms is based upon the following propositions:—

I. The logarithm of the product of two numbers is equal to the logarithm of one of the numbers + the logarithm of the other.

For, let $\log_a m = x$ and $\log_a n = y$,

then, $\log_a (m \times n) = \log_a (a^x \times a^y) = \log_a (a^{x+y}) = x + y$
$= \log_a m + \log_a n.$

II. The logarithm of the quotient of two numbers is the logarithm of the dividend − the logarithm of the divisor.

For, $\log_a \left(\frac{m}{n}\right) = \log_a \left(\frac{a^x}{a^y}\right) = \log_a (a^{x-y}) = x - y$ [as above]
$= \log_a m - \log_a n.$

III. The logarithm of a number raised to a power k is k times the logarithm of the number.

For, $\log_a (m^k) = \log_a \{(a^x)^k\} = \log_a (a^{kx}) = kx$
$= k$ times $\log_a m$.

Examples. Given

$\log_{10} 2 = \cdot 3010300$, $\log_{10} 3 = \cdot 4771213$, $\log_{10} 7 = \cdot 8450980$

find the values of the following:

(i) $\log_{10} 6 = \log_{10} (2 \times 3) = \log_{10} 2 + \log_{10} 3$ [by I.]
$= \cdot 3010300 + \cdot 4771213$
$= \cdot 7781513.$

(ii) $\log_{10} \frac{7}{3} = \log_{10} 7 - \log_{10} 3 = \cdot 8450980 - \cdot 4771213$ [by II.]
$= \cdot 3679767.$

(iii) $\log_{10} 3^5 = 5$ times $\log_{10} 3 = 5 \times \cdot 4771213$ [by III.]
$= 2 \cdot 3856065.$

(iv) $\log_{10} \sqrt[3]{\frac{3 \times 4}{7}} = \log_{10} \left(\frac{3 \times 4}{7}\right)^{\frac{1}{3}} = \frac{1}{3}$ of $\log_{10} \frac{3 \times 4}{7}$ [by III.]
$= \frac{1}{3}$ of $(\log 3 + \log 4 - \log 7)$ [by I. and II.]
$= \frac{1}{3}$ of $\{\cdot 4771213 +$ twice $\cdot 30103 - \cdot 8450980\}$
$= \frac{1}{3}$ of $\cdot 2340833 = \cdot 0780278.$

(v) $\log_{10} 5 = \log_{10} \frac{10}{2} = \log_{10} 10 - \log_{10} 2 = 1 - \cdot 3010300$
$= \cdot 6989700.$

EXAMPLES. LIII.

(1) Find the logarithms to the base a of a^3, $a^{1.6}$, $\sqrt[4]{a}$, $\sqrt[3]{a^2}$, $\frac{1}{a^{\frac{5}{2}}}$.

(2) Find the logarithms to the base 2 of 8, 64, $\frac{1}{2}$, ·125, ·015625, $\sqrt[3]{64}$.

(3) Find the logarithms to the base 3 of 9, 81, $\frac{1}{3}$, $\frac{1}{27}$, ·$\dot{1}$, $\frac{1}{81}$.

(4) Find the logarithms to base 4 of 8, $\sqrt[3]{16}$, $\sqrt[4]{\cdot 5}$, $\sqrt[3]{\cdot 015625}$.

(5) Find the value of

$\log_2 8$, $\log_2 \cdot 5$, $\log_3 243$, $\log_5 (\cdot 04)$, $\log_{10} 1000$, $\log_{10} \cdot 001$.

(6) Find the value of

$\log_a a^{\frac{4}{3}}$, $\log_b \sqrt[3]{b^2}$, $\log_8 2$, $\log_{27} 3$, $\log_{100} 10$.

Given that

$\log_{10} 2 = \cdot 3010300$, $\log_{10} 3 = \cdot 4771213$ and $\log_{10} 7 = \cdot 8450980$, find the values of

(7) $\log_{10} 6$, $\log_{10} 42$, $\log_{10} 16$. (8) $\log_{10} 49$, $\log_{10} 36$, $\log_{10} 63$.

(9) $\log_{10} 200$, $\log_{10} 600$, $\log_{10} 70$. (10) $\log_{10} 5$, $\log_{10} 3\cdot 3$, $\log_{10} 50$.

(11) $\log_{10} 35$, $\log_{10} 150$, $\log_{10} \cdot 2$. (12) $\log_{10} 3\cdot 5$, $\log_{10} 7\cdot 29$, $\log_{10} \cdot 081$.

(13) Given $\log_{10} 2$, $\log_{10} 3$, $\log_{10} 7$, find the value (i) of $\sqrt[3]{6} \times \sqrt[4]{7} \times \sqrt[5]{9}$, (ii) of $\sqrt[10]{2} \times 3^{-\frac{5}{4}} \times 7^{\frac{7}{11}}$

[$\cdot 6615067 = \log_{10} 4\cdot 5868$; $-\cdot 0285094 = \log_{10} \cdot 93646$].

(14) Prove that (i) $\log\{\sqrt[3]{2} \times \sqrt[4]{7} \div \sqrt[5]{9}\} = \frac{1}{3}\log 2 + \frac{1}{4}\log 7 - \frac{2}{5}\log 3$,

(ii) $\log\{\sqrt[10]{2} \times 3^{-\frac{5}{4}} \times 7^{\frac{7}{11}}\} = \frac{1}{10}\log 2 - \frac{5}{4}\log 3 + \frac{7}{11}\log 7$.

(15) If $\log_{10} a = 2\cdot 6560982$ and $\log_{10} b = 2\cdot 8129134$, show that

(i) $\log_{10} ab = 5\cdot 4690116$, (ii) $\log_{10} a^4 = 10\cdot 6243928$,

(iii) $\log_{10} a^2 b^3 = 13\cdot 7509366$, (iv) $\log_{10} \sqrt[3]{a} = \cdot 8853661$.

(v) $\log_{10} (a^3 b)^{\frac{1}{3}} = 1\cdot 7968680$, (vi) $\log_{10} a^{\frac{1}{2}} b^3 = 8\cdot 9699598$.

(16) Show (i) that $\log_{10} \sqrt[3]{21} \times \sqrt[4]{18} = \cdot 7545579$, (ii) that $\log_{10} \sqrt[7]{(\cdot 49 \times 4^5)} \times \sqrt[3]{(3^4 \times 2^{10})} = 2\cdot 989843$.

Common Logarithms.

194. That System of Logarithms whose base is 10, is called the Common System of Logarithms.

In speaking of logarithms hereafter, *common* logarithms are referred to unless the contrary is expressly stated.

195. We shall assume that a power of ten can be found which is practically equivalent to any number.

196. The indices of these powers of 10, *i.e.* the Common Logarithms, are in general *incommensurable* numbers.

Their value for every whole number, from 1 to 100000, has been calculated to 7 significant figures. Thus any calculation made with the aid of logarithms is as exact as the most carefully observed measurement (cf. Arts. 17, 216).

197. Now, the greater the index of any power of 10, the greater will be the numerical value of that power; and the less the index, the less will be the numerical value of the power.

Hence, if one number be less than another, the logarithm of the first will be less than the logarithm of the second.

But the student should notice that logarithms (or indices) are *not proportional* to the corresponding numbers.

Example. 1000 is less than 10000; and the logarithm to base 10 of the first is 3 and of the second is 4

But 1000, 10000, 3, 4 are not in proportion.

198. We know from Algebra that

$$1 = 10^0,$$

$10 = 10^1$ and that $\cdot 1 = \frac{1}{10} = 10^{-1}$

$100 = 10^2$ ..... $\cdot 01 = \frac{1}{100} = 10^{-2}$

$1000 = 10^3$ ..... $\cdot 001 = \frac{1}{1000} = 10^{-3}$

$10000 = 10^4$ ..... $\cdot 0001 = \frac{1}{10000} = 10^{-4}$

and so on.

Hence, the logarithm of 1 is 0.

The (common) logarithm of any number greater than 1 is *positive.*

The logarithm of any positive number less than 1 is *negative.*

199. We observe also

that the logarithm of any number between 1 and 10 is a positive decimal fraction;

that the logarithm of any number between 10 and 100, *i.e.* between 10^1 and 10^2, is of the form 1 + a decimal fraction;

that the logarithm of any number between 1000 and 10000, *i.e.* between 10^3 and 10^4, is of the form 3 + a decimal fraction;

and so on.

200. We observe also

that the logarithm of any number between 1 and ·1, *i.e.* between 10^0 and 10^{-1}, can be written in the form -1 + a decimal fraction;

that the logarithm of any number between ·1 and ·01, *i.e.* between 10^{-1} and 10^{-2}, can be written in the form -2 + a decimal fraction;

and so on.

Example 1. How many digits are there in the integral part of the number whose logarithm is 3·67192?

We know that $3 = \log 1000$ and $4 = \log 10000$;

$\therefore$ 3·67192 = log of some number between 1000 and 10000; that is, the integral part of the number has 4 digits.

Example 2. Given that $\log 3 = \cdot 4771213$, find the number of the digits in the integral part of 3^{20}.

Here, $\log (3^{20}) = 20$ times $\log 3 = 9 \cdot 542426$;

hence, $\log (3^{20})$ lies between 9 and 10;

therefore, as in Example 1, the number lies between 10^9 and 10^{10}; that is, its integral part has 10 digits.

Example 3. Supposing that the decimal part of the logarithm is to be kept positive, find the integral part of the logarithm of ·0001234.

This number is greater than ·0001 *i.e.* than 10^{-4} and less than ·001, *i.e.* than 10^{-3}.

Therefore its logarithm lies between -3 and -4, and therefore it is $-4 +$ a fraction; the integral part is therefore -4.

EXAMPLES. LIV.

Note. The *decimal* part of a logarithm is to be kept *positive.*

(1) Write down the integral part of the common logarithms of 17601, 361·1, 4·01, 723000, 29.

(2) Write down the integral part of the common logarithms of ·04, ·0000612, ·7963, ·001201. (See Note above.)

(3) Write down the integral part of the common logarithms of 7963, ·1, 2·61, 79·6341, 1·0006, ·00000079.

(4) How many digits are there in the integral part of the numbers whose common logarithms are respectively

3·461, ·3020300, 5·4712301, 2·6710100?

(5) Give the position of the first *significant* figure in the numbers whose logarithms are

$$-2+\cdot4612310,\quad -1+\cdot2793400,\quad -6+\cdot1763241.$$

(6) Give the position of the first *significant* figure in the numbers whose common logarithms are $4\cdot2990713$, $\cdot3040595$, $2\cdot5860244$, $-3+\cdot1760913$, $-1+\cdot3180633$, $\cdot4980347$.

(7) Given $\log_{10}2=\cdot30103$, find the number of digits in the integral part of 8^{10}, 2^{12}, 16^{20}, 2^{100}.

(8) Given that $\log 7=\cdot8450980$, find the number of digits in the integral part of 7^{10}, 49^6, $343^{\frac{100}{3}}$, $(\frac{10}{7})^{20}$, $(4\cdot9)^{12}$, $(3\cdot43)^{10}$.

(9) Find the position of the first significant figure in

$$\sqrt[10]{2},\ (\tfrac{1}{2})^{10},\ (\tfrac{10}{7})^{20},\ (\cdot02)^4,\ (\cdot49)^6.$$

(10) Find the position of the first significant figure in the numerical value of

$$20^7,\ (\cdot02)^7,\ (\cdot007)^2,\ (3\cdot43)^{\frac{1}{10}},\ (\cdot0343)^8,\ (\cdot0343)^{\frac{1}{10}}.$$

201. PROP. *To prove that if two numbers expressed in the decimal notation have the same digits (so that they differ only in the position of the decimal point), their logarithms to the base* 10 *will differ only by an integer.*

The decimal point in a number is moved by multiplying or dividing the number by some *integral* power of 10.

Let the numbers be m and n; then $m=n\times10^k$, where k is a whole number positive or negative; and

$$\begin{aligned}\log m &= \log(n\times10^k)=\log n+\log 10^k\\ &=\log n+k.\end{aligned}$$

Example 1. Let the numbers be $1\cdot2345$ and $1234\cdot5$.

Then, $1234\cdot5=1\cdot2345\times10^3$,

therefore $\log(1234\cdot5)=\log(1\cdot2345)+3$.

Example 2. Given $\log 1{\cdot}7692 = {\cdot}24776$, find

(i) $\log 17692$, (ii) $\log {\cdot}0017692$, (iii) $176{\cdot}92$.

Here, $\log 17692 = \log\{(1{\cdot}7692) \times 10^4\} = \log(1{\cdot}7692) + 4$
$= {\cdot}24776 + 4 = 4{\cdot}24776.$

$\log {\cdot}0017692 = \log\{(1{\cdot}7692) \times 10^{-3}\} = -3 + {\cdot}24776.$

$\log 176{\cdot}92 = \log\{(1{\cdot}7692) \times 10^2\} = 2{\cdot}24776.$

202. It is *convenient* to keep the *decimal part* of common logarithms always *positive*, because then the *decimal part* of the logarithms of any numbers expressed by the same digits will be *always* the same.

203. The decimal part of a logarithm is called the **mantissa.**

204. The integral part is called the **characteristic.**

205. The characteristic of a logarithm can be always obtained by the following rule, which is evident from page 167.

RULE. The characteristic of the logarithm of a number greater than unity is **one less** than the number of *integral* figures in that number.

The characteristic of a number less than unity is negative, and (when the number is expressed as a decimal,) is **one more** than the number of cyphers between the decimal point and the first significant figure to the right of the decimal point.

206. When the characteristic is negative, as for example in the logarithm $-3 + {\cdot}1760913$, the logarithm is abbreviated thus, $\bar{3}{\cdot}1760913$.

Example 1. The characteristics of 36741, $36{\cdot}741$, ${\cdot}0036741$, $3{\cdot}6741$ and ${\cdot}36741$ are respectively 4, 1, -3, 0, and -1.

Example 2. Given that the mantissa of the logarithm of 36741 is 5651510, we can at once write down the logarithm of any number whose digits are 36741.

Thus $\log 3674100 = 6{\cdot}5651510,$
$\log 36741 = 4{\cdot}5651510,$
$\log 367{\cdot}41 = 2{\cdot}5651510,$
$\log {\cdot}36741 = \bar{1}{\cdot}5651510,$
$\log {\cdot}00036741 = \bar{4}{\cdot}5651510,$

and so on.

207. In any set of tables of common logarithms the student will find the *mantissa only* corresponding to any set of digits.

It would obviously be superfluous to give the *characteristic.*

208. It is most important to remember to keep the **mantissa** always **positive.**

Example. Find the fifth root of ·00065061.

Here $\log_{10} {\cdot}00065061 = \bar{4}{\cdot}8133207,$

$\therefore \log_{10} ({\cdot}00065061)^{\frac{1}{5}} = \frac{1}{5}(\bar{4}{\cdot}8133207) = \frac{1}{5}(-4 + {\cdot}8133207)$
$= \frac{1}{5}(-5 + 1{\cdot}8133207) = -1 + {\cdot}3626641 = \bar{1}{\cdot}3626641,$

and $\bar{1}{\cdot}3626641 = \log {\cdot}23050,$

$\therefore$ the fifth root of ·00065061 = ·23050 nearly.

EXAMPLES. LV.

(1) Write down the logarithms of 776·43, 7·7643, ·00077643 and 776430. (The table gives opposite the numbers 77643, the figures 8901023.)

(2) Given that $\log_{10} 59082 = 4{\cdot}7714552$, write down the logarithms of 5908200, 5·9082, ·00059082, 590·82 and 5908·2.

(3) Find the fourth root of ·0059082, having given that

$$\log 5{\cdot}9082 = {\cdot}7714552\,;\quad 4{\cdot}4428638 = \log_{10} 27724.$$

(4) Find the product of ·00059082 and ·027724, having given that ·21431 = log 16380 (cf. Question 3).

(5) Find the 10th root of ·077643 (cf. Question 1), having given that ·8890102 = log 7·7448.

(6) Find the product of $(\cdot 27724)^2$ and ·077643. (See Questions 1 and 3; 7758288 = log 59680.)

**209. *To transform a system of logarithms having a given base, to another system with a different base.*

If we are given a list of logarithms calculated to a given base, we can deduce from it a list of logarithms calculated to any other base.

Let a be the given base; let b be any other base.

Let m be any number. Then the logarithm of m to the base a is in the given list. Let this logarithm be l. Then

$$m = a^l.$$

We wish to find the logarithm of m to the base b. Let it be x.

Then $m = b^x$. But $m = a^l$;

$$\therefore\ a^l = b^x\,;\ \text{or, } b = a^{\frac{l}{x}}\,;$$

or, $\frac{l}{x}$ is the logarithm of b to the base a.

Now the logarithm of b to the base a is given. For it is in the list of logarithms to the base a.

Thus, $$\frac{l}{x} = \log_a b\,;\ \text{or, } x = \frac{l}{\log_a b}\,;$$

or, $$\log_b m = \frac{\log_a m}{\log_a b}.$$

Hence, to calculate the logarithms of a series of numbers to a new base b, we have only to divide each of the logarithms of the numbers to any given base a, by a certain divisor, viz. $\log_a b$.

If then a list of logarithms to some base e can be made, we can deduce from it a list of common logarithms, by multiplying each logarithm in the given list by $\frac{1}{\log_e 10}$.

Example. Show how to transform logarithms, having 5 for base to logarithms having 125 for base.

Suppose $m = 5^l$, so that $l = \log_5 m$.

Now $125 = 5^3$, so that $3 = \log_5 125$,

and $m = 5^{3 \times \frac{l}{3}} = 125^{\frac{l}{3}}$, so that $\frac{l}{3} = \log_{125} m$.

Thus the logarithm of any number to base 5, divided by 3 (*i.e.* by $\log_5 125$), is the logarithm of the same number to the base 125.

**210. The student will find that the logarithms of numbers cannot be calculated to the base 10 directly.

They are first calculated to the base 2·7182818, etc., which is the sum of the series

$$1 + \frac{1}{1} + \frac{1}{1.2} + \frac{1}{1.2.3} + \frac{1}{1.2.3.4} + \text{etc. } ad\ inf.$$

This number is called e.

And the constant divisor in this case is $\log_e 10$,

and $$\frac{1}{\log_e 10} = \frac{1}{2{\cdot}30258509} = {\cdot}43429448, \text{ etc.}$$

When this constant divisor is transformed into a *multiplier*, this constant **multiplier** is called a **modulus.**

Example. Given that $\log_{10} 12 = 1{\cdot}0791812$, shew how to transform common logarithms to logarithms having 12 for base.

Here $$10^{1{\cdot}0791812} = 12;$$

$$\therefore 10 = 12^{\frac{1}{1{\cdot}0791812}} = 12^{\cdot 92662, \text{ etc.}}$$

$$\therefore 10^h = 12^{h \times \cdot 92662}.$$

EXAMPLES. LVI.

Show how to transform

(1) Logarithms with base 2 to logarithms with base 8.

(2) Logarithms with base 9 to logarithms with base 3.

(3) Common logarithms to logarithms with base 2.

(4) Logarithms with base 3 to common logs.

(5) Common logs to logarithms with base 3.

(6) Given $\log_{10} 2 = \cdot 3010300$, find $\log_2 10$.

(7) Given $\log_{10} 7 = \cdot 8450980$, find $\log_7 10$.

(8) Given $\log_{10} 2 = \cdot 3010300$, find $\log_8 10$ and $\log_{32} 10$.

**MISCELLANEOUS EXAMPLES. LVII.

(1) Find $\log_2 8$, $\log_5 1$, $\log_8 2$, $\log_7 1$, $\log_{32} 128$.

(2) Show that the logarithms of all except eight of the numbers from 1 to 30 inclusive, can be calculated in terms of log 2, log 3 and log 7.

(3) Show that the logarithms of the numbers 1 to 10 inclusive may be found in terms of the logarithms of 8, 14, 21.

(4) The mantissa of the log of 85762 is 9332949. Find the log of $\sqrt[17]{\cdot 0085762}$.

Find how many figures there are in $(85762)^{11}$, when it is multiplied out.

(5) Find the product of 47·609, 476·09, ·47609, ·000047609, having given that $\log 4{\cdot}7609 = \cdot 6776891$ and $\cdot 7107564 = \log 5{\cdot}1375$.

(6) What are the characteristics of the logarithm of 3742 to the bases 3, 6, 10 and 12 respectively.

(7) Having given that $\log 2 = \cdot 3010300$, $\log 3 = \cdot 4771213$ and $\log 7 = \cdot 8450980$, solve the following equations :

(i) $2^x \times 3^{4x} = 7^2$, (ii) $3^{2x} = 128 \times 7^{4-x}$,

(iii) $12^x = 49$, (iv) $2^{8x} = 21^{4-3x}$.

(8) Given $\log_{10} 7$, find $\log_7 490$.

(9) Given $\log_{10} 3$, find $\log_9 270$.

(10) Given $\log_{10} 2$, find $\log_5 10$.

(11) Given $\log_8 9 = a$, $\log_2 5 = b$, $\log_5 7 = c$; find the logs to base 10 of numbers 1 to 7 inclusive.

(12) How many positive integers are there whose logarithms to base 2 have 5 for a characteristic?

(13) If a be an integer, how many positive integers are there whose logs to base a have 10 for their characteristic?

(14) Given log 2 and log 7, find the eleventh root of $(39{\cdot}2)^2$.

log 1·9485 = ·289688.

(15) Prove that $7 \log \frac{15}{16} + 6 \log \frac{8}{9} + 5 \log \frac{2}{5} + \log \frac{32}{25} = \log 3$.

(16) Prove that

$$2 \log a + 2 \log a^2 + 2 \log a^3 \ldots + 2 \log a^n = n(n+1) \log a.$$

(17) Prove that $\log_a b \cdot \log_b a = 1$; and that $\log_a b \cdot \log_b c \cdot \log_c a = 1$.

(18) Prove that $\log_a r = \log_a b \cdot \log_b c \cdot \log_c d \ldots \log_q r$.

(19) Given that the integral part of $(3{\cdot}456)^{100000}$ contains 53856 digits, find log 345·6 correct to five places of decimals.

(20) Given that the integral part of $(3{\cdot}981)^{100000}$ contains sixty thousand digits, find log 39810 correct to five places of decimals.

(21) If the number of births in a year be $\frac{1}{48}$ of the population at the beginning of the year, and the number of deaths $\frac{1}{60}$, find in what time the population will be doubled.

Given log 2, log 3, and that log 241 = 2·3820170.

(22) Prove that

$$\log s + \log (s-a) - \log b - \log c = 2 \log \sqrt{\frac{s(s-a)}{bc}}.$$

(23) Prove that

$$\log (a^2 + x^2) + \log (a+x) + \log (a-x) = \log (a^4 - x^4).$$

(24) Prove that

$$\log \sin 4A = \log 4 + \log \sin A + \log \cos A + \log \cos 2A.$$

CHAPTER XV.

On the Use of Mathematical Tables.

213. The Logarithms referred to in this chapter, and in future throughout the book, are *Common* Logarithms.

214. Books of Mathematical Tables usually give an explanation of their own contents, but there are some points common to all such Tables which we proceed to explain.

215. The student will be supposed to have access to a book containing the following:

(i) A list of the logarithms of all whole numbers from 1 to 99999, calculated to seven significant figures;

(ii) A list of the numerical values, calculated to seven significant figures, of the Trigonometrical Ratios of all angles, between 0° and 90°, which differ by $1'$;

(iii) A list of the logarithms of these Ratios calculated to seven significant figures.

These will be found in Chambers' *Mathematical Tables*.

At the end of this Chapter we give the logarithms to five figures of all numbers from 100 to 999. This Table will be found useful in questions involving certain kinds of numerical calculations.

216. We have said that logarithms are in general incommensurable numbers. Their values can therefore only be given approximately.

If the value of any number is given to seven significant figures, then the error (i.e. the difference between the *given* value and the *exact* value of the number) is less than a millionth part of the number.

Example. 3·141592 is the value of π correct to seven significant figures. The *error* is less than ·000001; for π is less than 3·141593, and greater than 3·141592.

The ratio of ·000001 to 3·141592 is equal to 1 : 3141592. The ratio of ·000001 to π is less than this; i.e. much less than the ratio of one to one million.

217. An actual measurement of any kind must be made with the greatest care, with the most accurate instruments, by the most skilful observers, if it is to attain to anything like the accuracy represented by 'seven significant figures.'

Therefore the value of any quantity given correct to 'seven significant figures' is exact for all practical purposes.

218. We are given in the Tables the logarithms of all numbers from 1 to 99999; that is, of any number having *five* significant figures.

A Table consisting of the logarithms of all numbers from 1 to 9999999 (i.e. of any number having *seven* significant figures) would be *a hundred times as large.*

219. There is however a rule by which, if we are given a *complete* list of the logarithms of numbers having *five* significant figures, we can find the logarithms of numbers having *six* or *seven* significant figures.

Example. Suppose we require the logarithm of 4·804213.

From the Tables we find

$$\log 4{\cdot}8042 = {\cdot}6816211, \qquad \text{i.e. } 4{\cdot}8042 = 10^{{\cdot}6816211\cdots},$$
$$\log 4{\cdot}8043 = {\cdot}6816301, \qquad 4{\cdot}8043 = 10^{{\cdot}6816301\cdots}.$$

The number 4·804213 lies *between* the two numbers 4·8042, 4·8043 whose logarithms are found in the Tables, so that the required logarithm must lie *between* the two given logarithms.

Therefore we suppose that

$$\log 4{\cdot}804213 = {\cdot}6816211 + d, \qquad \text{i.e. } 4{\cdot}804213 = 10^{{\cdot}6816211\cdots + d}.$$

220. The **RULE** is as follows. **The differences between three numbers are proportional to the corresponding differences between the logarithms of those numbers,** provided that the *differences* between the numbers are *small* compared with the numbers.

Example. *Given* $\log 48042 = 4{\cdot}6816211$
$\log 48043 = 4{\cdot}6816301,$
find $\log 48042{\cdot}13.$

We may state the question thus.

Given $\log 48042 = 4{\cdot}6816211$
$\log (48042+1) = 4{\cdot}6816211 + {\cdot}0000890,$
find $\log (48042 + {\cdot}13).$
Let $\log (48042 + {\cdot}13) = 4{\cdot}6816211 + d.$

Thus corresponding to the **differences** 1 and ·13 in the numbers we have the **differences** ·0000090 and d in the logarithms.

But these differences are in proportion,

$$\therefore\ d = \frac{{\cdot}13}{1} \text{ times } {\cdot}000009$$
$$= \tfrac{13}{100} \text{ of } {\cdot}000009 = {\cdot}00000117\ldots;$$
$$\therefore \log 48042{\cdot}13 = 4{\cdot}6816211 + {\cdot}00000117$$
$$= 4{\cdot}68162227 = 4{\cdot}6816223 \text{ (to seven figures).}$$

221. We shall refer to the above rule as the **Rule of Proportional Differences.**

It is often called also 'The Principle of Proportional Parts.'

222. In Art. 197 we said that numbers are *not* proportional to their Logarithms. Hence the differences of numbers and the corresponding differences of their logarithms cannot be *exactly* in proportion. The rule is however true for all practical purposes. The proof of the rule belongs to a higher part of the subject than the present.

223. In the above example we said that

$$6{\cdot}68162227 = 6{\cdot}6816223\,;$$

and for this reason. We are *retaining* only *seven* significant figures in the decimal part of the logarithm.

If we put 6·6816222 for 6·68162227 the 'error' is greater than ·00000007.

If we put 6·6816223 for 6·68162227 the 'error' is less than ·00000003.

Thus the second error is less than the first.

In such a case, 1 must be added to the last digit which is retained, when the first digit which is neglected is 5 or greater than 5.

224. We give two more specimen examples.

Example 1. *Find the logarithm of* ·004804213.

We first find as before, by the rule of proportional differences, that

$$\log 4{\cdot}804213 = {\cdot}6816223$$

$$\therefore \log {\cdot}004804213 = \bar{3}{\cdot}6816223.$$

Example 2. *Find the number whose logarithm is* 2·5354291.

In the Table we find that

$$\cdot5354207 = \log 3{\cdot}4310 \quad \text{.....(i)},$$

and $$\cdot5354334 = \log 3{\cdot}4311 \quad \text{.....(ii)}.$$

Let $$\cdot5354291 = \log (3{\cdot}4310 + d) \quad \text{.....(iii)}.$$

Here we have three logarithms and three numbers.

Subtracting

·5354334	·5354291
·5354207	·5354207
·0000127	·0000084

The corresponding differences are ·0001 and d.

Hence,
$$d = \frac{\cdot0000084}{\cdot0000127} \text{ of } \cdot0001$$
$$= \tfrac{84}{127} \text{ of } \cdot0001$$
$$= \cdot0000661.$$

Therefore from (iii) $\cdot5354291 = \log (3{\cdot}4310 + \cdot000066\ldots)$
$$= \log 3{\cdot}431066.$$

Hence, $$2{\cdot}5354291 = \log 343{\cdot}1066,$$

or, the required number is 343·1066.

EXAMPLES. LVIII.

(1) Find log 7·65432, having given that

$$\log 7{\cdot}6543 = \cdot8839055,$$

$$\log 7{\cdot}6544 = \cdot8839112.$$

(2) Find log 564·123, having given that

$$\log 5{\cdot}6412 = \cdot7513715,$$

$$\log 5{\cdot}6413 = \cdot7513792.$$

(3) Find log ·0008736416, having given that

$$\log 8{\cdot}7364 = \cdot9413325,$$

$$\log 8{\cdot}7365 = \cdot9413375.$$

(4) Find log 6437125, having given that
log 6·4371 = ·8086903,
log 6·4372 = ·8086970.

(5) Find log 3·72456, having given that
log 37245 = 4·5710680,
log 37246 = 4·5710796.

(6) Find the number whose logarithm is ·5686760, having given that ·5686710 = log 3·7040,
·5686827 = log 3·7041.

(7) Find the number whose logarithm is 4·6602987, having given that ·6602962 = log 4·5740,
·6603057 = log 4·5741.

(8) Find the number whose logarithm is 6·3966938, having given that ·3966874 = log 2·4928,
·3967049 = log 2·4929.

(9) Find the number whose logarithm is $\bar{4}$·6431150, having given that ·6431071 = log 4·3965,
·6431170 = log 4·3966.

(10) Find the number whose logarithm is ·7550480, having given that 3·7550436 = log 5689·1,
2·7550512 = log 568·92.

225. The same Rule of Proportional Differences is used in the case of **angles** and their **Trigonometrical Ratios**; and therefore also in the case of **angles** and the **logarithms of their Ratios.**

Thus the (small) differences between three angles are assumed to be proportional to the corresponding differences between the sines of those three angles; also, proportional to the corresponding differences between the logarithms of the sines of those angles.

226. Sines and cosines are always less than unity, as also are the tangents of all angles between 0° and 45°.

The logarithms of these Ratios must therefore have *negative* characteristics.

To avoid the inconvenience of having to print these negative characteristics, the whole number 10 is added to each logarithm of the Trigonometrical Ratios, before it is set down in the Table.

The numbers thus recorded are called the **tabular logarithms** of the sine, cosine, etc., of an angle.

They are indicated by the letter 'L.'

Thus L *sin* 31° 15′, stands for the tabular logarithm of sin 31° 15′, and is equal to {log (sin 31° 15′) + 10}.

The words **logarithmic sine** are used as abbreviation for **tabular logarithm of the sine.**

Thus in the Tables we find

$$L \sin 31^\circ 15' = 9{\cdot}7149776.$$

Therefore $\log (\sin 31^\circ 15') = 9{\cdot}7149776 - 10 = \bar{1}{\cdot}7149776.$

Example 1. *Find* sin 31° 6′ 25″.

The Tables give sin 31° 6′ = ·5165333 .....(i),

sin 31° 7′ = ·5167824 .....(ii).

Let sin 31° 6′ 25″ = ·5165333 + d .....(iii).

Subtracting

$$\left.\begin{array}{l}\cdot 5167824\\ \cdot 5165333\end{array}\right\} \text{ The other difference is } d, \text{ and the corresponding differences in the angles are } 60'' \text{ and } 25'';$$
$$\cdot 0002491$$

$$\therefore\ d = \tfrac{25}{60} \text{ of } \cdot 0002491$$
$$= \cdot 0001038\ldots$$

Hence, from (iii) sin 31°6′25″ = ·5165333 + ·0001038 = ·5166371.

Example 2. Find the angle whose logarithmic cosine is 9·7858083.

The Table gives $9{\cdot}7857611 = L\cos 52^0\,22'$ .....(i),

$9{\cdot}7859249 = L\cos 52^0\,21'$ .....(ii).

The cosine diminishes as the angle increases. Hence corresponding to an increase in the angle there is a diminution of the cosine.

Hence, let $9{\cdot}7858083 = L\cos(52^0\,22' - D)$ .....(iii).

Subtracting the first tabular logarithm from the second the **difference** is ·0001638.

Subtracting the first tabular logarithm from the third, the **difference** is ·0000472.

Subtracting the first angle from the second, the **difference** is $-60''$.

Subtracting the first angle from the third, the **difference** is $-D$.

By the Rule these four **differences** are in proportion.

Therefore $D = \frac{472}{1638}$ of $60''$

$= 17{\cdot}3''$.

Hence $9{\cdot}7858083 = L\cos(52^0\,22' - 17'')$

$= L\cos 52^0\,21'\,43''$

EXAMPLES. LIX.

(1) Find $\sin 42^0\,21'\,30''$

having given that $\sin 42^0\,21' = {\cdot}6736577$

$\sin 42^0\,22' = {\cdot}6738727$.

(2) Find $\cos 47^0\,38'\,30''$

having given that $\cos 47^0\,38' = {\cdot}6738727$

$\cos 47^0\,39' = {\cdot}6736577$.

(3) Find $\cos 21^0\,27'\,45''$

having given that $\cos 21^0\,27' = {\cdot}9307370$

$\cos 21^0\,28' = {\cdot}9306306$.

(4) Find the angle whose sine is ·6666666
having given that ·6665325 = sin 41° 48′
·6667493 = sin 41° 49′.

(5) Find the angle whose cosine is ·3333333
having given that ·3332584 = cos 70° 32′
·3335326 = cos 70° 31′.

(6) Find the angle whose cosine is ·25
having given that ·2498167 = cos 75° 32′
·2500984 = cos 75° 31′.

(7) Find $L \sin 45^\circ 16' 30''$
having given that $L \sin 45^\circ 16' = 9{\cdot}8514969$
$L \sin 45^\circ 17' = 9{\cdot}8516220$.

(8) Find $L \tan 27^\circ 13' 45''$
having given that $L \tan 27^\circ 13' = 9{\cdot}7112148$
$L \tan 27^\circ 14' = 9{\cdot}7115254$.

(9) Find $L \cot 36^\circ 18' 20''$
having given that $L \cot 36^\circ 18' = 10{\cdot}1339650$
$L \cot 36^\circ 19' = 10{\cdot}1337003$.

(10) Find the angle whose Logarithmic tangent is 9·8464028,
having given that $9{\cdot}8463018 = L \tan 35^\circ 4'$
$9{\cdot}8465705 = L \tan 35^\circ 5'$.

(11) Find the angle whose Logarithmic cosine is 9·9448230,
having given that $9{\cdot}9447862 = L \cos 28^\circ 17'$
$9{\cdot}9448541 = L \cos 28^\circ 16'$.

(12) Find the angle whose Logarithmic cosecant is 10·4274623,
having given that $10{\cdot}4273638 = L \operatorname{cosec} 21^\circ 57'$
$10{\cdot}4276774 = L \operatorname{cosec} 21^\circ 56'$.

227. Problems in which each of the lines involved contains an *exact* number of feet, and each angle an *exact* number of degrees, **do not occur** in practical work.

As from time to time the skill of observers and of instrument-makers has increased, so also has the number of significant figures by which observations have been recorded.

Thus the want was felt of some method by which the labour involved in the multiplication and division of long numerical quantities could be avoided. At the end of the Seventeenth Century a celebrated Scotch mathematician, John Napier, Baron of Merchiston, proposed his method of 'Logarithms'; i.e. the method of representing numbers by **indices**; 'which, by reducing to a few days the labour of 'many months, doubles, as it were, the life of an Astronomer, 'besides freeing him from the errors and disgust inseparable 'from long calculations.' *Laplace.*

228. We shall now give a few examples of the **practical use** of logarithms.

Example 1. The sides containing the right angle C in a right angled triangle ABC contain 3456·4 ft. and 4543·5 feet respectively; find the angles of the triangle, and the length of the hypotenuse.



Let a, b, c be the lengths of the sides of the triangle opposite the angles A, B, C respectively.

Then, $a = 3456{\cdot}4$ feet, $b = 4543{\cdot}5$ feet,

$$\tan A = \frac{a}{b}.$$

Therefore $\log \tan A = \log \frac{a}{b} = \log a - \log b$

$$= \log 3456{\cdot}4 - \log 4543{\cdot}5.$$

In the Tables we find

$$\log 3456{\cdot}4 = 3{\cdot}5386240$$
$$\log 4543{\cdot}5 = 3{\cdot}6573905.$$
$$\therefore \log \tan A = 3{\cdot}5386240 - 3{\cdot}6573905$$
$$= \bar{1}{\cdot}8812335.$$
$$\therefore L \tan A = 9{\cdot}8812335.$$

In the Table we find

$$9{\cdot}8813144 = L \tan 37^0\ 16'$$
$$9{\cdot}8810522 = L \tan 37^0\ 15'.$$

Whence by the rule of Proportional Differences

$$9{\cdot}8812335 = L \tan 37^0\ 15'\ 42''.$$

Therefore $A = 37^0\ 15'\ 42''$
and $B = 90^0 - A = 52^0\ 44'\ 18''$ } .....(I).

To find c, we may take the square root of $a^2 + b^2$, or we may find it by the aid of logarithms thus:

$$\frac{c}{a} = \text{cosec } A = \text{cosec } 37^0\ 15'\ 42'',$$
$$\therefore \log c = \log a + \log \text{cosec } 37^0\ 15'\ 42''$$
$$= \log a + L \text{ cosec } 37^0\ 15'\ 42'' - 10$$
$$= 3{\cdot}5386240 + 10{\cdot}2179174 - 10$$
$$= 3{\cdot}7565414$$
$$= \log 5708{\cdot}8,$$
$$\therefore \text{the hypotenuse contains } 5708{\cdot}8 \text{ feet.}$$

Thus we have found the angles and the third side of the triangle.

229. There are some formulæ which are seldom used in practical work, because they are not adapted to logarithmic calculation. They are those in which powers of quantities are connected by the signs + or −.

Example. In the above example we might have found the length of the hypotenuse by means of the formula

$$c^2 = a^2 + b^2.$$

But we should have had to go through the process of calculating by multiplication the values of a^2 and b^2.

For this reason, a formula which consists entirely of **factors** is always preferred to one which consists of **terms**, when any of those terms contain any power of the quantities involved.

If in the above example, the lengths of the hypotenuse c and of one side a were given, then the formula

$$b^2 = c^2 - a^2 = (c-a)(c+a)$$

will give the length of b. For

$$\log b^2 = \log\{(c-a)(c+a)\},$$

or,

$$2\log b = \log(c-a) + \log(c+a).$$

And the values of $(c+a)$ and $(c-a)$ are easily written down from the given values of c and a.

EXAMPLES. LX.

In the following questions A, B, C are the angles of a right angled triangle of which C is a right angle, and a, b, c are the lengths of the sides opposite those angles respectively.

(1) Given that $a = 1046{\cdot}7$ yards, $c = 1856{\cdot}2$ yards, $C = 90^0$, find A.

$\log 1046{\cdot}7 = 3{\cdot}0198222$, $\log 1856{\cdot}2 = 3{\cdot}2686248$,
$L\sin 34^0\,19' = 9{\cdot}7510991$,
$L\sin 34^0\,20' = 9{\cdot}7512842$.

(2) Given that $a = 843{\cdot}2$ feet, $C = 90^0$, and $A = 34^0\,15'$; find c.

$\log 843{\cdot}2 = 2{\cdot}9259306$, $L\operatorname{cosec} 34^0\,15' = 10{\cdot}2496421$,

$\log 1{\cdot}4982 = {\cdot}17557$.

(3) Given that $a = 4845$ yards, $b = 4742$ yards, and $C = 90$, find A.

$\log 4845 = 3{\cdot}6852938$, $\log 4742 = 3{\cdot}6759615$,

$L\tan 45^0\,36' = 10{\cdot}0090965$, $L\tan 45^0\,37' = 10{\cdot}0093492$.

(4) Given that $c = 8762$ feet, $C = 90$, and $A = 37^0\,10'$, find a and b.

$\log 8762 = 3{\cdot}9426032$, $L\sin 37^0\,10' = 9{\cdot}7811344$,

$L\cos 37^0\,10' = 9{\cdot}9013938$, $\log 5{\cdot}2934 = {\cdot}72373$,

$\log 6{\cdot}9823 = {\cdot}843997$.

(5) Given that $b = 1694{\cdot}2$ chains, $C = 90^0$, and $A = 18^0\,47'$, find a.

$\log 1694{\cdot}2 = 3{\cdot}2289647$, $L\cot 18^0\,47' = 10{\cdot}4683893$,

$\log 5{\cdot}7620 = {\cdot}76057$.

(6) Given that $a = 1072$ chains, $c = 4849$ chains, and $C = 90^0$, find b.

$\log 5921 = 3{\cdot}7723951$, $\log 3777 = 3{\cdot}5771470$,

$\log 4{\cdot}729 = {\cdot}67477$.

(7) Given that $b = 841$ feet, $c = 3762$ feet, and $C = 90^0$, find a.

$\log 4603 = 3{\cdot}6630410$, $\log 2921 = 3{\cdot}4655316$,

$\log 3{\cdot}6668 = {\cdot}56428$.

(8) Given that $a = 7694{\cdot}5$ chains, $b = 8471$ chains, $C = 90^0$, find A and c.

$\log 7694{\cdot}5 = 3{\cdot}8861804$, $\log 8471 = 3{\cdot}9279347$,

$L\tan 42^0\,15' = 9{\cdot}95824$, $L\operatorname{cosec} 42^0\,15' = 10{\cdot}1723937$,

$\log 1{\cdot}1444 = {\cdot}05857$.

230. In the following examples the student must find the necessary logarithms from the Tables.

* MISCELLANEOUS EXAMPLES. LXI.

(1) A balloon is at a height of 2500 feet above a plain and its angle of elevation at a point in the plain is $40^0\,35'$. How far is the balloon from the point of observation?

(2) A tower standing on a horizontal plain subtends an angle of $37^0\,19'\,30''$ at a point in the plain distant 369·5 feet from the foot of the tower. Find the height of the tower.

(3) The shadow of a tower on a horizontal plain in the sunlight is observed to be 176·23 feet and the elevation of the sun at that moment is $33^0\,12'$. Find the height of the tower.

(4) From the top of a tower 163·5 feet high by the side of a river the angle of depression of a post on the opposite bank of the river is $29^0\,47'\,18''$. Find the distance of the post from the foot of the tower.

(5) Given $a=673{\cdot}12$, $b=415{\cdot}89$ chains, $C=90^0$, find A and B.

(6) Given $a=576{\cdot}12$, $c=873{\cdot}14$ chains, $C=90^0$, find b and A.

(7) From the top of a light-house 112·5 feet high, the angles of depression of two ships, when the line joining the ships points to the foot of the light-house, are $27^0\,18'$ and $20^0\,36'$ respectively. Find the distance between the ships.

(8) From the top of a cliff the angles of depression of the top and bottom of a light-house 97·25 feet high are observed to be $23^0\,17'$ and $24^0\,19'$ respectively. How much higher is the cliff than the light-house?

(9) Find the distance in space travelled in an hour, in consequence of the earth's rotation, by St Paul's cathedral. (Latitude of London $=51^0\,25'$, earth's diameter $=7914$ miles.)

(10) The angle of elevation of a balloon from a station due south of it is $47^0\,18'\,30''$, and from another station due west of the former and distant 671·38 feet from it the elevation is $41^0\,14'$. Find the height of the balloon.

On the four following pages are given a Table of the mantissæ of the logarithms of all numbers from 100 to 1000 correct to 5 significant figures.

By the aid of the principle of proportional parts this Table can be used to work correctly to 4 significant figures.

It is only in observations or measurements, made with the most elaborate care and under special circumstances, that a greater degree of accuracy is attained than that indicated by 4 significant figures [see Lock's *Arithmetic*, chapter VIII.]; hence the Table here given will be found sufficient for most numerical calculations in which Arithmetic only is involved.

Example i. *Find to three significant figures the length of the diagonal of a cube whose side contains* 14·7 *inches.*

Let x be the number of inches in the diagonal,

then $$x^2 = 3 \times (14{\cdot}7)^2,$$

$$\therefore \quad x = \sqrt{3} \times 14{\cdot}7\,;$$

$$\therefore \quad \log x = \tfrac{1}{2} \log 3 + \log 14{\cdot}7$$

$$= \tfrac{1}{2}\,(\cdot 47712) + 1{\cdot}16732 \text{ [from the Table]}$$

$$= \cdot 23856 + 1{\cdot}16732 = 1{\cdot}40588$$

$$= \log 25{\cdot}45 \text{ nearly [from the Table].}$$

Hence the diagonal is 25·45... inches.

Example ii. *Find the value of* $(43{\cdot}72)^{\frac{5}{9}}$.

$$\left.\begin{matrix}\log 4{\cdot}37 = \cdot 64048 \\ \log 4{\cdot}38 = \cdot 64147\end{matrix}\right\} \text{ difference for } \cdot 01 = \cdot 00099,$$

$$\therefore \quad \text{difference for } \cdot 002 = \cdot 00020\,;$$

$$\therefore \quad \log 43{\cdot}72 = 1{\cdot}64068\,;$$

$$\therefore \quad \log (43{\cdot}72)^{\frac{5}{9}} = \tfrac{5}{9} \times 1{\cdot}64068 = \cdot 9115\ldots$$

$$= \log 8{\cdot}156\ldots \text{ [from the Table].}$$

Therefore $(43{\cdot}72)^{\frac{5}{9}} = 8{\cdot}156\ldots$

TABLE OF THE LOGARITHMS OF ALL NUMBERS FROM 100 TO 1000

No.	Log.	No.	Log.	No.	Log	No.	Log.	No.	Log.
100	00000	143	15534	186	26951	229	35983	272	43457
101	00432	144	15836	187	27184	230	36172	273	43616
102	00860	145	16137	188	27416	231	36361	274	43775
103	01284	146	16435	189	27646	232	36549	275	43933
104	01703	147	16732	190	27875	233	36736	276	44091
105	02119	148	17026	191	28103	234	36922	277	44248
106	02531	149	17319	192	28330	235	37107	278	44404
107	02938	150	17609	193	28556	236	37291	279	44560
108	03342	151	17898	194	28780	237	37475	280	44716
109	03743	152	18184	195	29003	238	37658	281	44870
110	04139	153	18469	196	29226	239	37840	282	45025
111	04532	154	18752	197	29447	240	38021	283	45179
112	04922	155	19033	198	29667	241	38202	284	45332
113	05308	156	19312	199	29885	242	38382	285	45484
114	05690	157	19590	200	30103	243	38561	286	45637
115	06070	158	19866	201	30320	244	38739	287	45788
116	06446	159	20140	202	30535	245	38917	288	45939
117	06819	160	20412	203	30750	246	39094	289	46090
118	07188	161	20683	204	30963	247	39270	290	46240
119	07555	162	20951	205	31175	248	39445	291	46389
120	07918	163	21219	206	31387	249	39620	292	46538
121	08279	164	21484	207	31597	250	39795	293	46687
122	08636	165	21748	208	31806	251	39967	294	46835
123	08991	166	22011	209	32015	252	40140	295	46982
124	09342	167	22272	210	32222	253	40312	296	47129
125	09691	168	22531	211	32428	254	40483	297	47276
126	10037	169	22789	212	32634	255	40654	298	47422
127	10380	170	23045	213	32838	256	40824	299	47567
128	10721	171	23300	214	33041	257	40993	300	47712
129	11059	172	23553	215	33244	258	41162	301	47857
130	11394	173	23805	216	33446	259	41330	302	48001
131	11727	174	24055	217	33646	260	41497	303	48144
132	12057	175	24304	218	33846	261	41664	304	48287
133	12385	176	24551	219	34044	262	41830	305	48430
134	12710	177	24797	220	34242	263	41996	306	48572
135	13033	178	25042	221	34439	264	42160	307	48714
136	13354	179	25285	222	34635	265	42325	308	48855
137	13672	180	25527	223	34830	266	42488	309	48996
138	13988	181	25768	224	35025	267	42651	310	49136
139	14301	182	26007	225	35218	268	42813	311	49276
140	14613	183	26245	226	35411	269	42975	312	49415
141	14921	184	26482	227	35602	270	43136	313	49554
142	15229	185	26717	228	35793	271	43297	314	49693

No.	Log.	No.	Log.	No.	Log.	No.	Log.	No.	Log.
315	49831	361	55751	407	60959	453	65610	499	69810
316	49969	362	55871	408	61066	454	65705	500	69897
317	50106	363	55991	409	61172	455	65801	501	69984
318	50243	364	56110	410	61278	456	65896	502	70070
319	50379	365	56229	411	61384	457	65992	503	70157
320	50515	366	56348	412	61470	458	66087	504	70243
321	50651	367	56467	413	61595	459	66181	505	70329
322	50786	368	56585	414	61700	460	66276	506	70415
323	50920	369	56703	415	61805	461	66370	507	70501
324	51055	370	56820	416	61909	462	66464	508	70586
325	51188	371	56937	417	62014	463	66558	509	70672
326	51322	372	57054	418	62118	464	66652	510	70757
327	51455	373	57171	419	62221	465	66745	511	70842
328	51587	374	57287	420	62325	466	66839	512	70927
329	51720	375	57403	421	62428	467	66932	513	71011
330	51851	376	57519	422	62531	468	67025	514	71096
331	51983	377	57634	423	62634	469	67117	515	71181
332	52114	378	57749	424	62737	470	67210	516	71265
333	52244	379	57864	425	62839	471	67302	517	71349
334	52375	380	57978	426	62941	472	67394	518	71433
335	52504	381	58093	427	63043	473	67486	519	71517
336	52634	382	58206	428	63144	474	67578	520	71600
337	52763	383	58320	429	63246	475	67669	521	71684
338	52892	384	58433	430	63347	476	67761	522	71767
339	53020	385	58546	431	63448	477	67852	523	71850
340	53148	386	58659	432	63548	478	67943	524	71933
341	53275	387	58771	433	63649	479	68034	525	72016
342	53403	388	58883	434	63749	480	68124	526	72099
343	53529	389	58995	435	63849	481	68215	527	72181
344	53656	390	59106	436	63949	482	68305	528	72263
345	53782	391	59218	437	64048	483	68395	529	72346
346	53908	392	59329	438	64147	484	68485	530	72428
347	54033	393	59439	439	64246	485	68574	531	72509
348	54158	394	59550	440	64345	486	68664	532	72591
349	54283	395	59660	441	64444	487	68753	533	72673
350	54407	396	59770	442	64542	488	68842	534	72754
351	54531	397	59879	443	64640	489	68931	535	72835
352	54654	398	59988	444	64738	490	69020	536	72916
353	54777	399	60097	445	64836	491	69108	537	72997
354	54900	400	60206	446	64933	492	69197	538	73078
355	55023	401	60314	447	65031	493	69285	539	73159
356	55145	402	60423	448	65128	494	69373	540	73239
357	55267	403	60530	449	65225	495	69461	541	73320
358	55388	404	60638	450	65321	496	69548	542	73400
359	55509	405	60746	451	65418	497	69636	543	73480
360	55630	406	60853	452	65514	498	69723	544	73560

No.	Log.	No.	Log.	No.	Log.	No.	Log.	No.	Log.
545	73640	591	77159	637	80414	683	83442	729	86273
546	73719	592	77232	638	80482	684	83506	730	86332
547	73799	593	77305	639	80550	685	83569	731	86392
548	73878	594	77379	640	80618	686	83632	732	86451
549	73957	595	77452	641	80686	687	83696	733	86510
550	74036	596	77525	642	80754	688	83759	734	86570
551	74115	597	77597	643	80821	689	83822	735	86629
552	74194	598	77670	644	80889	690	83885	736	86688
553	74273	599	77743	645	80956	691	83948	737	86747
554	74351	600	77815	646	81023	692	84011	738	86806
555	74429	601	77887	647	81090	693	84073	739	86864
556	74507	602	77960	648	81158	694	84136	740	86923
557	74586	603	78032	649	81224	695	84198	741	86982
558	74663	604	78104	650	81291	696	84261	742	87040
559	74741	605	78176	651	81358	697	84323	743	87099
560	74819	606	78247	652	81425	698	84385	744	87157
561	74896	607	78319	653	81491	699	84448	745	87216
562	74974	608	78390	654	81558	700	84510	746	87274
563	75051	609	78462	655	81624	701	84572	747	87332
564	75128	610	78533	656	81690	702	84634	748	87390
565	75205	611	78604	657	81757	703	84696	749	87448
566	75281	612	78675	658	81823	704	84757	750	87506
567	75358	613	78746	659	81889	705	84819	751	87564
568	75435	614	78817	660	81954	706	84880	752	87622
569	75511	615	78888	661	82020	707	84942	753	87680
570	75587	616	78958	662	82086	708	85003	754	87737
571	75664	617	79029	663	82151	709	85065	755	87795
572	75740	618	79099	664	82217	710	85126	756	87852
573	75815	619	79169	665	82282	711	85187	757	87910
574	75891	620	79239	666	82347	712	85248	758	87967
575	75967	621	79309	667	82413	713	85309	759	88024
576	76042	622	79379	668	82478	714	85370	760	88081
577	76118	623	79449	669	82543	715	85431	761	88138
578	76192	624	79518	670	82607	716	85491	762	88196
579	76268	625	79588	671	82672	717	85552	763	88252
580	76343	626	79657	672	82737	718	85612	764	88309
581	76418	627	79727	673	82802	719	85673	765	88366
582	76492	628	79796	674	82866	720	85733	766	88423
583	76567	629	79865	675	82930	721	85794	767	88480
584	76641	630	79934	676	82995	722	85854	768	88536
585	76716	631	80003	677	83059	723	85914	769	88593
586	76790	632	80072	678	83123	724	85974	770	88649
587	76864	633	80140	679	83189	725	86034	771	88705
588	76938	634	80209	680	83251	726	86094	772	88762
589	77012	635	80277	681	83315	727	86153	773	88818
590	77085	636	80346	682	83378	728	86213	774	88874

No.	Log.	No.	Log.	No.	Log.	No.	Log.	No.	Log.
775	88930	820	91381	865	93702	910	95904	955	98000
776	88986	821	91434	866	93752	911	95952	956	98046
777	88042	822	91487	867	93802	912	95999	957	98091
778	89098	823	91540	868	93852	913	96047	958	98137
779	89154	824	91593	869	93902	914	96095	959	98182
780	89209	825	91645	870	93952	915	96142	960	98227
781	89265	826	91698	871	94002	916	96190	961	98272
782	89321	827	91751	872	94052	917	96237	962	98318
783	89376	828	91803	873	94101	918	96284	963	98363
784	89432	829	91855	874	94151	919	96332	964	98408
785	89487	830	91908	875	94201	920	96379	965	98453
786	89542	831	91960	876	94250	921	96426	966	98498
787	89597	832	92012	877	94300	922	96473	967	98543
788	89653	833	92065	878	94349	923	96520	968	98588
789	89708	834	92117	879	94399	924	96567	969	98632
790	89763	835	92169	880	94448	925	96614	970	98677
791	89818	836	92221	881	94498	926	96661	971	98722
792	89873	837	92273	882	94547	927	96708	972	98767
793	89927	838	92324	883	94596	928	96754	973	98811
794	89982	839	92376	884	94645	929	96801	974	98856
795	90037	840	92428	885	94694	930	96848	975	98900
796	90091	841	92480	886	94743	931	96895	976	98945
797	90146	842	92531	887	94792	932	96941	977	98990
798	90200	843	92583	888	94841	933	96988	978	99034
799	90255	844	92634	889	94890	934	97035	979	99078
800	90309	845	92686	890	94949	935	97081	980	99123
801	90363	846	92737	891	94988	936	97128	981	99167
802	90417	847	92788	892	95036	937	97174	982	99211
803	90472	848	92840	893	95085	938	97220	983	99255
804	90526	849	92891	894	95134	939	97267	984	99300
805	90580	850	92942	895	95182	940	97313	985	99344
806	90634	851	92993	896	95231	941	97359	986	99388
807	90687	852	93044	897	95279	942	97405	987	99432
808	90741	853	93095	898	95328	943	97451	988	99476
809	90795	854	93146	899	95376	944	97497	989	99520
810	90849	855	93197	900	95424	945	97543	990	99564
811	90902	856	93247	901	95472	946	97589	991	99607
812	90956	857	93298	902	95521	947	97635	992	99651
813	91009	858	93349	903	95569	948	97681	993	99695
814	91062	859	93399	904	95617	949	97727	994	99739
815	91116	860	93450	905	95665	950	97772	995	99782
816	91169	861	93500	906	95713	951	97818	996	99826
817	91222	862	93551	907	95761	952	97864	997	99870
818	91275	863	93601	908	95809	953	97909	998	99913
819	91328	864	93651	909	95856	954	97955	999	99957

EXAMPLES. LXI. a.

Prove that the following statements are correct to four significant figures :

(1) $\sqrt[3]{451} = 7{\cdot}669.$

(2) $\sqrt[5]{802} = 3{\cdot}809.$

(3) $(273)^{\frac{4}{3}} \times (234)^{\frac{1}{4}} = 47{\cdot}32.$

(4) $(451)^{\frac{3}{5}} \times (231)^{\frac{4}{3}} = 55460.$

(5) $\left(\frac{192{\cdot}5}{84}\right)^3 = 12{\cdot}03.$

(6) $\frac{(34{\cdot}79)^{\frac{2}{3}}}{(41{\cdot}25)^{\frac{3}{2}}} = {\cdot}04023.$

(7) $\frac{(24{\cdot}76)^{\frac{2}{7}}}{({\cdot}0045)^{\frac{3}{2}}} = 8287.$

(8) $\frac{7{\cdot}89}{{\cdot}0345} \times (89130)^{\frac{1}{7}} = 1165.$

(9) $\frac{\frac{3}{2}\sqrt{(5{\cdot}2)}}{5\sqrt{(11{\cdot}31)}} \times (\tfrac{3}{7})^{-\frac{1}{2}} = {\cdot}3107.$

(10) $\sqrt[5]{\left\{\frac{2\sqrt{(34)}}{3\sqrt{(791)}}\right\}} = {\cdot}6731.$

(11) $\frac{\sqrt[4]{3}}{\sqrt[6]{3}} = 1{\cdot}096.$

(12) $\left(\frac{21^3 \times 45^5}{2^7 \times 3^9}\right)^{\frac{1}{2}} = 823{\cdot}6.$

Solve the equations correct to 4 figures :

(13) $10^x = 421$ $[x = 2{\cdot}624]$.

(14) $(\frac{21}{20})^x = 3$ $[x = 22{\cdot}51]$.

(15) $(\frac{203}{200})^{2x} = 2$ $[x = 23{\cdot}28]$.

(16) $(\frac{26}{25})^x = 3$ $[x = 28{\cdot}01]$.

(17) $\log 37^{x+8} = 3{\cdot}412$ $[x = -{\cdot}8243]$.

(18) $x = 10\sqrt[3]{(31{\cdot}2)}$ $[x = 31{\cdot}48]$.

Example i. *Find the* amount at Compound Interest *on* £1 *for* 8 years *at* 5 per cent.

To find the amount for 1 year we multiply by $\frac{105}{100}$, i.e. by $\frac{21}{20}$.

The amount for 2 years will be $£\frac{21}{20} \times \frac{21}{20}$ and the amount for 8 years $= (\frac{21}{20})^8$.

Let x be the required amount in pounds, then

$$x = (\tfrac{21}{20})^8,$$

$$\therefore \log x = 8(\log 21 - \log 20)$$

$$= 8(1{\cdot}32222 - 1{\cdot}30103) = 8({\cdot}02119)$$

$$= {\cdot}16852 = \log 1{\cdot}474\ldots$$

Hence, to find the amount at Compound Interest for 8 years at 5 per cent. we multiply the Principal expressed in pounds by $1{\cdot}474+\ldots$

Example ii. *In how many years will the Principal be doubled at* 5 per cent. *Compound Interest?*

Let x be the number of years, then

$(\frac{21}{20})^x$ is the amount at the end of x years,

hence $(\frac{21}{20})^x = 2$,

or $x(\log 21 - \log 20) = \log 2$, $\therefore\ x = \dfrac{\cdot 30103}{\cdot 02119} = \dfrac{30103}{2119} = 14\cdot 2$.

Example iii. *Find the present value of an annuity of* £100 a year *payable yearly for* 10 years, *the first payment to be at the end of a year, Compound Interest being reckoned at* 4 *per cent. per annum.*

The principal which will amount to £1 at the end of n years at 4 per cent. Comp. Int. is [See Lock's *Arithmetic*, p. 193, Ex. iv.]

$$£(\tfrac{100}{104})^n.$$

Hence the required present value is

$$£100\{r + r^2 + r^3 + \ldots + r^{10}\}, \text{ where } r = \tfrac{100}{104}$$

$$= £100\,\frac{1 - r^{11}}{1 - r} - £100 = £100\,\{1 - (\tfrac{100}{104})^{11}\} \times 26 - £100.$$

$$\log(\tfrac{100}{104})^{11} = 11(\log 100 - \log 104) = (2 - 2\cdot 01703) \times 11$$

$$= (\bar{1}\cdot 98297) \times 11 = \bar{1}\cdot 81267$$

$$= \log \cdot 6496\ldots$$

$\therefore$ the required present value $= £100 \times \cdot 3503\ldots \times 26 - £100$,

that is £810 about.

EXAMPLES. LXI. b.

(1) Find the Compound Interest on £100 for 10 years at 4 per cent. [*Result*, £48.]

(2) Find the Compound Interest on £1 for 8 years at 5 per cent. [*Result*, 9*s.* 6½*d.*]

(3) In how many years will a sum of money be doubled at 3 per cent. Compound Interest? [*Result*, 23·4.]

(4) In how many years will a sum of money be doubled at 4 per cent. Compound Interest? [*Result*, 17·7.]

(5) Find the present value of £100 to be paid 8 years hence reckoning Compound Interest at 4 per cent. [*Result*, £73·07.]

(6) If the number of births in a town are 25 per 1000 and the deaths 20 per 1000 annually, in how many years will the population be doubled? [*After* 140 *years.*]

(7) On the birth of an infant £1000 is invested at Compound Interest in the Funds (3 per cent. payable half-yearly); calculate what it will be worth when the child is 21 years old. [*Result*, £1869.]

(8) In what time will a sum of money treble itself at 3 per cent. Compound Interest payable half-yearly? [*Result*, 36·9 *years.*]

(9) A sum of 1 shilling lent on condition of 1 penny interest being paid monthly, accumulates at Compound Interest at the same rate for 12 years; what will be then the amount? [*Result, about* £5066.]

(10) A man puts by 2*d.* at the end of the second week of the year, 4*d.* at the end of the fourth week, 8*d.* at the end of the sixth week; what sum would be put by for the last fortnight in the year? [*About* 67,100,000 *pence.*]

(11) A train starting from rest has at the end of 1 second velocity ·001 ft. per sec. and at the end of each second its velocity is greater by one-third than at the end of the preceding second; find the velocity in miles per hour at the end of 25 seconds. [·679 *miles per hour.*]

(12) The volume of a sphere is $\frac{4}{3}\pi \times$(cube of the radius); find the diameter of the sphere which contains a cubic yard. [*Result*, 1·24 *yards.*]

(13) Find the present value at 4 per cent. per annum Compound Interest of a Fellowship of £250 a year for six years, payable half-yearly, the first payment being due in six months. [*Result*, £1322.]

CHAPTER XVI.

On the Relations between the Sides and Angles of a Triangle.

231. The **three sides** and the **three angles** of any **triangle,** are called its six **parts.**

By the letters A, B, C we shall indicate *geometrically*, the three angular points of the triangle ABC; *algebraically*, the three angles at those angular points respectively.



By the letters a, b, c we shall indicate the measures of the sides BC, CA, AB opposite the angles A, B, C respectively.

232. I. We know that, $A + B + C = 180^\circ$. [Euc. I. 32.]

233. Also if A be an angle of a triangle, then A may have any value between 0° and 180°. Hence,

(i) **sin A** must be positive (and less than 1),

(ii) **cos A** may be positive or negative (but must be numerically less than 1),

(iii) **tan A** may have any value whatever, positive or negative.

234. Also, if we are given the value of

(i) **sin A**, there are **two** angles, each less than 180°, which have the given positive value for their sine.

(ii) **cos A**, or (iii) **tan A**, then there is only **one** value of A, which value can be found from the Tables.

235. $\frac{A}{2}+\frac{B}{2}+\frac{C}{2}=90°$. Therefore $\frac{A}{2}$ is less than 90°, and its Trigonometrical Ratios are all positive. Also, $\frac{A}{2}$ is known, when the value of any one of its Ratios is given. Similar remarks of course apply to the angles B and C.

Example 1. *To prove* $\sin(A+B)=\sin C$.

$$A+B+C=180° \quad \therefore \quad A+B=180°-C,$$

and $\therefore$ $\sin(A+B)=\sin(180°-C)=\sin C$. [p. 104.]

Example 2. *To prove* $\sin\frac{A+B}{2}=\cos\frac{C}{2}$,

Now $\frac{A+B+C}{2}=90°$. $\therefore$ $\frac{A+B}{2}=90°-\frac{C}{2}$,

and $\therefore$ $\sin\frac{A+B}{2}=\sin\left(90°-\frac{C}{2}\right)=\cos\frac{C}{2}$. [Art. 118.]

Example 3. *To prove*

$$\sin A+\sin B+\sin C=4\cos\frac{A}{2}\cdot\cos\frac{B}{2}\cdot\cos\frac{C}{2}.$$

Now $\sin A+\sin B=2\sin\frac{A+B}{2}\cdot\cos\frac{A-B}{2}$. [Art. 157.]

$=2\cos\frac{C}{2}\cdot\cos\frac{A-B}{2}$. [Art. 118.]

and $\sin C=2\sin\frac{C}{2}\cdot\cos\frac{C}{2}$. [Art. 166.]

$=2\cos\frac{A+B}{2}\cdot\cos\frac{C}{2}$. [Art. 118.]

$$\therefore \sin A+\sin B+\sin C=2\cos\frac{C}{2}\cdot\cos\frac{A-B}{2}+2\cos\frac{C}{2}\cdot\cos\frac{A+B}{2},$$

$$=2\cos\frac{C}{2}\left(\cos\frac{A-B}{2}+\cos\frac{A+B}{2}\right).$$

$$=2\cos\frac{C}{2}\left(2\cos\frac{A}{2}\cdot\cos\frac{B}{2}\right). \quad \text{[Art. 157.]}$$

$$=4\cos\frac{A}{2}\cdot\cos\frac{B}{2}\cdot\cos\frac{C}{2}. \quad \text{Q.E.D.}$$

EXAMPLES. LXII.

Find A from each of the six following equations, A being an angle of a triangle.

(1) $\cos A=\frac{1}{2}$. (2) $\cos A=-\frac{1}{2}$. (3) $\sin A=\frac{1}{2}$.

(4) $\tan A=-1$. (5) $\sin A=\frac{1}{\sqrt{2}}$ (6) $\tan A=-\sqrt{3}$.

Prove the following statements, A, B, C being the angles of a triangle.

(7) $\sin(A+B+C)=0$. (8) $\cos(A+B+C)=-1$.

(9) $\sin\frac{A+B+C}{2}=1$. (10) $\cos\frac{A+B+C}{2}=0$.

(11) $\tan(A+B)=-\tan C$. (12) $\cot\frac{B+C}{2}=\tan\frac{A}{2}$.

(13) $\cos(A+B)=-\cos C$. (14) $\cos(A+B-C)=-\cos 2C$.

(15) $\tan A-\cot B=\cos C\cdot\sec A\cdot\operatorname{cosec} B$.

(16) $\frac{\sin A-\sin B}{\sin A+\sin B}=\tan\frac{C}{2}\cdot\tan\frac{A-B}{2}$.

(17) $\frac{\sin 3B-\sin 3C}{\cos 3C-\cos 3B}=\tan\frac{3A}{2}$.

(18) $\sin A+\sin B-\sin C=4\sin\frac{A}{2}\cdot\sin\frac{B}{2}\cdot\cos\frac{C}{2}$.

* (19) $\sin A - \sin B + \sin C = 4 \sin \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \sin \frac{C}{2}$.

(20) $\sin \frac{A}{2} \cdot \cos \frac{A}{2} + \sin \frac{B}{2} \cdot \cos \frac{B}{2} + \sin \frac{C}{2} \cdot \cos \frac{C}{2}$
$= 2 \cos \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2}$.

(21) $\cos A + \cos B + \cos C - 1 = 4 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}$.

(22) $\cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} - \cos^2 \frac{C}{2} = 2 \cos \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \sin \frac{C}{2}$.

(23) $\sin^2 \frac{A}{2} - \sin^2 \frac{B}{2} + \sin^2 \frac{C}{2} = 1 - 2 \cos \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \cos \frac{C}{2}$.

(24) $\sin \frac{B+C-A}{2} + \sin \frac{C+A-B}{2} + \sin \frac{A+B-C}{2} - 1$
$= 4 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}$.

(25) $\sin 2A + \sin 2B + \sin 2C = 4 \sin A \cdot \sin B \cdot \sin C$.

(26) $\sin A \cdot \cos A - \sin B \cdot \cos B + \sin C \cdot \cos C$
$= 2 \cos A \cdot \sin B \cdot \cos C$.

(27) $\sin (B+C-A) - \sin (C+A-B) + \sin (A+B-C)$
$= 4 \cos A \cdot \sin B \cdot \cos C$.

(28) $\cos 2A + \cos 2B + \cos 2C = -1 - 4 \cos A \cdot \cos B \cdot \cos C$.

(29) $\sin^2 A - \sin^2 B + \sin^2 C = 2 \sin A \cdot \cos B \cdot \sin C$.

(30) $\cos (B+C-A) + \cos (C+A-B) - \cos (A+B-C) + 1$
$= 4 \sin A \cdot \sin B \cdot \cos C$.

(31) $\sin \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2} + \sin \frac{B}{2} \cdot \cos \frac{C}{2} \cdot \cos \frac{A}{2}$
$+ \sin \frac{C}{2} \cdot \cos \frac{A}{2} \cdot \cos \frac{B}{2} = \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2} + 1$.

(32) $\tan A + \tan B + \tan C = \tan A \cdot \tan B \cdot \tan C$.

(33) $\tan \frac{B}{2} \cdot \tan \frac{C}{2} + \tan \frac{C}{2} \cdot \tan \frac{A}{2} + \tan \frac{A}{2} \cdot \tan \frac{B}{2} = 1$.

236. If ABC be a **right-angled** triangle having $C = 90°$, then $(A + B) = 90°$.



Hence, $\sin A = \sin (90° - B) = \cos B.$

Also, $$\sin A = \frac{a}{c} = \cos B,$$

and so on. (See page 52.)

EXAMPLES. LXIII.

In a right-angled triangle ABC, in which C is a right angle, prove the following statements.

(1) $\tan A = \cot B.$

(2) $\tan B = \cot A + \cos C.$

(3) $\sin 2A = \sin 2B.$

(4) $\cos 2A + \cos 2B = 0.$

(5) $\sin 2A = \frac{2ab}{c^2}.$

(6) $\operatorname{cosec} 2B = \frac{a}{2b} + \frac{b}{2a}.$

(7) $\cos 2A = \frac{b^2 - a^2}{c^2}.$

(8) $\cos 2B = \frac{\sin^2 A - \sin^2 B}{\sin^2 A + \sin^2 B}.$

(9) $\sin^2 \frac{B}{2} = \frac{c - a}{2c}.$

(10) $\cos^2 \frac{A}{2} = \frac{c + b}{2c}.$

(11) $\left(\cos \frac{A}{2} + \sin \frac{A}{2}\right)^2 = \frac{a + c}{c}.$

(12) $\frac{a - b}{a + b} = \tan \frac{A - B}{2}.$

(13) $\sin(A - B) + \cos 2A = 0.$

(14) $\sin(A - B) + \sin(2A + C) = 0.$

(15) $(\sin A - \sin B)^2 + (\cos A + \cos B)^2 = 2.$

(16) $\sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}} = \frac{2 \sin A}{\sqrt{\cos 2B}}.$

237. II. *To prove* $a = b\cos C + c\cos B$.

From A, any one of the angular points, draw AD perpendicular to BC, or to BC produced if necessary.

There will be three cases. Fig. i. when both B and C are acute angles; Fig. ii. when one of them (B) is obtuse; Fig. iii. when one of them (B) is a right angle. Then,



Fig. i. $\dfrac{CD}{CA} = \cos ACD$; or, $CD = b\cos C$,

and $\dfrac{DB}{AB} = \cos ABD$; or, $DB = c\cos B$,

$\therefore a = CD + DB = b\cos C + c\cos B$.

Fig. ii. $\dfrac{CD}{CA} = \cos ACD$; or, $CD = b\cos C$,

$\dfrac{BD}{AB} = \cos ABD$; or, $BD = c\cos(180^\circ - B)$,

$\therefore a = CD - BD = b\cos C - c\cos(180^\circ - B)$

$= b\cos C + c\cos B$.

Fig. iii. $a = CB = b\cos C$

$= b\cos C + c\cos B$. [For, $\cos B = \cos 90^\circ = 0$.]

Similarly it may be proved that,

$b = c\cos A + a\cos C$; $c = a\cos B + b\cos A$.

238. III. *To prove that in any triangle, the sides are proportional to the sines of the opposite angles;* or, *To prove that* $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$.

From A, any one of the angular points, draw AD perpendicular to BC, or to BC produced if necessary. Then,

Fig. i. $AD = b \sin C$; for, $\frac{AD}{AC} = \sin C$ [Def.]:

also $AD = c \sin B$; for, $\frac{AD}{AB} = \sin B$.

$\therefore b \sin C = c \sin B$;

or, $\frac{b}{\sin B} = \frac{c}{\sin C}$.

Fig. ii. $AD = b \sin C$,

and $AD = c \sin ABD = c \sin (180° - B)$

$\therefore AD = c \sin B$;

$\therefore b \sin C = c \sin B$;

or, $\frac{b}{\sin B} = \frac{c}{\sin C}$.

Fig. iii. $AB = AC \,.\, \sin C$; or, $c = b \sin C$;

$\therefore \frac{c}{\sin C} = \frac{b}{\sin B}$. [For $\sin B = \sin 90° = 1$.]

Similarly it may be proved that

$$\frac{a}{\sin A} = \frac{b}{\sin B};$$

$$\therefore \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$ Q.E.D.

239. IV. *To prove that* $a^2 = b^2 + c^2 - 2bc \cos A$.

Take one of the angles A. Then of the other two, one must be acute. Let B be an acute angle. From C draw CF perpendicular to BA, or to BA produced if necessary.

There will be three figures according as A is less, greater than, or equal to a right angle. Then,



Fig. i. $BC^2 = CA^2 + AB^2 - 2 \,.\, BA \,.\, FA$; [Euc. II. 13]

or, $a^2 = b^2 + c^2 - 2c \,.\, FA$

$= b^2 + c^2 - 2cb \cos A$. [For $FA = b \,.\, \cos A$.]

Fig. ii. $BC^2 = CA^2 + AB^2 + 2 \,.\, BA \,.\, AF$; [Euc. II. 12]

or, $a^2 = b^2 + c^2 + 2cb \cos FAC$

$= b^2 + c^2 - 2bc \cos A$. [For $FAC = 180^\circ - A$.]

Fig. iii. $BC^2 = CA^2 + AB^2$; [Euc. I. 47]

or, $a^2 = b^2 + c^2 - 2bc \cos A$. [For $\cos A = \cos 90^\circ = 0$.]

Similarly it may be proved that

$$b^2 = c^2 + a^2 - 2ca \cos B,$$

and that $$c^2 = a^2 + b^2 - 2ab \cos C.$$

240. V. Hence,

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}, \quad \cos B = \frac{c^2 + a^2 - b^2}{2ca}, \quad \cos C = \frac{a^2 + b^2 - c^2}{2ab}.$$

241. VI. *To prove that* $\sin^2\frac{A}{2} = \frac{(a+c-b)(a+b-c)}{4bc}$,

and that $\cos^2\frac{A}{2} = \frac{(a+b+c)(b+c-a)}{4bc}$.

Since $\cos A = 1 - 2\sin^2\frac{A}{2}$, [Art. 166]

and $\cos A = \frac{b^2+c^2-a^2}{2bc}$; [Art. 240]

$$\therefore\ 2\sin^2\frac{A}{2} = 1 - \cos A = 1 - \frac{b^2+c^2-a^2}{2bc}$$

$$= \frac{2bc-(b^2+c^2-a^2)}{2bc} = \frac{a^2-(b^2-2bc+c^2)}{2bc}$$

$$= \frac{a^2-(b-c)^2}{2bc} = \frac{\{a-(b-c)\}\{a+(b-c)\}}{2bc}$$

$$\therefore\ \sin^2\frac{A}{2} = \frac{(a+c-b)(a+b-c)}{4bc}.$$ Q.E.D.

Again, $\cos A = 2\cos^2\frac{A}{2} - 1$; [Art. 166]

$$\therefore\ 2\cos^2\frac{A}{2} = 1 + \cos A = 1 + \frac{b^2+c^2-a^2}{2bc}$$

$$\therefore\ \cos^2\frac{A}{2} = \frac{(b+c)^2-a^2}{4bc} = \frac{(b+c+a)(b+c-a)}{4bc}.$$ Q.E.D.

242. Similarly it may be proved that

$$\sin^2\frac{B}{2} = \frac{(b+c-a)(b+a-c)}{4ca},\quad \cos^2\frac{B}{2} = \frac{(a+b+c)(a+c-b)}{4ca},$$

and that

$$\sin^2\frac{C}{2} = \frac{(c+a-b)(c+b-a)}{4ab},\quad \cos^2\frac{C}{2} = \frac{(a+b+c)(a+b-c)}{4ab}.$$

243. VII. Now let s stand for $\frac{a+b+c}{2}$, so that $(a+b+c)=2s$.

Then, $(b+c-a)=(b+c+a-2a)=(2s-2a)=2(s-a)$,

and $(c+a-b)=(c+a+b-2b)=(2s-2b)=2(s-b)$,

and $(a+b-c)=(a+b+c-2c)=(2s-2c)=2(s-c)$.

Then the result of Art. 241 may be written

$$\sin^2\frac{A}{2}=\frac{2(s-b)\,2(s-c)}{4bc};\quad \text{or, } \sin\frac{A}{2}=\sqrt{\frac{(s-b)(s-c)}{bc}},$$

and $\cos^2\frac{A}{2}=\frac{2s\,2(s-a)}{4bc}$; or, $\cos\frac{A}{2}=\sqrt{\frac{s(s-a)}{bc}}$,

and so on.

Hence, $$\tan\frac{A}{2}=\frac{\sin\frac{A}{2}}{\cos\frac{A}{2}}=\frac{\sqrt{(s-b)(s-c)}}{\sqrt{s(s-a)}}.$$

244. VIII. Again

$$\sin A=2\sin\frac{A}{2}\cdot\cos\frac{A}{2};\qquad \text{[Art. 166]}$$

$$\therefore \sin A=2\sqrt{\frac{(s-b)(s-c)}{bc}}\cdot\sqrt{\frac{s(s-a)}{bc}}$$

$$=\frac{2}{bc}\sqrt{s(s-a)(s-b)(s-c)}.$$

The letter S usually stands for $\sqrt{s(s-a)(s-b)(s-c)}$, so that the above may be written $\frac{\sin A}{a}=\frac{2S}{abc}$.

Similarly, $$\frac{\sin B}{b}=\frac{2S}{abc}=\frac{\sin C}{c}.$$

245. IX. *To prove that* $\frac{b-c}{b+c} \cdot \cot\frac{A}{2} = \tan\frac{B-C}{2}$.

Since $\frac{b}{\sin B} = \frac{c}{\sin C}$, let each of these fractions $= d$.

Then $b = d \sin B$, and $c = d \sin C$.

$$\therefore \frac{b-c}{b+c} = \frac{d\sin B - d\sin C}{d\sin B + d\sin C} = \frac{\sin B - \sin C}{\sin B + \sin C}$$

$$= \frac{2\sin\frac{B-C}{2} \cdot \cos\frac{B+C}{2}}{2\sin\frac{B+C}{2} \cdot \cos\frac{B-C}{2}} = \frac{\tan\frac{B-C}{2}}{\tan\frac{B+C}{2}}$$

$$= \frac{\tan\frac{B-C}{2}}{\cot\frac{A}{2}}. \quad \left[\text{Since } \tan\frac{B+C}{2} = \tan\left(90^\circ - \frac{A}{2}\right).\right]$$

$$\therefore \frac{b-c}{b+c} \cdot \cot\frac{A}{2} = \frac{\tan\frac{B-C}{2}}{\cot\frac{A}{2}} \cdot \cot\frac{A}{2} = \tan\frac{B-C}{2}. \quad \text{Q.E.D.}$$

Similarly, $\frac{c-a}{c+a} \cdot \cot\frac{B}{2} = \tan\frac{C-A}{2}$,

and $\frac{a-b}{a+b} \cdot \cot\frac{C}{2} = \tan\frac{A-B}{2}$.

246. NOTE. In the next chapter we shall prove, that given three of the parts of a triangle (one of which must be a side), we can find the remaining three parts by means of the formulæ of this chapter. Hence, these formulæ cannot be equivalent to more than three *independent* equations. It is an instructive, but somewhat difficult problem to take three of these formulæ (*e.g.* $A+B+C=180^\circ$, $b = c\cos A + a\cos C$, $c = b\cos A + a\cos B$) and deduce all the others from them.

247. The following examples are important.

Example 1. Suppose we are given that

$$\left.\begin{aligned} a &= b\cos C + c\cos B, \\ b &= c\cos A + a\cos C, \\ c &= a\cos B + b\cos A. \end{aligned}\right\}$$

Then, taking a times the first $+b$ times the second $-c$ times the third, we get

$$a^2+b^2-c^2 = (ab\cos C + ac\cos B) + (bc\cos A + ba\cos C) - (ca\cos B + cb\cos A),$$
$$= 2ab\cos C.$$

Example 2. Suppose we are given that $A+B+C=180^0$,

and that $$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}\,[=d],$$

then $$\frac{b^2+c^2-a^2}{2bc} = \frac{d^2\sin^2 B + d^2\sin^2 C - d^2\sin^2 A}{2d^2\sin B\sin C},$$

$$= \frac{\sin^2 B + \sin^2 C - \sin^2 A}{2\sin B\sin C} = \frac{2\cos A\,.\,\sin B\,.\,\sin C}{2\sin B\,.\,\sin C}$$

[Ex. 29, p. 194, since $(A+B+C)=180^0$.]

$$= \cos A.$$

248. The formula

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = d$$

is very frequently of use in solving examples.

Example. Prove that

$$a\cos A + b\cos B + c\cos C = 2a\sin B\sin C.$$

Since $a = d\sin A$, $b = d\sin B$, $c = d\sin C$,

the above may be written

$$d\sin A\,.\,\cos A + d\sin B\,.\,\cos B + d\sin C\,.\,\cos C = 2d\sin A\,.\,\sin B\,.\,\sin C,$$

or $$\sin 2A + \sin 2B + \sin 2C = 4\sin A\,.\,\sin B\,.\,\sin C,$$

which is Example (25) on page 194.

249. The student is advised to make himself thoroughly familiar with the following **formulæ**:

$$a = b\cos C + c\cos B \quad \text{..... (ii),}$$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}\,[= d] = \frac{abc}{2S} \quad \text{..... (iii),}$$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc} \quad \text{..... (v).}$$

$$\left.\begin{aligned} \sin\frac{A}{2} &= \sqrt{\frac{(s-b)(s-c)}{bc}} \\ \cos\frac{A}{2} &= \sqrt{\frac{s(s-a)}{bc}} \end{aligned}\right\} \quad \text{..... (vii),}$$

$$\sin A = \frac{2}{bc}\sqrt{s(s-a)(s-b)(s-c)} = \frac{2S}{bc} \quad \text{..... (viii}$$

$$\tan\frac{B-C}{2} = \frac{b-c}{b+c}\cdot\cot\frac{A}{2} \quad \text{..... (ix).}$$

EXAMPLES. LXIV.

In any triangle ABC prove the following statements:

(1) $\dfrac{\sin A + 2\sin B}{a + 2b} = \dfrac{\sin C}{c}$.

(2) $\dfrac{\sin^2 A - m\cdot\sin^2 B}{a^2 - m\cdot b^2} = \dfrac{\sin^2 C}{c^2}$.

(3) $a\cos A + b\cos B - c\cos C = 2c\cos A\cdot\cos B$.

(4) $(a+b)\sin\dfrac{C}{2} = c\cos\dfrac{A-B}{2}$.

(5) $(b-c)\cos\dfrac{A}{2} = a\sin\dfrac{B-C}{2}$.

(6) $\dfrac{\cos A}{\sin B\cdot\sin C} + \dfrac{\cos B}{\sin C\cdot\sin A} + \dfrac{\cos C}{\sin A\cdot\sin B} = 2$.

(7) $a\sin(B-C)+b\sin(C-A)+c\sin(A-B)=0.$

(8) $\dfrac{a-b}{c}=\dfrac{\cos B-\cos A}{1+\cos C}.$

(9) $\dfrac{b+c}{a}=\dfrac{\cos B+\cos C}{1-\cos A}.$

(10) $\sqrt{bc\sin B.\sin C}=\dfrac{b^2\sin C+c^2\sin B}{b+c}.$

(11) $a+b+c=(b+c)\cos A+(c+a)\cos B+(a+b)\cos C.$

(12) $b+c-a=(b+c)\cos A-(c-a)\cos B+(a-b)\cos C.$

(13) $\tan A=\dfrac{a\sin C}{b-a\cos C}.$

(14) $a(b^2+c^2)\cos A+b(c^2+a^2)\cos B+c(a^2+b^2)\cos C=3abc.$

(15) $a\cos(A+B+C)-b\cos(B+A)-c\cos(A+C)=0.$

(16) $\dfrac{\cos A}{a}+\dfrac{\cos B}{b}+\dfrac{\cos C}{c}=\dfrac{a^2+b^2+c^2}{2abc}.$

(17) $\dfrac{\tan B}{\tan C}=\dfrac{a^2+b^2-c^2}{a^2-b^2+c^2}.$ (18) $b\cos^2\dfrac{C}{2}+c\cos^2\dfrac{B}{2}=s.$

(19) $\tan\dfrac{B}{2}.\tan\dfrac{C}{2}=\dfrac{b+c-a}{b+c+a}.$

(20) $\tan\dfrac{A}{2}(b+c-a)=\tan\dfrac{B}{2}(c+a-b).$

(21) $c^2=(a+b)^2\sin^2\dfrac{C}{2}+(a-b)^2\cos^2\dfrac{C}{2}.$

** MISCELLANEOUS EXAMPLES. LXV.

(1) If p is the length of the perpendicular from A on BC,

$$\sin A=\frac{ap}{bc}.$$

(2) If $2\cos B.\sin C=\sin A$, then $B=C$.

(3) If $A=3B$, then $\sin B=\frac{1}{2}\sqrt{\dfrac{3b-a}{b}}$.

(4) If $\sqrt{bc}\sin B \,.\, \sin C = \dfrac{b^2 \sin B + c^2 \sin C}{b+c}$, then $B=C$.

(5) $a\cos\dfrac{B}{2}\,.\,\cos\dfrac{C}{2}\,.\,\operatorname{cosec}\dfrac{A}{2} = b\cos\dfrac{C}{2}\,.\,\cos\dfrac{A}{2}\,.\,\operatorname{cosec}\dfrac{B}{2}$

$$= c\cos\frac{A}{2}\,.\,\cos\frac{B}{2}\,.\,\operatorname{cosec}\frac{C}{2} = s.$$

(6) Given $\cos A = \frac{3}{5}$ and $\cos B = \frac{12}{13}$, prove that $\cos C = -\frac{16}{65}$.

(7) If $\sin^2 B + \sin^2 C = \sin^2 A$, then $A = 90^0$.

(8) If $\sin 2B + \sin 2C = \sin 2A$, then either $B = 90^0$ or $C = 90^0$.

(9) If $A : B : C = 1 : 2 : 5$, then $1 + 4\cos A \,.\, \cos B \,.\, \cos C = 0$, and a^2, b^2, c^2 are in A.P.

(10) $a\sin\dfrac{A}{2}\,.\,\sin\dfrac{B-C}{2} + b\sin\dfrac{B}{2}\,.\,\sin\dfrac{C-A}{2}$

$$+ c\sin\frac{C}{2}\,.\,\sin\frac{A-B}{2} = 0.$$

(11) If D is the middle point of BC, prove that

$$4AD^2 = 2b^2 + 2c^2 - a^2.$$

(12) Given that $a = 2b$, and that $A = 3B$, prove that $C = 60^0$.

(13) $abc\,(a\cos A + b\cos B + c\cos C) = 8S^2$.

(14) If $b\cos^2\dfrac{C}{2} + c\cos^2\dfrac{B}{2} = \dfrac{3a}{2}$, then b, a, c are in A.P.

(15) If D, E, F are the middle points of the sides BC, CA, AB, then

$$4\,(AD^2 + BE^2 + CF^2) = 3\,(a^2 + b^2 + c^2).$$

(16) If D is the middle point of BC, $\cot ADB = \dfrac{b^2 - c^2}{4S}$.

(17) If d, e, f are the perpendiculars from A, B, C on the opposite sides of the triangle, then

$$a\sin A + b\sin B + c\sin C = 2\,(d\cos A + e\cos B + f\cos C).$$

[Some of the Examples in the Appendix might be worked by the student at this stage.]

CHAPTER XVII.

On the Solution of Triangles.

250. The problem known as the **Solution of Triangles** may be stated thus: *When a sufficient number of the parts of a triangle are given, to find the magnitude of each of the other parts.*

251. When **three** parts of a Triangle (one of which must be a side) are given, the other parts can in general be determined.

There are four cases.

I. Given **three sides.** [Compare Euc. I. 8.]

II. Given **one side** and **two angles.** [Euc. I. 26.]

III. Given **two sides** and the **angle between** them. [Euc. I. 4.]

IV. Given **two sides** and the **angle opposite** one of them. [Compare Euc. VI. 7.]

Case I.

252. Given **three sides,** a, b, c. [Euc. I. 8; VI. 5.]

We find two of the angles from the formulæ

$$\tan\frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$$

$$\tan\frac{B}{2} = \sqrt{\frac{(s-c)(s-a)}{s(s-b)}}.$$

The third angle $C = 180 - A - B$.

253. In practical work we proceed as follows:

$$\log \tan \frac{A}{2} = \log \sqrt{\frac{(s-b)(s-c)}{s(s-a)}};$$

or,

$$L \tan \frac{A}{2} - 10 = \tfrac{1}{2}\{\log (s-b) + \log (s-c) - \log s - \log (s-a)\}.$$

Similarly,

$$L \tan \frac{B}{2} - 10 = \tfrac{1}{2}\{\log (s-c) + \log (s-a) - \log s - \log (s-b)\}.$$

254. Either of the formulæ $\sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}}$, $\cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}$ may also be used as above.

The $\sin \frac{A}{2}$ and the $\cos \frac{A}{2}$ formulæ are either of them as convenient as the $\tan \frac{A}{2}$ formula, when *one* of the angles only is to be found. If *all* the angles are to be found the tangent formula is convenient, because we can find the L tangents of two half angles from the same *four* logs, viz. $\log s$, $\log (s-a)$, $\log (s-b)$, $\log (s-c)$. To find the L sines of *two* half angles we require the *six* logarithms, viz. $\log (s-a)$, $\log (s-b)$, $\log (s-c)$, $\log a$, $\log b$, $\log c$.

Example. Given $a = 275{\cdot}35$, $b = 189{\cdot}28$, $c = 301{\cdot}47$ chains, find A and B.

Here, $s = 383{\cdot}05$, $s-a = 107{\cdot}70$, $s-b = 193{\cdot}77$, $s-c = 81{\cdot}58$.

Then

$$\begin{aligned} L \tan \frac{A}{2} &= 10 + \tfrac{1}{2}\{\log 193{\cdot}77 + \log 81{\cdot}58 - \log 383{\cdot}05 - \log 107{\cdot}70\} \\ &= 10 + \tfrac{1}{2}\{2{\cdot}2872865 + 1{\cdot}9115837 - 2{\cdot}5832555 - 2{\cdot}0322157\} \\ &= 9{\cdot}7916995 \qquad \text{[from the tables]}, \end{aligned}$$

whence $\frac{A}{2} = 31^\circ\, 45'\, 28{\cdot}5''$; $\therefore A = 63^\circ\, 30'\, 57''$.

Also

$$L\tan\frac{B}{2}=10+\tfrac{1}{2}\{\log 81{\cdot}58+\log 107{\cdot}70-\log 383{\cdot}05-\log 193{\cdot}77\}$$

$$=9{\cdot}5366287=L\tan 18^0\,59'\,9{\cdot}8'';$$

$$\therefore\ B=37^0\,58'\,20'';\ C=180^0-A-B=78^0\,30'\,43''.$$

255. This Case may also be solved by the formula

$$\cos A=\frac{b^2+c^2-a^2}{2bc}.$$

But this formula is not adapted for logarithmic calculation, and therefore is seldom used in practice.

It may sometimes be used with advantage, when the given lengths of a, b, c each contain less than three digits.

Example. Find the greatest angle of the triangle whose sides are 13, 14, 15.

Let $a=15$, $b=14$, $c=13$. Then the greatest angle is A.

$$\text{Now, } \cos A=\frac{14^2+13^2-15^2}{2\times 14\times 13}=\frac{140}{2\times 14\times 13}=\tfrac{5}{13}=\cdot 384615.$$

$=\cos 67^0\,23'$, nearly.

[By the Table of natural cosines.]

$\therefore$ the greatest angle $=67^0\,23'$.

EXAMPLES. LXVI.

(1) If $a=352{\cdot}25$, $b=513{\cdot}27$, $c=482{\cdot}68$ yards, find the angle A, having given

$\log 674{\cdot}10=2{\cdot}8287243$, $\log 321{\cdot}85=2{\cdot}5076535$,

$\log 160{\cdot}83=2{\cdot}2063401$, $\log 191{\cdot}42=2{\cdot}2819873$,

$L\tan 20^0\,38'=9{\cdot}5758104$, $L\tan 20^0\,39'=9{\cdot}5761934$.

(2) Find the two largest angles of the triangle whose sides are 484, 376, 522 chains, having given that

$$\log 6{\cdot}91 = {\cdot}8394780,\quad \log 3{\cdot}15 = {\cdot}4983106,$$
$$\log 2{\cdot}07 = {\cdot}3159703,\quad \log 1{\cdot}69 = {\cdot}2278867,$$
$$L\tan 36^0\,46'\,6'' = 9{\cdot}8734581,\quad L\tan 31^0\,23'\,9'' = 9{\cdot}7853745.$$

(3) If $a=5238$, $b=5662$, $c=9384$ yards, find the angles A and B, having given

$$\log 1{\cdot}0142 = {\cdot}0061236,\quad \log 4{\cdot}904 = {\cdot}6905505,$$
$$\log\ 4{\cdot}48 = {\cdot}6512780,\quad \log\ 7{\cdot}58 = {\cdot}8796692,$$
$$L\tan 14^0\,38' = 9{\cdot}4168099,\quad L\tan 15^0\,57' = 9{\cdot}4560641,$$
$$L\tan 14^0\,39' = 9{\cdot}4173265,\quad L\tan 15^0\,58' = 9{\cdot}4565420.$$

(4) If $a=4090$, $b=3850$, $c=3811$ yards, find A, having given

$$\log 5{\cdot}8755 = {\cdot}7690448,\quad \log\ 3{\cdot}85 = {\cdot}5854607,$$
$$\log 1{\cdot}7855 = {\cdot}2517599,\quad \log 3{\cdot}811 = {\cdot}5810389,$$
$$L\cos 32^0\,15' = 9{\cdot}9272306,\quad L\cos 32^0\,16' = 9{\cdot}9271509.$$

(5) Find the greatest angle in a triangle whose sides are 7 feet, 8 feet, and 9 feet, having given

$$\log 3 = {\cdot}4771213,\quad L\cos 36^0\,42' = 9{\cdot}9040529,$$
$$\log 1{\cdot}4 = {\cdot}146128,\quad \text{diff. for } 60'' = {\cdot}0000942.$$

(6) Find the smallest angle of the triangle whose sides are 8 feet, 10 feet, and 12 feet, having given that

$$\log 2 = {\cdot}30103,\quad L\sin 20^0\,42' = 9{\cdot}5483585,\quad \text{diff. for } 60'' = {\cdot}0003342.$$

(7) If $a : b : c = 4 : 5 : 6$, find C, having given

$$\log 2 = {\cdot}3010300,\quad \log 3 = {\cdot}4771213,$$
$$L\cos 41^0\,25' = 9{\cdot}8750142,\quad \text{diff. for } 60'' = {\cdot}0001115.$$

(8) The sides of a triangle are 2, $\sqrt{6}$, and $1+\sqrt{3}$, find the angles.

(9) The sides of a triangle are 2, $\sqrt{2}$ and $\sqrt{3}-1$, find the angles.

Case II.

256. Given **one side** and **two angles**, as a, B, C.

[Euc. I. 26; VI. 4.]

First, $A = 180^\circ - B - C$; which determines A.

Next, $$\frac{b}{\sin B} = \frac{a}{\sin A}, \quad \text{or, } b = \frac{a \,.\, \sin B}{\sin A};$$

and, $$\frac{c}{\sin C} = \frac{a}{\sin A}, \quad \text{or, } c = \frac{a \,.\, \sin C}{\sin A}.$$

These determine b and c.

257. In practical work we proceed as follows:

$$\text{Since } b = \frac{a \,.\, \sin B}{\sin A},$$

$$\therefore \log b = \log \frac{a \,.\, \sin B}{\sin A}$$

$$\therefore \log b = \log a + \log (\sin B) + 10 - (10 + \log \sin A).$$

or, $$\log b = \log a + L \sin B - L \sin A.$$

Similarly, $\log c = \log a + L \sin C - L \sin A$.

Example. Given that $c = 1764{\cdot}3$ feet, $C = 18^\circ\,27'$, and $B = 66^\circ\,39'$, find b.

From the Tables we find $\log 1764{\cdot}3 = 3{\cdot}2465724$.

$L \sin 18^\circ\,27' = 9{\cdot}5003421$, $L \sin 66^\circ\,39' = 9{\cdot}9628904$;

$\therefore \log b = 3{\cdot}2465724 + 9{\cdot}9628904 - 9{\cdot}5003421$

$= 3{\cdot}7091207 = \log 5118{\cdot}2$;

$\therefore b = 5118{\cdot}2$ feet.

EXAMPLES. LXVII.

(1) If $A=53^0\,24'$, $B=66^0\,27'$, $c=338{\cdot}65$ yards, find C and a, having given that

$$L\sin 53^0\,24'=9{\cdot}9046168,\quad \log 3{\cdot}3865=\cdot5297511,$$
$$L\sin 60^0\;\;9'=9{\cdot}9381851,\quad \log 3{\cdot}1346=\cdot4961821,$$
$$\log 3{\cdot}1347=\cdot4961960.$$

(2) If $A=48^0$, $B=54^0$, and $c=38$ inches, find a and b, having given that

$$\log 38=1{\cdot}5797836,\quad \log 2{\cdot}88704=\;\cdot4604527,$$
$$\log 3{\cdot}14295=\;\cdot4973368,\quad L\sin 54^0=9{\cdot}9079576,$$
$$L\sin 78^0=9{\cdot}9904044,\quad L\sin 48^0=9{\cdot}8710735.$$

(3) Find c, having given that $a=1000$ yards, $A=50^0$, $C=66^0$, and that

$$L\sin 50^0=9{\cdot}8842540,\quad L\sin 66^0=9{\cdot}9607302,$$
$$\log 1{\cdot}19255=\cdot0764762.$$

(4) Find b, having given that $B=32^0\,15'$, $C=21^0\,47'\,20''$, $a=34$ feet.

$$\log 3{\cdot}4=\cdot531479,\quad L\sin 32^0\,15'=9{\cdot}727228,$$
$$\log 2{\cdot}241=\cdot350442,\quad L\sin 54^0\;\;2'=9{\cdot}908141,$$
$$\log 2{\cdot}242=\cdot350636,\quad L\sin 54^0\;\;3'=9\;908233.$$

(5) Find a, b, C, having given $A=72^0\,4'$, $B=41^0\,56'\,18''$, $c=24$ feet.

$$\log 2{\cdot}4\;\;=\cdot3802112,\quad L\sin 72^0\,4'\;\;=9{\cdot}9783702,$$
$$\log 1{\cdot}755\;=\cdot2442771,\quad L\sin 41^0\,56'\,10''=9{\cdot}8249725,$$
$$\log 1{\cdot}756\;=\cdot2445245,\quad L\sin 41^0\,56'\,20''=9{\cdot}8249959,$$
$$\log 2{\cdot}4995=\cdot3978531,\quad L\sin 65^0\,59'\;\;=9{\cdot}9606739,$$
$$\log 2{\cdot}4996=\cdot3978705,\quad L\sin 66^0\;\;\;\;=9{\cdot}9607302.$$

Case III.

258. Given **two sides** and the **included angle**, as b, c, A. [Euc. I. 4; VI. 6.]

First, $B + C = 180^\circ - A$. Thus $(B + C)$ is determined.

Next, $$\tan\frac{B-C}{2} = \frac{b-c}{b+c}\cot\frac{A}{2}.$$

Thus $(B - C)$ is determined.

And B and C can be found when the values of $(B + C)$ and $(B - C)$ are known.

Lastly, $$\frac{a}{\sin A} = \frac{b}{\sin B}, \text{ or } a = \frac{b \,.\, \sin A}{\sin B}.$$

Whence a is determined.

259. In practical work we proceed as follows;

Since $$\tan\frac{B-C}{2} = \frac{b-c}{b+c}\cot\frac{A}{2},$$

$$\therefore \log\left(\tan\frac{B-C}{2}\right) + 10$$

$$= \log(b-c) - \log(b+c) + \log\left(\cot\frac{A}{2}\right) + 10,$$

or, $$L\tan\frac{B-C}{2} = \log(b-c) - \log(b+c) + L\cot\frac{A}{2}.$$

Also, since $$a = \frac{b \,.\, \sin A}{\sin B},$$

$\therefore \log a = \log b + L\sin A - L\sin B$, as in Case II.

Example. Given $b=456{\cdot}12$ chains, $c=296{\cdot}86$ chains, and $A=74^0\,20'$, find the other angles.

Here, $b-c=159{\cdot}26$, $b+c=752{\cdot}98$.

From the Table we find,

$$\log 159{\cdot}26=2{\cdot}2021067, \text{ and } \log 752{\cdot}98=2{\cdot}8767834,$$

$$L\cot 37^0\,10'=10{\cdot}1202593\,;$$

$$\therefore L\tan\frac{B-C}{2}=2{\cdot}2021067-2{\cdot}8767834+10{\cdot}1202593$$

$$=9{\cdot}4455826=L\tan 15^0\,35'\,18''.$$

$$\therefore B-C=31^0\,10'\,36'', \text{ and } B+C=180^0-74^0\,20'.$$

Thus $B+C=105^0\,40'$;

$$\therefore 2B=136^0\,50'\,36''; \quad 2C=74^0\,29'\,24'',$$

or, $B=68^0\,25'\,18''$; or, $C=37^0\,14'\,42''$.

260. The formula $a^2=b^2+c^2-2bc\cos A$ may be used in simple cases; or it may be adapted to logarithmic calculation by the use of a subsidiary angle. [Cf. Example LXXVI. (20), p. 256.]

Example. If $b=35$ feet, $c=21$ feet, and $A=50^0$, find a, given that $\cos 50^0=\cdot643$.

Here $a^2=35^2+21^2-2\times35\times21\times\cos 50^0$;

$$\therefore \frac{a^2}{7^2}=5^2+3^2-2\times5\times3\times\cos 50^0,$$

$$=25+9-30\times\cdot643, \quad =14{\cdot}71.$$

$$\therefore \frac{a}{7}=3{\cdot}82 \text{ nearly}; \quad \text{or, } a=26{\cdot}74=\text{about } 26\tfrac{3}{4} \text{ feet.}$$

EXAMPLES. LXVIII.

(1) Find B and C, having given that $A=40^0$, $b=131$, $c=72$.

$\log 5{\cdot}9 = \cdot7708520$, $L\cot 20^0 = 10{\cdot}4389341$,

$\log 2{\cdot}03 = \cdot3074960$, $L\tan 38^0\,36' = 9{\cdot}9021604$,

$L\tan 38^0\,37' = 9{\cdot}9024195$.

(2) Find A and B, having given that $a=35$ feet, $b=21$ feet, $C=50^0$. $\log 2=\cdot301030$, $L\tan 28^0\,11'=9\cdot729020$,

$L\tan 65^0=10\cdot331327$, $L\tan 28^0\,12'=9\cdot729323$.

(3) If $b=19$ chains, $c=20$ chains, $A=60^0$, find B and C, having given that $\log 3\cdot9=\cdot591065$, $L\tan 2^0\,32'=8\cdot645853$,

$L\cot 30^0=10\cdot238561$, $L\tan 2^0\,33'=8\cdot648704$.

(4) Given that $a=376\cdot375$ chains, $b=251\cdot765$ chains, and $C=78^0\,26'$, find A and B. $L\cot 39^0\,13'=10\cdot0882755$,

$\log 1\cdot2461=\cdot0955529$, $L\tan 13^0\,39'=9\cdot3853370$,

$\log 6\cdot2814=\cdot7980565$, $L\tan 13^0\,40'=9\cdot3858876$.

(5) If $a=135$, $b=105$, $C=60^0$, find A, having given that

$\log 2=\cdot3010300$, $L\tan 12^0\,12'=9\cdot3348711$,

$\log 3=\cdot4771213$, $L\tan 12^0\,13'=9\cdot3354823$.

(6) If $a=21$ chains, $b=20$ chains, $C=60^0$, find c.

(7) Find c in the triangle of example (5).

(8) In a triangle the ratio of two sides is 5 : 3 and the included angle is $70^0\,30'$. Find the other angles.

$\log 2=\cdot3010300$, $L\cot 35^0\,15'=10\cdot1507464$,

$L\tan 19^0\,28'\,50''=9\cdot5486864$.

Case IV.

261. Given **two sides** and the **angle opposite** one of them, as b, c, B. [Omitted in Euc. I; Euc. VI. 7.]

First, since $\dfrac{c}{\sin C}=\dfrac{b}{\sin B}$; $\therefore\ \sin C=\dfrac{c\sin B}{b}$.

C must be found from this equation.

When C is known, $A = 180^\circ - B - C$,

and, $$a = \frac{b \sin A}{\sin B}.$$

Which solves the triangle.

262. We remark however that the angle C, found from the **trigonometrical** equation *sin* $C = a$ *given quantity*, where C is an angle of a triangle, has **two** values, one less than 90°, and one greater than 90°. [Art. 234.]

The question arises, Are both these values admissible?

This may be decided as follows:

If B is not less than 90°, C must be less than 90°; and the smaller value for C **only** is admissible.

If B is less than 90° we proceed thus.

1. If b is less than $c \sin B$, then $\sin C$, which $= \frac{c \sin B}{b}$, is greater than 1. This is impossible. Therefore if b is less than $c \sin B$, there is **no** solution whatever.

2. If b is equal to $c \sin B$, then $\sin C = 1$, and therefore $C = 90^\circ$; and there is only **one** value of C, viz. 90°.

3. If b is greater than $c \sin B$, and less than c, then B is less than C, and C may be obtuse or acute. In this case C may have either of the values found from the equation $\sin C = \frac{c \sin B}{b}$. Hence there are **two** solutions, and the triangle is said to be **ambiguous.**

4. If b is equal to or greater than c, then B is equal to or greater than C, so that C must be an acute angle; and the smaller value for C **only** is admissible.

263. The same results may be obtained **geometrically**.

Construction. Draw $AB = c$; make the angle ABD = the given angle B; with centre A and radius $= b$ describe a circle; draw AD perpendicular to BD.



Then $AD = c \sin B$.

1. If b is less than $c \sin B$, i.e. less than AD, the circle will not cut BD at all, and the construction **fails**. (Fig. i.)

2. If b is equal to AD, the circle will *touch* the line BD in the point D, and the required triangle is the **right-angled** triangle ABD. (Fig. ii.)

3. If b is greater than AD and less than AB, i.e. than c, the circle will cut the line BD in *two* points C_1, C_2 each on the same side of B. And we get **two** triangles ABC_1, ABC_2 each satisfying the given condition. (Fig. iii.)

4. If b is equal to c, the circle cuts BD in B and in one other point C; if b is greater than c the circle cuts BD in *two* points, but on *opposite* sides of B. In either case there is only **one** triangle satisfying the given condition. (Fig. iv.)

264. We may also obtain the same results **algebraically**, from the formula

$$b^2 = c^2 + a^2 - 2ca\cos B.$$

In this b, c, B are given, a is unknown. Write x for a and we get the **quadratic equation**

$$x^2 - 2c\cos B \,.\, x = b^2 - c^2.$$

Whence, $x^2 - 2c\cos B \,.\, x + c^2\cos^2 B = b^2 - c^2 + c^2\cos^2 B$

$$= b^2 - c^2\sin^2 B;$$

$$\therefore\ x = c\cos B \pm \sqrt{b^2 - c^2\sin^2 B}.$$

Let a_1, a_2 be the two values of x thus obtained, then

$$\left.\begin{aligned} a_1 &= c\cos B + \sqrt{b^2 - c^2\sin^2 B} \\ a_2 &= c\cos B - \sqrt{b^2 - c^2\sin^2 B} \end{aligned}\right\}.$$

Which of these two solutions is admissible, may be decided as follows;

1. If b is less than $c\sin B$, then $(b^2 - c^2\sin^2 B)$ is negative, so that a_1, a_2 are **impossible** quantities.

2. If b is equal to $c\sin B$, then $(b^2 - c^2\sin^2 B) = 0$, and $a_1 = a_2$; thus the *two* solutions become **one**.

3. If b is greater than $c\sin B$, then the **two** values a_1, a_2 are different and positive unless

$$\sqrt{b^2 - c^2\sin^2 B} \text{ is } > c\cos B,$$

i.e. unless $\quad b^2 - c^2\sin^2 B \quad > c^2\cos^2 B,$

i.e. unless $\quad b^2 \quad > c^2.$

4. If b is equal to c, then $a_2 = 0$; if b is greater than c, then a_2 is negative and is therefore inadmissible. In either of these cases a_1 is the **only** available solution.

265. We give two examples. In the first there are two solutions, in the second there is only one.

Example 1. Find A and C, having given that $b=379{\cdot}41$ chains, $c=483{\cdot}74$ chains, and $B=34^0\,11'$.

$$\begin{aligned} L\sin C &= \log c + L\sin B - \log b \\ &= 2{\cdot}6846120 + 9{\cdot}7496148 - 2{\cdot}5791088 \\ &= 9{\cdot}8551180 = L\sin 45^0\,45'; \end{aligned}$$

$$\therefore C = 45^0\,45', \text{ or, } 180^0 - 45^0\,45' = 134^0\,15'.$$

Since b is less than c, each of these values is admissible.

When $C = 45^0\,45'$, then $A = 100^0\,4'$.

When $C = 134^0\,15'$, then $A = 11^0\,34'$.

Example 2. Find A and C, when $b=483{\cdot}74$ chains, $c=379{\cdot}41$ chains, and $B=34^0\,11'$.

$$\begin{aligned} L\sin C &= \log c + L\sin B - \log b \\ &= 2{\cdot}5791088 + 9{\cdot}7496148 - 2{\cdot}6846120 \\ &= 9{\cdot}6441116 = L\sin 26^0\,9'; \end{aligned}$$

$$\therefore C = 26^0\,9', \text{ or, } 180^0 - 26^0\,9' = 153^0\,51'.$$

Since b is greater than c, C must be less than 90^0, and the larger value for C is inadmissible.

[It is also clear that $(153^0\,51' + 34^0\,11')$ is $> 180^0$].

$$\therefore C = 26^0\,9', \quad A = 119^0\,40'.$$

EXAMPLES. LXIX.

(1) If $B=40^0$, $b=140{\cdot}5$ feet, $a=170{\cdot}6$ feet, find A and C.

$\log 1{\cdot}405 = {\cdot}1476763$, $L\sin 40^0 = 9{\cdot}8080675$,

$\log 1{\cdot}706 = {\cdot}2319790$, $L\sin 51^0\,18' = 9{\cdot}8923342$,

$L\sin 51^0\,19' = 9{\cdot}8924354$.

(2) Find B and C, having given that $A=50^0$, $b=119$ chains, $a=97$ chains, and that

$$\log 1{\cdot}19=\cdot075547,\ L\sin 50^0\ =9{\cdot}884254,$$
$$\log 9{\cdot}7\ =\cdot986772,\ L\sin 70^0\ =9{\cdot}972986,$$
$$L\sin 70^0\,1'=9{\cdot}973032.$$

(3) Find B, C, and c, having given that $A=50^0$, $b=97$, $a=119$ (see example (2)).

$$\log 1{\cdot}553=\cdot191169,\ L\sin 38^0\,38'\,24''=9{\cdot}795479,$$
$$L\sin 88^0\,37'\,24''=9{\cdot}999876.$$

(4) Find A, having given that $a=24$, $c=25$, $C=65^0\,59'$, and that

$$\log 2{\cdot}5=\cdot3979400,\ L\sin 65^0\,59'=9{\cdot}9606739,$$
$$\log 2{\cdot}4=\cdot3802112,\ L\sin 61^0\,16'=9{\cdot}9429335,$$
$$L\sin 61^0\,17'=9{\cdot}9430028.$$

(5) If $a=25$, $c=24$, and $C=65^0\,59'$, find A, B and the greater value of b.

$$\log 1{\cdot}755=\cdot2442771,\ L\sin 72^0\,4'=9{\cdot}9783702,$$
$$\log 1{\cdot}756=\cdot2445245,\ L\sin 72^0\,5'=9{\cdot}9784111,$$
$$L\sin 41^0\,56'\,10''=9{\cdot}8249725,$$
$$L\sin 41^0\,56'\,26''=9{\cdot}8249959\ ;$$

for other logs see example (4).

(6) Supposing the data for the solution of a triangle to be as in the three following cases (α), (β), (γ), point out whether the solution will be ambiguous or not, and find the third side in the obtuse-angled triangle in the ambiguous case :

(α) $A=30^0$, $a=125$ feet, $c=250$ feet,

(β) $A=30^0$, $a=200$ feet, $c=250$ feet,

(γ) $A=30^0$, $a=200$ feet, $c=125$ feet.

$$\log 2=\cdot3010300,\quad L\sin 38^0\,41'=9{\cdot}7958800,$$
$$\log 6{\cdot}0389=\cdot7809578,\quad L\sin 8^0\,41'\ =9{\cdot}1789001,$$
$$\log 6{\cdot}0390=\cdot7809650.$$

266. It saves a little trouble in practice when using the formula $a = \frac{b \sin A}{\sin B}$, to write it thus ; $a = b \sin A \,.\, \text{cosec}\, B$.

For then, $\log a = \log b + L \sin A + L \text{cosec}\, B - 20$.

Thus the *subtraction* of a logarithm is avoided.

* 267. In the following Examples the student must find the necessary logarithms etc. from the Tables.

*MISCELLANEOUS EXAMPLES. LXX.

(1) Find A when $a = 374{\cdot}5$, $b = 576{\cdot}2$, $c = 759{\cdot}3$ feet.

(2) Find B when $a = 4001$, $b = 9760$, $c = 7942$ yards.

(3) Find C when $a = 8761{\cdot}2$, $b = 7643$, $c = 4693{\cdot}8$ chains.

(4) Find B when $A = 86^0\ 19'$, $b = 4930$, $c = 5471$ chains.

(5) Find C when $B = 32^0\ 58'$, $c = 1873{\cdot}5$, $a = 764{\cdot}2$ chains.

(6) Find c when $C = 108^0\ 27'$, $a = 36541$, $b = 89170$ feet.

(7) Find c when $B = 74^0\ 10'$, $C = 62^0\ 45'$, $b = 3720$ yards.

(8) Find b when $B = 100^0\ 19'$, $C = 44^0\ 59'$, $a = 1000$ chains.

(9) Find a when $B = 123^0\ 7'\ 20''$, $C = 15^0\ 9'$, $c = 9964$ yards.

Find the other two angles in the six following triangles.

(10) $C = 100^0\ 37'$, $b = 1450$, $c = 6374$ chains.

(11) $C = 52^0\ 10'$, $b = 643$, $c = 872$ chains.

(12) $A = 76^0\ 2'\ 30''$, $b = 1000$, $a = 2000$ chains.

(13) $C = 54^0\ 23'$, $b = 873{\cdot}4$, $c = 752{\cdot}8$ feet.

(14) $C = 18^0\ 21'$, $b = 674{\cdot}5$, $c = 269{\cdot}7$ chains.

(15) $A = 29^0\ 11'\ 43''$, $b = 7934$, $a = 4379$ feet.

(16) The difference between the angles at the base of a triangle is $17^0\ 48'$, and the sides subtending those angles are 105·25 feet and 76·75 feet ; find the third angle.

(17) If $b : c = 4 : 5$, $a = 1000$ yards and $A = 37^0\ 19'$, find b.

The student will find some Examples of Solution of Triangles without the aid of logarithms, in an Appendix.

*MISCELLANEOUS EXAMPLES. LXX b.

In the following Examples Logarithms are not required; they are not *practical* Examples, but some of them require for their solution considerable neatness and ingenuity; for which reason they are often set in examinations.

PART I.

(1) Simplify the formulæ

$$\cos A = \frac{b^2+c^2-a^2}{2bc}, \qquad \cos \tfrac{1}{2}A = \sqrt{\left\{\frac{s(s-a)}{bc}\right\}}$$

in the case of an equilateral triangle.

(2) The sides of a triangle are as $2 : \sqrt{6} : 1+\sqrt{3}$, find the angles.

(3) The sides of a triangle are as $4, 2\sqrt{2}, 2(\sqrt{3}-1)$, find the angles.

(4) Given $C=120^0$, $c=\sqrt{19}$, $a=2$, find b.

(5) Given $A=60^0$, $b=4\sqrt{7}$, $c=6\sqrt{7}$, find a.

(6) Given $A=45^0$, $B=60^0$ and $a=2$, find c.

(7) The sides of a triangle are as $7 : 8 : 13$, find the greatest angle.

(8) The sides of a triangle are $1, 2, \sqrt{7}$, find the greatest angle.

(9) The sides of a triangle are as $a : b : \sqrt{(a^2+ab+b^2)}$, find the greatest angle.

(10) When $a : b : c$ as $3 : 4 : 5$, find the greatest and least angles; given $\cos 36^0\,52' = \cdot 8$.

(11) If $a=5$ miles, $b=6$ miles, $c=10$ miles, find the greatest angle. [$\cos 49^0\,33' = \cdot 65$.]

(12) If $a=4$, $b=5$, $c=8$, find C; given that $\cos 54^0\,54' = \cdot 575$.

(13) The two sides a and b are $\sqrt{3}+1$ and 2 respectively; $C=30^0$; solve the triangle.

(14) Given $C=18^0$, $a=\sqrt{5}+1$, $c=\sqrt{5}-1$, find the other angles.

(15) If $b=3$, $C=120^0$, $c=\sqrt{13}$, find a and the sines of the other angles.

(16) Given $A=105^0$, $B=45^0$, $c=\sqrt{2}$, solve the triangle.

(17) Given $B=75^0$, $C=30^0$, $c=\sqrt{8}$, solve the triangle.

(18) Given $B=45^0$, $c=\sqrt{75}$, $b=\sqrt{50}$, solve the triangle.

(19) Given $B=30^0$, $c=150$, $b=50\sqrt{3}$, show that of the two triangles which satisfy the data one will be isosceles and the other right-angled. Find the third side in the greatest of these triangles.

(20) Is the solution ambiguous when $B=30^0$, $c=150$, $b=75$?

(21) If the angles adjacent to the base of a triangle are $22\frac{1}{2}^0$ and $112\frac{1}{2}^0$, show that the perpendicular altitude will be half the base.

(22) If $a=2$, $b=4-2\sqrt{3}$, $c=\sqrt{6}(\sqrt{3}-1)$, solve the triangle.

(23) If $A=9^0$, $B=45^0$, $b=\sqrt{6}$, find c.

(24) Given $B=15^0$, $b=\sqrt{3}-1$, $c=\sqrt{3}+1$, solve the triangle.

(25) Given $\sin B=\cdot 25$, $a=5$, $b=2\cdot 5$, find A. Draw a figure to explain the result.

(26) Given $C=15^0$, $c=4$, $a=4+\sqrt{48}$, solve the triangle.

(27) Two sides of a triangle are $3\sqrt{6}$ yards and $3(\sqrt{3}+1)$ yards, and the included angle 45^0, solve the triangle.

(28) If $C=30^0$, $b=100$, $c=45$, is the triangle ambiguous?

(29) Prove that if $A=45^0$ and $B=60^0$ then $2c=a(1+\sqrt{3})$.

(30) The cosines of two of the angles of a triangle are $\frac{1}{2}$ and $\frac{3}{5}$, find the ratio of the sides.

CHAPTER XVIII.

On the Measurement of Heights and Distances.

268. We have said (Art. 97) that the measurement, with scientific accuracy, of a line of any considerable length involves a long and difficult process.

On the other hand, sometimes it is required to find the *direction* of a line that it may point to an object which is not visible from the point from which the line is drawn. As for example when a tunnel has to be constructed.

By the aid of the Solution of Triangles

we can find the length of the distance between points which are inaccessible;

we can calculate the magnitude of angles which cannot be practically observed;

we can find the relative heights of distant and inaccessible points.

The method on which the Trigonometrical Survey of a country is conducted affords the following illustration.

269. *To find the distance between two distant objects.*



Two convenient positions A and B, on a level plain as far apart as possible, having been selected, the distance between A and B is measured with the greatest possible care. This line AB is called the **base** line. (In the survey of England, the base line is on Salisbury plain, and is about 36,578 feet long).

Next, the two distant objects, P and Q (church spires, for instance) visible from A and B, are chosen.

The angles PAB, PBA are observed. Then by Case II. Chapter XVII, the lengths of the lines PA, PB are calculated.

Again, the angles QAB, QBA are observed; and by Case II. the lengths of QA and QB are calculated.

Thus the lengths of PA and QA are found.

The angle PAQ is observed; and then by Case III. the length of PQ is calculated.

270. Thus the distance between two points P and Q has been found. The points P and Q are not necessarily accessible; the only condition being that P and Q must be *visible* from both A and B.

271. In practice, the points P and Q will generally be accessible, and then the line PQ, whose length has been calculated, may be used as a new base to find other distances.

272. *To find the height of a distant object above the point of observation.*



Let B be the point of observation; P the distant object. From B measure a base line BA of any convenient length, in any convenient direction; observe the angles PAB, PBA, and by Case II. calculate the length of BP. Next observe at B the 'angle of elevation' of P; that is, the angle which the line BP makes with the horizontal line BM, M being the point in which the vertical line through P cuts the horizontal plane through B.

Then PM, which is the vertical height of P above B can be calculated, for $PM = BP \cdot \sin MBP$.

Example 1. *The distance between a church spire* A *and a milestone* B *is known to be* 1764·3 *feet; C is a distant spire. The angle* CAB *is* 94° 54′. *and the angle* CBA *is* 66° 39′. *Find the distance of* C *from* A.

ABC is a triangle and we know one side c and two angles (A and B), and therefore it can be solved by Case II.

The angle $ACB = 180^0 - 94^0\ 54' - 66^0\ 39'$

$= 18^0\ 27'$.

Therefore the triangle is the same as that solved on page 210. Therefore $AC = 5118{\cdot}2$ feet.

Example 2. *If the spire* C *in the last Example stands on a hill, and the angle of elevation of its highest point is observed at* A *to be* $4^0\ 19'$; *find how much higher* C *is than* A.

The required height $x = AC \cdot \sin 4^0\ 19'$ [Art. 268], and AC is 5118·2 feet,

$$\therefore \log x = \log (AC \cdot \sin 4^0\ 19')$$
$$= \log 5118{\cdot}2 + L\sin 4^0\ 19' - 10$$
$$= 3{\cdot}7091173 + 8{\cdot}8766150 - 10$$
$$= 2{\cdot}5857323 = \log 385{\cdot}24.$$

Therefore $x = 385$ ft. 3 in. nearly.

EXAMPLES. LXXI.

(Exercises XIV. and LXI. consist of easy examples on this subject).

(1) Two straight roads inclined to one another at an angle of 60^0, lead from a town A to two villages B and C; B on one road distant 30 miles from A, and C on the other road distant 15 miles from A. Find the distance from B to C. Ans. 25·98 m.

(2) Two ships leave harbour together, one sailing N.E. at the rate of $7\frac{1}{2}$ miles an hour and the other sailing North at the rate of 10 miles an hour. Prove that the distance between the ships after an hour and a half is 10·6 miles.

(3) A and B are two consecutive milestones on a straight road and C is a distant spire. The angles ABC and BAC are observed to be 120^0 and 45^0 respectively. Show that the distance of the spire from A is 3·346 miles.

(4) If the spire C in the last question stands on a hill, and its angle of elevation at A is 15^0, show that it is ·866 of a mile higher than A.

(5) If in Question (3) there is another spire D such that the angles DBA and DAB are 45^0 and 90^0 respectively and the angle DAC is 45^0; prove that the distance from C to D is $2\frac{3}{4}$ miles very nearly.

(6) A and B are two consecutive milestones on a straight road, and C is the chimney of a house visible from both A and B. The angles CAB and CBA are observed to be $36^0\ 18'$ and $120^0\ 27'$ respectively. Show that C is 2639·5 yards from B,

log 1760 = 3·2455127	$L \sin 36^0\ 18' = 9{\cdot}7723314$
log 2639·5 = 3·42152	$L \operatorname{cosec} 23^0\ 15' = 10{\cdot}4036846.$

(7) A and B are two points on opposite sides of a mountain, and C is a place visible from both A and B. It is ascertained that C is distant 1794 feet and 3140 feet from A and B respectively and the angle ACB is $58^0\ 17'$. Show that the angle which the line pointing from A to B makes with AC is $86^0\ 55'\ 49''$,

log 1346 = 3·1290451	$L \cot 29^0\ 8'\ 30'' = 10{\cdot}2537194$
log 4934 = 3·6931991	$L \tan 26^0\ 4'\ 19'' = 9{\cdot}6895654$

(8) A and B are two hill-tops 34920 feet apart, and C is the top of a distant hill. The angles CAB and CBA are observed to be $61^0\ 53'$ and $76^0\ 49'$ respectively. Prove that the distance from A to C is 51515 feet,

log 34920 = 4·5430742	$L \sin 76^0\ 49' = 9{\cdot}9884008$
log 51515 = 4·71193	$L \operatorname{cosec} 41^0\ 18' = 10{\cdot}1804552.$

(9) From two stations A and B on shore, 3742 yards apart, a ship C is observed at sea. The angles BAC, ABC are simultaneously observed to be $72^0\ 34'$ and $81^0\ 41'$ respectively. Prove that the distance from A to the ship is 8522·7 yards,

log 3742 = 3·5731038	$L \sin 81^0\ 41' = 9{\cdot}9954087$
log 8522·7 = 3·9305774	$L \operatorname{cosec} 25^0\ 45' = 10{\cdot}3620649.$

(10) The distance between two mountain peaks is known to be 4970 yards, and the angle of elevation of one of them when seen from the other is $9^0\ 14'$. How much higher is the first than the second? $\sin 9^0\ 14' = {\cdot}1604555$. *Ans.* 797·5 yards.

(11) Two straight railways intersect at an angle of 60^0. From their point of intersection two trains start, one on each line, one at the rate of 40 miles an hour. Find the rate of the second train that at the end of an hour they may be 35 miles apart. *Ans.* Either 25 or 15 miles an hour. (Art. 264.)

(12) A and B are two positions on opposite sides of a mountain; C is a point visible from A and B; AC and BC are 10 miles and 8 miles respectively, and the angle BCA is 60^0. Prove that the distance between A and B is 9·165 miles.

(13) In the last question, if the angle of elevation of C at A is 8^0, and at B is $2^0\ 48'\ 24''$: show that the height of B above A is one mile very nearly.

$$\sin 8^0 = \cdot1391731 \quad \sin 2^0\ 48'\ 24'' = \cdot0489664.$$

(14) Show that the angles which a tunnel going through the mountain from A to B, in Questions (12) and (13), would make (i) with the horizon, (ii) with the line joining A and C, are respectively $6^0\ 16'$ and $49^0\ 6'\ 24''$.

$$\sin 6^0\ 16' = \cdot1091\ ; \quad \tan 10^0\ 53'\ 36'' = \cdot192450.$$

(15) A and B are consecutive milestones on a straight road; C is the top of a distant mountain. At A the angle CAB is observed to be $38^0\ 19'$; at B the angle CBA is observed to be $132^0\ 42'$, and the angle of elevation of C at B is $10^0\ 15'$. Show that the top of the mountain is 1243·5 yards higher than B.

$L \sin 38^0\ 19' = 9{\cdot}7923968$ $\qquad$ $\log 1760 = 3{\cdot}2455127$

$L \operatorname{cosec} 8^0\ 59' = 10{\cdot}8064659$ $\qquad$ $\log 1243{\cdot}5 = 3{\cdot}09465$

$L \sin 10^0\ 15' = 9{\cdot}2502822.$

(16) A base line AB, 1000 feet long is measured along the straight bank of a river; C is an object on the opposite bank; the angles BAC, and CBA are observed to be $65^0\ 37'$ and $53^0\ 4'$ respectively. Prove that the perpendicular breadth of the river at C is 829·87 feet; having given

$L \sin 65^0\ 37' = 9{\cdot}9594248$, $\qquad$ $L \sin 53^0\ 4' = 9{\cdot}9027289$

$L \operatorname{cosec} 61^0\ 19' = 10{\cdot}0568589$, $\qquad$ $\log 8{\cdot}2987 = \cdot91901.$

* MISCELLANEOUS EXAMPLES. LXXII.

(1) A man walking along a straight road at the rate of three miles an hour sees, in front of him at an elevation of 60^0, a balloon which is travelling horizontally in the same direction at the rate of six miles an hour; ten minutes after he observes that the elevation is 30^0. Prove that the height of the balloon above the road is $440\sqrt{3}$ yards.

(2) A person standing at a point A, due south of a tower built on a horizontal plain, observes the altitude of the tower to be 60^0. He then walks to a point B due west from A and observes the altitude to be 45^0, and then at the point C in AB produced he observes the altitude to be 30^0. Prove that $AB = BC$.

(3) The angle of elevation of a balloon, which is ascending uniformly and vertically, when it is one mile high is observed to be $35^0\ 20'$; 20 minutes later the elevation is observed to be $55^0\ 40'$. How fast is the balloon moving?

Ans. 3 (sin $20^0\ 20'$) (sec $55^0\ 40'$) (cosec $35^0\ 20'$) miles per hour.

(4) The angular elevation of a tower at a place A due south of it is 30^0; and at a place B due west of A, and at a distance a from it, the elevation is 18^0; show that the height of the tower is

$$\frac{a}{\sqrt{2\,(1+\sqrt{5})}}.$$

(5) The angular elevation of the top of a steeple at a place due south of it is 45^0, and at another place due west of the former station and distant a feet from it the elevation is 15^0; show that the height of the steeple is $\frac{a}{2}(3^{\frac{1}{2}} - 3^{-\frac{1}{2}})$ feet.

(6) A tower stands at the foot of an inclined plane whose inclination to the horizon is 9^0; a line is measured up the incline from the foot of the tower of 100 feet in length. At the upper extremity of this line the tower subtends an angle of 54^0. Find the height of the tower. *Ans.* 114·4 ft.

(7) The altitude of a certain rock is observed to be 47^0, and after walking 1000 feet towards the rock, up a slope inclined at an angle of 32^0 to the horizon, the observer finds that the altitude is 77^0. Prove that the vertical height of the rock above the first point of observation is 1034 ft. Sin $47^0 = \cdot 73135$.

(8) At the top of a chimney 150 feet high standing at one corner of a triangular yard, the angle subtended by the adjacent sides of the yard are 30^0 and 45^0 respectively; while that subtended by the opposite side is 30^0. Show that the lengths of the sides are 150 ft. 86·6 ft. and 106 ft. respectively.

(9) A flag staff h feet high stands on the top of a tower. From a point in the plain on which the tower stands the angles of elevation of the top and bottom of the flagstaff are observed to be α and β respectively. Prove that the height of the tower is $\dfrac{h \tan \beta}{\tan \alpha - \tan \beta}$ feet, i.e. $\dfrac{h \sin \beta . \cos \alpha}{\sin (\alpha - \beta)}$ feet.

(10) From the top of a cliff h feet high the angles of depression of two ships at sea in a line with the foot of the cliff are α and β respectively. Show that the distance between the ships is $h (\cot \beta - \cot \alpha)$ feet.

(11) The angular elevation of a tower at a place due south of it is α, and at another place due west of the first and distant d from it, the elevation is β. Prove that the height of the tower is

$$\frac{d}{\sqrt{\cot^2 \beta - \cot^2 \alpha}}, \text{ i.e. } \frac{d \sin \alpha . \sin \beta}{\sqrt{\sin (\alpha - \beta) . \sin (\alpha + \beta)}}.$$

(12) A man stands on the top of the wall of height h, and observes the angular elevation (α) of the top of a telegraph post; he then descends from the wall, and finds that the angular elevation is now β; prove that the height of the post exceeds the height of the man by $h \dfrac{\sin \beta . \cos \alpha}{\sin (\beta - \alpha)}$.

(13) Find the height of a cloud by observing (i) its elevation (α) and (ii) the depression (β) of its reflexion in a lake h feet below the point of observation. *Ans.* Height above the lake $h \sin(\alpha+\beta) \operatorname{cosec}(\beta-\alpha)$ feet.

(14) If the angular elevation of the summits of two spires (which appear in a straight line) is α, and the angular depressions of their reflexions in a lake, h feet below the point of observation, are β and γ, then the horizontal distance between the spires is $2h \cos^2 \alpha \sin(\beta-\gamma) . \operatorname{cosec}(\beta-\alpha) . \operatorname{cosec}(\gamma-\alpha)$ feet.

(15) A statue AB on the top of a pillar BC situated on level ground is found to subtend the greatest angle (α) at the eye of an observer E when he is distant c feet from the pillar. Prove that the height of the statue is $(2c \tan \alpha)$ feet; and, if b be the height of the observer's eye from the ground find the height of the pillar. (See Note in the Answers.)

(16) A man walking along a straight road observes that the greatest angle which two objects subtend at his eye is α. From the place where this happens he walks a yards; the objects then appear in a straight line making a right angle with the road. Prove that the distance between the objects is $(2a \tan \alpha)$ yards.

(17) A man standing on a horizontal plain at a distance h feet from a tower, observes that a flagstaff on the tower subtends an angle α at his eye, and that on walking $2k$ feet towards the tower the flagstaff subtends again the same angle. Prove that the height of the flagstaff is $2(h-k) \tan \alpha$ feet.

(18) A man walking along a straight road observes that the greatest angle which two objects subtend at his eye is α; from the place where this happens he walks a yards, and the objects then appear in a straight line making an angle β with the road. Prove that the distance between the objects is $\dfrac{2a \sin \alpha . \sin \beta}{\cos \alpha + \cos \beta}$ yds.

CHAPTER XIX.

On Triangles and Circles.

273. *To find the* **Area** *of a Triangle.*

The area of the triangle ABC is denoted by Δ.



Through A draw HK parallel to BC, and through ABC draw lines AD, BK, CH perpendicular to BC.

The area of the triangle ABC is half that of the rectangular parallelogram $BCHK$ [Euc. I. 41].

Therefore $$\Delta = \frac{BC \,.\, CH}{2} = \frac{BC \,.\, DA}{2}$$

$$= \frac{a \,.\, b \sin C}{2} \quad \text{.....(i).}$$

But $$\sin C = \frac{2}{ab} \,.\, \sqrt{s\,(s-a)\,(s-b)\,(s-c)}\,;$$

$$\therefore\ \Delta = \sqrt{s\,(s-a)\,(s-b)\,(s-c)} = S \text{.....(ii).}$$

274. *To find the Radius of the* **Circumscribing Circle.**

Fig. I.



Fig. II.



Let a circle $AA'CB$ be described about the triangle ABC. Let R stand for its radius. Let O be its centre. Join BO, and produce it to cut the circumference in A'. Join $A'C$.

Then, Fig. I. the angles BAC, $BA'C$ in the same segment are equal; Fig. II. the angles BAC, $BA'C$ are supplementary; also the angle BCA' in a semicircle is a right angle.

Therefore $$\frac{CB}{A'B} = \sin CA'B = \sin CAB = \sin A,$$

or, $$\frac{a}{2R} = \sin A;$$

$$\therefore\ 2R = \frac{a}{\sin A}.$$

275. Similarly, it may be proved that

$$2R = \frac{b}{\sin B};\ \text{and that}\ 2R = \frac{c}{\sin C}.$$

Hence, $$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R.$$

Thus d, the value of each of these fractions, is the diameter of the circumscribing circle.

276. *To find the radius of the* **Inscribed Circle.**



Let D, E, F be the points in which the circle inscribed in the triangle ABC touches the sides. Let I be the centre of the circle; let r be its radius. Then $ID = IE = IF = r$.

The area of the triangle ABC

$$= \text{area of } IBC + \text{area of } ICA + \text{area of } IAB.$$

And the area of the triangle $IBC = \frac{1}{2}ID \,.\, BC = \frac{1}{2}r \,.\, a$,

$$\therefore \text{area of } ABC = \tfrac{1}{2}ID \,.\, BC + \tfrac{1}{2}IE \,.\, CA + \tfrac{1}{2}IF \,.\, AB$$

$$= \tfrac{1}{2}ra + \tfrac{1}{2}rb + \tfrac{1}{2}rc;$$

or,
$$\Delta = \tfrac{1}{2}r(a + b + c) = \tfrac{1}{2}r \,.\, 2s = rs.$$

$$\therefore r = \frac{\Delta}{s} = \frac{S}{s}.$$

277. A circle which touches one of the sides of a triangle and the other two sides produced is called an **Escribed Circle** of the triangle.

278. *To find the radius of an* **Escribed Circle.**

Let an escribed circle touch the side BC and the sides AC, AB produced in the points D_1, E_1, F_1 respectively. Let I_1 be its centre, r_1 its radius. Then

$$I_1D_1 = I_1E_1 = I_1F_1 = r_1.$$

The area of the triangle ABC

$$= \text{area of } ABI_1C - \text{area of } I_1BC,$$

$$= \text{area of } I_1CA + \text{area of } I_1AB - \text{area of } I_1BC,$$

or
$$\Delta = \tfrac{1}{2}I_1E_1 \,.\, CA + \tfrac{1}{2}I_1F_1 \,.\, AB - \tfrac{1}{2}I_1D_1 \,.\, BC$$

$$= \tfrac{1}{2}r_1b + \tfrac{1}{2}r_1c - \tfrac{1}{2}r_1a$$

$$= \tfrac{1}{2}r_1(b + c - a) = \tfrac{1}{2}r_1(2s - 2a)$$

$$= r_1(s - a).$$

$$\therefore r_1 = \frac{\Delta}{s-a} = \frac{S}{s-a}.$$

279. Similarly if r_2 and r_3 be the radii of the other two escribed circles of the triangle ABC, then

$$r_2 = \frac{S}{s-b}; \quad r_3 = \frac{S}{s-c}.$$

* 280. The following results are often useful.

With the construction of the last two articles; the lines IA, IB, IC bisect the angles A, B, C respectively; the lines I_1A, I_1B, I_1C bisect the angles BAC, CBF_1, BCE_1 respectively; so that AII_1 is a straight line.



(i) Two tangents drawn from an external point to touch a circle are equal; therefore

$$AE = AF; \quad BD = BF; \quad CE = CD;$$
$$AE_1 = AF_1; \; BD_1 = BF_1; \; CE_1 = CD_1.$$

(ii) $$2AE_1 = AE_1 + AF_1 = (AC + CD_1) + (AB + BD_1)$$
$$= AC + CB + BA = a + b + c = 2s.$$

(iii) $$2AE = AE + AF = (AC - CD) + (AB - BD)$$
$$= AC + AB - BC = b + c - a = 2(s - a).$$

Similarly, $BF = BD = (s - b)$; $CD = CE = (s - c)$.

(iv) $$BD_1 = BF_1 = AF_1 - AB = (s - c).$$

Similarly, $$CE_1 = CD_1 = (s - b).$$

(v) Hence, $BD_1 = CD$, and $CD_1 = BD$.

(vi) $$FF_1 = AF_1 - AF = s - (s - a) = a.$$

Similarly, $$EE_1 = a.$$

(vii) $II_1 = AI_1 - AI = AF_1 \sec\frac{A}{2} - AF\sec\frac{A}{2} = FF_1 \sec\frac{A}{2}$

$$\therefore\ II_1 = \frac{a}{\cos\frac{A}{2}}.$$

(viii) $IF = AF\tan\frac{A}{2}$; or, $r = (s-a)\tan\frac{A}{2}$,

$I_1F = AF_1\tan\frac{A}{2}$; or, $r_1 = s\tan\frac{A}{2}$.

(ix) $AF = IF\cot\frac{A}{2}$, and $BF = IF\cot\frac{B}{2}$;

$$\therefore\ c = AF + BF = r\left(\cot\frac{A}{2} + \cot\frac{B}{2}\right);$$

$$\therefore\ r = \frac{c}{\cot\frac{A}{2} + \cot\frac{B}{2}} = \frac{c\sin\frac{A}{2}.\sin\frac{B}{2}}{\sin\frac{A+B}{2}} = \frac{c\sin\frac{A}{2}.\sin\frac{B}{2}}{\cos\frac{C}{2}}.$$

(x) $F_1BC = 180^\circ - B$, $\therefore\ I_1BC = 90^\circ - \frac{B}{2}$. Hence,

$$r_1\left\{\cot\left(90^\circ - \frac{B}{2}\right) + \cot\left(90^\circ - \frac{C}{2}\right)\right\} = a;$$

or,
$$r_1 = \frac{a\cos\frac{B}{2}.\cos\frac{C}{2}}{\cos\frac{A}{2}}.$$

(xi) $r_1 = AF_1\tan\frac{A}{2}$, and $r = AF\tan\frac{A}{2}$; therefore

$$r_1 - r = FF_1\tan\frac{A}{2} = a\tan\frac{A}{2}.$$

* 281. *To find an expression for the* **area** *of a quadrilateral inscribed in a circle, in terms of the sides.*



Let $DEFG$ be the quadrilateral; and let d, e, f, g represent the sides.

The angle $F = 180° - D$. [Euc. III. 22.]

The area of $GDE = \frac{1}{2}dg \sin D$. [Art. 273.]

The area of $EFG = \frac{1}{2}ef \sin F = \frac{1}{2}ef \sin (180° - D)$
$= \frac{1}{2}ef \sin D$.

Thus the area of the quadrilateral is

$$\tfrac{1}{2}(dg + ef) \sin D.$$

We must find an expression for $\sin D$ in terms of the sides.

From the triangle EGD we have

$$EG^2 = d^2 + g^2 - 2dg \cos D.$$ [Art. 239.]

From the triangle EGF we have

$$EG^2 = e^2 + f^2 - 2ef \cos F$$
$$= e^2 + f^2 + 2ef \cos D; \qquad [\text{For } F = 180° - D]$$

$$\therefore\ d^2 + g^2 - 2dg \cos D = e^2 + f^2 + 2ef \cos D;$$

$$\therefore\ 2(dg+ef)\cos D = d^2+g^2-e^2-f^2;$$

$$\therefore\ 1-\cos^2 D = 1-\frac{(d^2+g^2-e^2-f^2)^2}{4(dg+ef)^2};$$

$$\therefore\ \sin^2 D = \frac{4(dg+ef)^2-(d^2+g^2-e^2-f^2)^2}{4(dg+ef)^2}.$$

The numerator of this expression is equal to

$$\{(2dg+2ef)+(d^2+g^2-e^2-f^2)\}\{(2dg+2ef)-(d^2+g^2-e^2-f^2)\}$$

$$=\{(d+g)^2-(e-f)^2\}\times\{(e+f)^2-(d-g)^2\}$$

$$=\{(d+g)+(e-f)\}\{(d+g)-(e-f)\}$$
$$\times\{(e+f)+(d-g)\}\{(e+f)-(d-g)\}$$

$$=(d+g+e-f)(d+g-e+f)(e+f+d-g)(e+f+g-d).$$

Let 2s stand for $(d+e+f+g)$.

Then the above expression may be written

$$(2s-2f)(2s-2e)(2s-2g)(2s-2d).$$

Therefore $\sin^2 D = \dfrac{4(s-d)(s-e)(s-f)(s-g)}{(dg+ef)^2}$.

or $\sin D = \dfrac{2\sqrt{(s-d)(s-e)(s-f)(s-g)}}{dg+ef}$.

Hence the area of the quadrilateral $DEFG$

which is $\frac{1}{2}(dg+ef)\sin D$

$$=\sqrt{(s-d)(s-e)(s-f)(s-g)}.$$

* 282. *To prove that if* I *and* O *are the centres of the inscribed and circumscribed circles of a triangle, then*

$$OI^2 = R^2 - 2Rr.$$



Let OI produced cut the circumscribing circle in H and K. Join IA, IC; produce IA to cut the circle ABC again in V. Draw the diameter VOW and join CW. Then the angle $VIC = IAC + ICA = \frac{1}{2}A + \frac{1}{2}C = \frac{1}{2}(A + C)$. The angle

$$ICV = ICB + BCV = ICB + BAV = \tfrac{1}{2}C + \tfrac{1}{2}A = \tfrac{1}{2}(A + C).$$

Therefore the angle $VCI = VIC$, and $\therefore\ VI = VC$.

Now $OK^2 - OI^2 = HI \,.\, IK$ [Euc. II. 5]

$= VI \,.\, IA = VC \,.\, IA.$ [Euc. III. 35.]

But $VC = VW \,.\, \sin CWV = 2R \,.\, \sin CAV = 2R \,.\, \sin \dfrac{A}{2}$,

and $$IA = \frac{r}{\sin \frac{A}{2}}.$$ [See Figure on page 234.]

Hence $$OK^2 - OI^2 = 2R \sin \frac{A}{2} \times \frac{r}{\sin \frac{A}{2}},$$

or $$R^2 - OI^2 = 2Rr.$$

EXAMPLES. LXXIII.

(1) Find the area of the triangle ABC when

(i) $a=4$, $b=10$ feet, $C=30^0$.

(ii) $b=5$, $c=20$ inches, $A=60^0$.

(iii) $c=66\frac{2}{3}$, $a=15$ yards, $B=17^0\ 14'$ [$\sin 17^0\ 14'=\cdot29626$].

(iv) $a=13$, $b=14$, $c=15$ chains.

(v) $a=10$, the perpendicular from A on $BC=20$ feet.

(vi) $a=625$, $b=505$, $c=904$ yards.

(2) Find the Radii of the Inscribed and each of the Escribed Circles of the triangle ABC when $a=13$, $b=14$, $c=15$ feet.

(3) Show that the triangles in which (i) $a=2$, $A=60^0$; (ii) $b=\frac{2}{3}.\sqrt{3}$, $B=30^0$ can be inscribed in the same circle.

(4) Prove that $R=\dfrac{abc}{4S}$; find R in the triangle of (2).

(5) Prove that if a series of triangles of equal perimeter are described about the same circle, they are equal in area.

(6) If $A=60^0$, $a=\sqrt{3}$, $b=\sqrt{2}$, prove that the area
$$=\tfrac{1}{4}(3+\sqrt{3}).$$

(7) Prove that each of the following expressions represents the area of the triangle ABC:

(i) $\dfrac{abc}{4R}$. (ii) $2R^2 \sin A\,.\sin B\,.\sin C$.

(iii) rs. (iv) $Rr(\sin A+\sin B+\sin C)$.

(v) $\frac{1}{2}a^2 \sin B\,.\sin C\,.\operatorname{cosec} A$. (vi) $ra\operatorname{cosec}\dfrac{A}{2}\cos\dfrac{B}{2}\cos\dfrac{C}{2}$.

(vii) $(rr_1r_2r_3)^{\frac{1}{2}}$. (viii) $\frac{1}{2}(a^2-b^2)\sin A\,.\sin B\,.\operatorname{cosec}(A-B)$.

Prove the following statements:

(8) If a, b, c are in A. P., then $ac=6rR$.

(9) The area of the greatest triangle, two of whose sides are 50 and 60 feet, is 1500 sq. feet.

(10) If the altitude of an isosceles triangle is equal to the base, R is five-eighths of the base.

(11) $R(\sin A+\sin B+\sin C)=s$.

(12) $bc=4R^2(\cos A+\cos B.\cos C)$.

(13) If C is a right angle, $2r+2R=a+b$.

(14) $r_2r_3+r_3r_1+r_1r_2=s^2$.

(15) $\dfrac{1}{bc}+\dfrac{1}{ca}+\dfrac{1}{ab}=\dfrac{1}{2rR}$.

(16) $r_1\cot\dfrac{A}{2}=r_2\cot\dfrac{B}{2}=r_3\cot\dfrac{C}{2}=r\cot\dfrac{A}{2}.\cot\dfrac{B}{2}.\cot\dfrac{C}{2}$.

(17) $r_1+r_2=c\cot\dfrac{C}{2}$.

(18) $\dfrac{1}{r_1}+\dfrac{1}{r_2}+\dfrac{1}{r_3}=\dfrac{1}{r}$.

(19) $\dfrac{r_1-r}{a}+\dfrac{r_2-r}{b}=\dfrac{c}{r_3}$. (20) $r_1+r_2+r_3-r=4R$.

(21) $a.b.c.r=4R(s-a)(s-b)(s-c)$.

(22) The distances of the centres of the escribed circles of the triangle ABC from that of the inscribed circle are

$$4R\sin\frac{A}{2},\quad 4R\sin\frac{B}{2},\quad 4R\sin\frac{C}{2}.$$

(23) If A is a right angle, $r_2+r_3=a$.

(24) In an equilateral triangle $3R=6r=2r_1$.

(25) $\dfrac{r_1}{bc}+\dfrac{r_2}{ca}+\dfrac{r_3}{ab}=\dfrac{1}{r}-\dfrac{1}{2R}$.

* CHAPTER XX.

On the Area of the Circle, the Construction of Trigonometrical Tables, &c.

283. Let r be the radius of the circle described about a regular polygon of n sides. Let O be the centre; HK one of the sides.



From O draw OD perpendicular to HK, bisecting both HK and the angle HOK. Then, since the polygon has n equal sides, n times the angle HOK make up four right angles

$$\therefore HOK = \frac{2\pi}{n}, \text{ and } \therefore DOK = \frac{\pi}{n}. \quad \text{Hence,}$$

the perimeter of the polygon, which is n times HK,

$$= 2n \,.\, DH = 2n \,.\, OH \sin DOH$$

$$= 2nr \sin \frac{\pi}{n};$$

the area of the polygon, which is n times the area of HOK,

$$= nOD \,.\, HD = nr \cos \frac{\pi}{n} \,.\, r \sin \frac{\pi}{n}$$

$$= nr^2 \sin \frac{\pi}{n} \,.\, \cos \frac{\pi}{n}.$$

284. Let r be the radius of the circle **inscribed** in a regular polygon of n sides. Let O be the centre, HK one of the sides.



From O draw OM perpendicular to HK, bisecting both HK and the angle HOK. Then since the polygon has n equal sides, n times the angle HOK make up four right angles,

$$\therefore HOK = \frac{2\pi}{n} \text{ and } \therefore MOH = \frac{\pi}{n}. \quad \text{Hence}$$

the **perimeter** of the polygon, which is n times HK,

$$= 2n \,.\, MH = 2n \,.\, OM \tan MOH$$

$$= 2nr \tan \frac{\pi}{n}.$$

The **area** of the polygon, which is n times the area of HOK,

$$= n \,.\, OM \,.\, MH = n \,.\, OM \,.\, OM \tan MOH$$

$$= nr^2 \tan \frac{\pi}{n}$$

= the radius × half the perimeter.

285. Let a circle of radius r have a regular polygon of n sides circumscribed about it, and also a regular polygon of the same number (n) of sides inscribed in it.

The **perimeter** of the circumscribing polygon is $2nr \tan \frac{\pi}{n}$.

The **perimeter** of the inscribed polygon is $2nr \sin \frac{\pi}{n}$.

The ratio of these perimeters is $1 : \cos \frac{\pi}{n}$ ..... (1).

The **area** of the circumscribed polygon is $nr^2 \tan \frac{\pi}{n}$.

The **area** of the inscribed polygon is $nr^2 \sin \frac{\pi}{n} \cos \frac{\pi}{n}$.

The ratio of these areas is $1 : \cos^2 \frac{\pi}{n}$ ..... (ii).

286. We must assume the two following axioms:

(i) The circumference of the circle lies in magnitude **between** the perimeter of a circumscribed and that of an inscribed polygon.

(ii) The area of the circle lies in magnitude **between** the area of a circumscribed and that of an inscribed polygon.

Now, when n is increased, $\frac{\pi}{n}$ is diminished, and therefore (by Art. 94) $\cos \frac{\pi}{n}$ approaches 1.

Hence, as the number of the sides of the two polygons in Art. 285 is increased, their perimeters approach to equality.

And since the circumference of the circle always lies in magnitude **between** them, each of these perimeters must approach the circumference of the circle.

Therefore, the **circumference** of a circle is that, to which the perimeter of a regular inscribed (or circumscribed) polygon approaches as the number of its sides is increased, and from which it can be made to differ by less than any assignable quantity however small. (Cf. Art. 32.)

287. In like manner it follows, from (ii) Art. 285, that the area of the circle is the '*limit*' of the area of a regular inscribed (or circumscribed) polygon when the number of the sides is indefinitely increased.

Now, twice the area of a polygon circumscribed about a circle is equal to (the radius × the perimeter). [Art. 284]

This is true however great be the number of the sides. It is therefore true when the number of the sides is indefinitely increased.

Therefore it is true for the circle itself. Hence,

twice the area of a circle = the radius × the circumference.

Or, the area of a circle $= \frac{1}{2}r\,(2\pi r)$ [Art. 34]

$= \pi r^2$.

EXAMPLES. LXXIV.

(1) Prove the surd expressions of Art. 37 for the ratios of the perimeters of certain regular polygons to the diameters of their circumscribing circles.

(2) Prove that the area of a regular polygon of twelve sides described about a circle whose radius is 1 foot is 3·215 sq. ft.

(3) Prove that the area of a square described about a circle is $\frac{4}{3}$ of the dodecagon inscribed in the same circle.

(4) Find the perimeter of a regular polygon of 100 sides (i) when described about a circle of 1 foot diameter, (ii) when inscribed in the same circle. *Ans.* (i) 3·14263, (ii) 3·14108 ft.

(5) An equilateral triangle and a regular hexagon have the same perimeter: show that the areas of their inscribed circles are as 4 : 9.

(6) Prove that the area of a regular polygon of n sides, each of whose sides is a, is $\frac{na^2}{4} \cdot \cot \frac{180^0}{n}$.

(7) If the areas of a regular pentagon and decagon are equal, the ratio of their sides is $\sqrt[4]{20} : 1$.

(8) If a be a side of a regular polygon and R and r the radii of the inscribed and described circles respectively, then $2R = a \operatorname{cosec} \frac{\pi}{n}$ and $2r = a \cot \frac{\pi}{n}$.

(9) If R and r be the radii of the inscribed and circumscribed circles of a regular polygon of n sides, each $= a$,

$$R + r = \frac{a}{2} \cot \frac{\pi}{2n}.$$

(10) The triangle formed from one side each of a regular pentagon, hexagon and decagon inscribed in the same circle, is right-angled.

(11) Prove that the area of an irregular polygon described about a circle is equal to the product of the radius and half the perimeter of the polygon.

(12) The area of an irregular polygon of an even number of sides circumscribed about a circle is equal to the radius × the sum of every alternate side.

(13) The diameter of the dome of St Paul's is 108 feet; prove that it covers an area of 1018 sq. yds.

(14) The radius of the circle whose area is one acre is $39\frac{1}{4}$ yds.

(15) A length of 300 yards of paper, the thickness of which is the hundred and fiftieth part of an inch, is rolled up into a solid cylinder. Find approximately the diameter of the cylinder. *Ans.* 9·575 in.

(16) The diameter of a roll of carpet is 2 feet and the thickness of the carpet is the eighth of an inch. What is the length of the carpet? *Ans.* 301·6 ft.

On the Construction of Trigonometrical Tables.

288. *To prove that, if θ be the circular measure of an angle less than a right angle,* $\sin\theta$, θ, $\tan\theta$ *are in ascending order of magnitude.*



Let ROP be an angle (θ) less than a right angle. Make the angle ROP' on the other side of OR equal to ROP. With centre O and any radius OR describe the arc $P'RP$. Draw PT, $P'T$ to touch the circle at P and P'. PT and $P'T$ will meet on OR. The line PP' is bisected by OR at right angles at M.

Then, since the circumference of a circle lies *between* the perimeter of the inscribed and circumscribed polygons, it follows that the arc $P'RP$ lies in magnitude between $P'MP$ and $P'T + TP$. In other words, $P'MP$, the arc $P'RP$, and $P'T + TP$ are in ascending order of magnitude.

Therefore also their halves MP, RP, TP are in ascending order of magnitude.

And so also are $$\frac{MP}{OP}, \frac{RP}{OP}, \frac{TP}{OP}.$$

That is, $\sin\theta$, θ, $\tan\theta$ are in ascending order of magnitude, where $\theta \left[= \frac{\text{arc}}{\text{radius}}\right]$ is the circular measure of the angle referred to.

289. Hence, 1, $\frac{\theta}{\sin\theta}$, $\frac{1}{\cos\theta}$ are in ascending order of magnitude.

290. *To prove that the 'limit' of* $\frac{\theta}{\sin\theta}$, *when* $\theta = 0$, *is* 1; *θ being the circular measure of the angle referred to.*

The value of $\frac{\theta}{\sin\theta}$ lies between 1 and $\frac{1}{\cos\theta}$. As θ is diminished, $\cos\theta$ approaches 1; and the smaller θ becomes, the more nearly does $\frac{1}{\cos\theta}$ approach 1. [Art. 94]

Therefore, by diminishing θ sufficiently, we can make $\frac{\theta}{\sin\theta}$ differ from 1 by less than any assignable quantity however small.

This is what is meant when it is said that '*the limit of* $\frac{\theta}{\sin\theta}$, *when* $\theta = 0$, *is* 1.'

291. The student must notice carefully that θ here is the **number of radians** in the angle referred to.*

Example. Prove that the limit of $\frac{n}{\sin n^0}$, *when* $n = 0$, *is* $\frac{180}{\pi}$.

Let θ radians $= n^0$, then

$$\frac{\theta}{\pi} = \frac{n}{180}, \quad \therefore n = \frac{180\theta}{\pi}, \quad \therefore \frac{n}{\sin n^0} = \frac{180}{\pi} \cdot \frac{\theta}{\sin\theta}.$$

When n is diminished, θ is diminished also, and the limit of $\frac{\theta}{\sin\theta}$, when $\theta = 0$, is 1. Therefore the limit of $\frac{n}{\sin n^0}$ $\left(\text{which} = \frac{180}{\pi} \cdot \frac{\theta}{\sin\theta}\right)$, when $n = 0$, is $\frac{180}{\pi}$.

* It is on this account that a radian is used as a unit of angle (Art. 62).

292. *To prove that if θ be the circular measure of a positive angle less than a right angle,* $\sin\theta$ *lies between* θ *and* $\theta - \frac{1}{4}\theta^3$.

It has been proved that $\sin\theta$ is less than θ ..... (i).

And that θ is less than $\tan\theta$.

Therefore $\frac{\theta}{2}$ is less than $\tan\frac{\theta}{2}$; or, $\frac{\theta}{2}\cos\frac{\theta}{2}$ is less than $\sin\frac{\theta}{2}$.

Now, $\sin\theta = 2\sin\frac{\theta}{2}\cdot\cos\frac{\theta}{2}$; [Art. 166]

$$\therefore \sin\theta \text{ is greater than } 2\left(\frac{\theta}{2}\cos\frac{\theta}{2}\right)\cos\frac{\theta}{2},$$

i.e. greater than $\theta\cos^2\frac{\theta}{2}$, i.e. greater than $\theta - \theta\sin^2\frac{\theta}{2}$.

But $\left(\frac{\theta}{2}\right)^2$ is greater than $\sin^2\frac{\theta}{2}$.

Still greater therefore is $\sin\theta$ than $\theta - \frac{1}{4}\theta^3$.....(ii).

Hence, $\sin\theta$ is less than θ, and greater than $\theta - \frac{\theta^3}{4}$.

293. *To find* $\sin 10''$.

In the above, θ is the *circular measure* of the angle.

The circular measure of $10''$, correct to three significant figures, is $\cdot 0000484....$ [Examples X. (17).]

Let $\theta = \cdot 0000484....$

Then,

$$\theta - \tfrac{1}{4}\theta^3 = \theta - \tfrac{1}{4}(\cdot 0000484....)^3 = \theta - \cdot 000000000000028....$$

Hence, θ and $(\theta - \frac{1}{4}\theta^3)$ are decimal fractions which agree in their first twelve figures at least.

And since $\sin 10''$ lies *between* these fractions, therefore the first twelve decimal places of $\sin 10''$ are the same as those of the circular measure of $10''$.

Hence, if the value of π be given to a sufficient number of decimal places, we can calculate the circular measure of $10''$, and therefore also $\sin 10''$, to 12 decimal places.

294. *To show how to construct a Table giving the Trigonometrical Ratios of angles which form an arithmetical progression having* $10''$ *for the common difference.*

In the identity

$$\sin(n+1)\alpha + \sin(n-1)\alpha = 2\sin n\alpha \,.\, \cos\alpha,$$

let $\alpha = 10''$, and suppose a table of the sines of all angles at intervals of $10''$ to have been calculated up to $n\,.\,10''$.

Then, $\sin(n-1)\alpha$, $\sin n\alpha$ and $\cos\alpha \left[= \sqrt{1-\sin^2 10''}\right]$ are known.

Therefore by the above formula $\sin(n+1)\alpha$ can be found.

Hence, since we know $\sin 10''$, the sine of $20''$ can be found; and then the value of $\sin 30''$; and so on.

295. When the sines of angles up to 45° have been calculated, the rest may be found from the formula,

$$\sin(45^\circ + A) - \sin(45^\circ - A) = \sqrt{2} \,.\, \sin A.$$

Also, when the sines of angles up to 60° have been calculated, the remainder up to 90° can be found still more easily from the formula, $\sin(60^\circ + A) - \sin(60^\circ - A) = \sin A$.

The other ratios may be found from the following:

$$\cos A = \sin(90^\circ - A), \quad \tan A = \frac{\sin A}{\cos A}, \quad \cot A = \tan(90^\circ - A),$$

$$\operatorname{cosec} A = \frac{1}{\sin A}, \quad \sec A = \operatorname{cosec}(90^\circ - A).$$

296. Since $3^\circ = 18^\circ - 15^\circ$, the sines of angles differing by 3°, or by any multiple of 3°, can be found independently. (See p. 120.) These values may be used to *test* the accuracy of the Tables calculated as above.

297. The following formulæ may be also used to test the accuracy of the Tables,

$$\sin(36^\circ+A)-\sin(36^\circ-A)-\sin(72^\circ+A)+\sin(72^\circ-A)=\sin A,$$

$$\cos(36^\circ+A)+\cos(36^\circ-A)-\cos(72^\circ+A)-\cos(72^\circ-A)=\cos A.$$

They are called **formulæ of verification.**

On the Limit of the Visible Horizon.

298. The surface of the sea is very nearly that of a sphere whose radius is 3957 miles.

The height of the highest mountains on the globe is less than 6 miles. Thus a point must be considered to be at a very considerable height above the surface of the sea if its height is a thousandth part of the earth's radius.

299. In the figure, let O be the centre of the earth, PRP' part of the surface of the sea, T a point of observation, TR its vertical height.

Draw TP, TP' tangents to the earth's surface. Then PMP' is parallel to the 'horizontal plane' at R.

The angle TPM is called the **dip of the horizon** at T. It is the angle of depression of the most distant point on the horizon seen from T. It must obviously be a very small angle, since TR is so small compared with RO.



300. If RO be produced to cut the circle again in L, then

$$TP^2 = TR \,.\, TL \qquad \text{[Euc. III. 36]}$$

$$= TR\,(RL + TR) = 2TR \,.\, RO + TR^2.$$

But TR will in general be much less than a thousandth part of RO, and therefore TR^2 will be much less than a thousandth part of $2TR \,.\, RO$.

Hence, the formula $TP^2 = 2TR \,.\, RO$, i.e. TP^2 = twice the earth's radius × vertical height, will give the value of TP correct to at least three significant figures.

Example. Three times the height in feet *of the place of observation above the sea is equal to twice the square of the distance of the horizon* in miles.

Here, $TP^2 = RL \times 7914$ miles.

Let f be the number of feet in RL, then the number of miles in RL is $\frac{f}{5280}$; let x be the number of miles in TP, then

$$x^2 = \frac{f}{5280} \times 7914 = \frac{3f}{2} \text{ nearly.} \quad \text{Q.E.D.}$$

EXAMPLES. LXXV.

(1) Show that the limit of $\frac{n}{2} R^2 \sin \frac{2\pi}{n}$ (i.e. the area of a polygon of n sides inscribed in a circle of radius R), when $n = \infty$ is πR^2.

(2) Prove that the limit of $nr^2 \tan \frac{\pi}{n}$, when $n = \infty$, is πr^2.

(3) Given that $\pi = 3{\cdot}141592653589793\ldots$ prove that the circular measure of 10″ is ·00004848136811...

(4) Prove that $2 \sin (72^0 + A) - 2 \sin (72^0 - A) = (\sqrt{5} - 1) \sin A$, and that $2 \sin (36^0 + A) - 2 \sin (36^0 - A) = (\sqrt{5} + 1) \sin A$.

(5) If a mast of a ship be 150 feet high, show that the greatest distance seen from its top is 15 miles nearly.

(6) Prove that if the dip of the horizon at the top of a mountain is $1^0\,26'$ [$= \tan^{-1} {\cdot}025$], the mountain is about 6530 feet high.

NOTE. The definitions given in Arts. 75, 78 of the Trigonometrical Ratios are now used exclusively.

The NAMES *tangent*, *secant*, *sine*, were given originally to quantities defined as follows.

Let *ROP* be any angle. With centre *O* and any radius describe the arc *RP*. Draw *PM* perpendicular to *OR* and *PT* perpendicular to *OP*. (See Figure on previous page.)

Then *PR* is called an *arc*, *PT* is the *tangent of the arc PR*, *OT* is the *secant of the arc PR*, *MP* is the *sine of the arc PR*.

The name *sine* is derived from the word *sinus*. For, in the figure, *PMP'* is the string of the "bow" (*arcus*), and the string of a bow when in use is pulled to the archer's *breast*.

The *co*-tangent, *co*-secant and *co*-sine are respectively the tangent, secant and sine of the *complement* of the arc or of the angle.

The sine, tangent, etc. of the *angle* are the same as the *measures* of the sine, tangent, etc. of the *arc*, when the radius of the circle is the unit of length.

EXAMPLES FOR EXERCISE. LXXVIa.

1. Define the terms sine, cotangent; and prove that if A be *any* angle, $\sin^2 A + \cos^2 A = 1$.

If $\tan A = \frac{3}{4}$, find $\sin A$ and $\cos A$.

2. Trace the changes in the sign and magnitude of $\cos\theta - \sec\theta$ for values of θ between 0 and π.

3. Prove geometrically that $\cos(180^0 - A) = -\cos A$.

Find A if $2\sin A = \tan A$.

4. Prove

(1) $\sin(A+B) \cdot \sin(A-B) = \sin^2 A - \sin^2 B$;

(2) $\dfrac{\sin A + \sin B}{\sin A - \sin B} = \dfrac{\tan \frac{1}{2}(A+B)}{\tan \frac{1}{2}(A-B)}$.

5. Prove that

$$\cos^2 A - \cos A \cos(60^0 + A) + \sin^2(30^0 - A) = \tfrac{3}{4}.$$

6. Find the greatest side of the triangle of which one side is 2183 feet and the adjacent angles are $78^0\,14'$ and $71^0\,24'$.

$\log 2183 = 3{\cdot}3390537$, $\log 42274 = 4{\cdot}6260733$,
$L\sin 78^0\,14' = 9{\cdot}9907766$, $\log 42275 = 4{\cdot}6260836$.
$L\sin 30^0\,22' = 9{\cdot}7037486$,

---

7. Express the other trigonometrical ratios in terms of the cosine.

8. Prove $\sin(180 + A) = -\sin A$;
$\tan(90 + A) = -\cot A$.

9. Write down the sines of all the angles which are multiples of 30^0 and less than 360^0.

10. Prove $\tan^2 A = \dfrac{1 - \cos 2A}{1 + \cos 2A}$.

11. If $\tan A + \sec A = 2$, prove that $\sin A = \frac{3}{5}$, when A is less than 90^0.

If $\sin A = \frac{4}{5}$, prove that $\tan A + \sec A = 3$, when A is less than 90^0.

12. The length of the greatest side of a triangle is 1035·43 feet, and the three angles are 44°, 66°, and 70°. Solve the triangle, having given

$L \sin 44^0 = 9{\cdot}8417713$, $L \sin 66^0 = 9{\cdot}9607302$,
$L \sin 70^0 = 9{\cdot}9729858$, $\log 1035{\cdot}43 = 3{\cdot}0151212$,
$\log 765432 = 5{\cdot}8839067$, $\log 10066 = 4{\cdot}0028656$.

---

13. Express the other trigonometrical ratios in terms of the cotangent.

14. Prove that $\cos(180^0 - A) = -\cos A$;
$\operatorname{cosec}(180^0 + A) = -\operatorname{cosec} A$.

15. Write down the tangents of all the angles which are multiples of 30° and less than 360°.

16. If $\tan A + \sec A = 3$, prove that $\sin A = \frac{4}{5}$ when A is less than 90°.

If $\sin A = \frac{4}{5}$, prove that $\tan A + \sec A = 2$, when A is less than 90°.

17. Find the sines of the three angles of the triangle whose sides are 193, 194, and 195 feet.

18. Investigate the following formulæ:

(1) $\cos \frac{3A}{2} = (2 \cos A - 1) \cos \frac{1}{2}A$;

(2) $\cos\theta - \cos(\theta + \delta) = \sin\theta \sin\delta \,(1 + \cot\theta \tan \frac{1}{2}\delta)$.

---

19. Define the secant of an angle.

Prove the formula $\dfrac{1}{\sec^2 A} + \dfrac{1}{\operatorname{cosec}^2 A} = 1$.

If $\sin A = \frac{1}{3}$, find $\sec A$.

20. Trace the changes in sign and magnitude of

$$\frac{\sin\theta + \cos\theta}{\sin\theta - \cos\theta},$$ as θ changes from π to 2π.

21. Find a formula to include all angles that have the same cotangent as the angle α.

Solve the equation $\tan\theta = \cot\theta$.

22. Prove the formula to express the cosine of the sum of two angles in terms of the sines and cosines of those angles.

Express $\cos 5\alpha$ in terms of $\cos\alpha$.

23. Prove the formula

$$2\sin\tfrac{1}{2}A = \pm\sqrt{(1+\sin A)} \pm \sqrt{(1-\sin A)}.$$

Account for the double signs in this formula, and examine which signs must be taken if A be an angle between 540^0 and 630^0.

24. A ring 10 inches in diameter is suspended from a point 1 foot above its centre by six equal strings attached to its circumference at equal intervals; find the cosine of the angle between two consecutive strings.

---

25. Define 1^0. Assuming that $\frac{22}{7}$ is the circular measure of two right angles, express the angle A^0 in circular measure.

Find the number of degrees in the angle whose circular measure is $\cdot 1$.

26. Find the trigonometrical ratios of the angle whose cosine is $\frac{3}{5}$.

27. Prove that

(1) $\cos(180^0+A) = \cos(180^0-A)$;
(2) $\tan(90^0+A) = \cot(180^0-A)$.

28. Prove $\sin x(2\cos x - 1) = 2\sin\frac{x}{2}\cos\frac{3x}{2}$.

29. Trace the changes in sign and magnitude of

$$\frac{2\sin\theta - \sin 2\theta}{2\sin\theta + \sin 2\theta},$$

as θ changes from 0 to 2π.

30. If the angle opposite the side a be 60^0, and if b, c be the remaining sides of the triangle, prove that

$$(a+b+c)(b+c-a) = 3bc.$$

---

31. Assuming $\frac{22}{7}$ to be the circular measure of two right angles, express in degrees the angle whose circular measure is θ. Find the number of degrees in an angle whose circular measure is $\frac{1}{3}$.

32. Prove that $\tan^{-1}x - \tan^{-1}y = \tan^{-1}\frac{x-y}{1+xy}$.

33. Prove $\sin x(2\cos x + 1) = 2\cos\frac{x}{2}\sin\frac{3x}{2}$.

34. Trace the changes in sign and magnitude of

$$\frac{\sin\theta + 2\sin\frac{1}{2}\theta}{\sin\theta - 2\sin\frac{1}{2}\theta},$$

as θ changes from 0 to 2π.

35. If $(\sin A+\sin B+\sin C)(\sin A+\sin B-\sin C)=3\sin A\sin B$, and $A+B+C=180^0$, prove that $C=60^0$.

36. Given $A=18^0$, $B=144^0$, and $b=1$, solve the triangle.

---

37. Give the trigonometrical definition of an angle.

What angle does the minute-hand of a clock describe between twelve o'clock and 20 minutes to four?

38. Express the cosine and the tangent of an angle in terms of the sine.

The angle A is greater than 90^0 but less than 180^0, and $\sin A=\frac{1}{3}$. Find $\cos A$.

39. Find an expression for all the values of θ for which

$$\cos\theta+\cos 2\theta=0.$$

40. If in a triangle $a\cos A=b\cos B$, the triangle will be either isosceles or right-angled.

41. The sides are 1 foot and $\sqrt{3}$ feet respectively, and the angle opposite to the shorter side is 30^0; solve the triangle.

42. The sides of a triangle are 2, 3, 4. Find the greatest angle, having given

$$\log 2= \cdot 3010300,$$
$$\log 3= \cdot 4771213,$$
$$L\tan 52^0.15'=10\cdot 1111004,$$
$$L\tan 52^0.14'=10\cdot 1108395.$$

---

43. Distinguish between Euclid's definition of an angle and the trigonometrical definition.

What angle does the minute-hand of a clock describe between half-past four and a quarter-past six?

44. Express the sine and the cosine of an angle in terms of the tangent.

The angle A is greater than 180^0 but less than 270^0, and $\tan A=\frac{1}{2}$. Find $\sin A$.

45. Prove (i) $\sin 2A=\dfrac{2\cot A}{1+\cot^2 A}$.

(ii) Show that if $A+B+C=90^0$,

$$\sin 2A+\sin 2B+\sin 2C=4\cos A\cos B\cos C.$$

46. Find an expression for all the values of θ for which

$$\sin\theta+\sin 2\theta=0.$$

47. If in a triangle $b\cos A=a\cos B$, show that the triangle is isosceles.

48. The sides are 1 foot and $\sqrt{2}$ feet respectively, and the angle opposite to the shorter side is 30^0; solve the triangle.

---

49. Express in degrees, minutes, etc. (1) the angle whose circular measure is $\frac{1}{20}\pi$; (2) the angle whose circular measure is 5.

If the angle subtended at the centre of a circle by the side of a regular heptagon be the unit of angular measurement, by what number is an angle of 45^0 represented?

50. Prove that

$$(\sin 30^0 + \cos 30^0)(\sin 120^0 + \cos 120^0) = \sin 30^0.$$

51. Prove the formulæ:

(1) $\cos^2(\alpha+\beta) - \sin^2\alpha = \cos\beta\cos(2\alpha+\beta)$;

(2) $1 + \cot\alpha\cot\frac{1}{2}\alpha = \operatorname{cosec}\alpha\cot\frac{1}{2}\alpha$.

52. Solve the equations:

(1) $5\tan^2 x - \sec^2 x = 11$; (2) $\sin 5\theta - \sin 3\theta = \sqrt{2}\,.\cos 4\theta$.

53. Two sides of a triangle are 10 feet and 15 feet in length, and the angle between them is 30^0. What is its area?

54. Given that

$$\sin 40^0\,29' = 0{\cdot}6492268, \quad \sin 40^0\,30' = 0{\cdot}6494480,$$

find $\sin^{-1}(0{\cdot}6493000)$.

---

55. Express in circular measure (1) 10′, (2) $\frac{1}{5}$ of a right angle.

If the angle subtended at the centre of a circle by the side of a regular pentagon be the unit of angular measurement, by what number is a right angle represented?

56. If $\sec\alpha = 7$, find $\tan\alpha$ and $\operatorname{cosec}\alpha$.

57. Prove the formulæ:

(1) $\cos^2(\alpha-\beta) - \sin^2(\alpha+\beta) = \cos 2\alpha\cos 2\beta$;

(2) $1 + \tan\alpha\tan\frac{1}{2}\alpha = \sec\alpha$.

58. Solve the equations:

(1) $5\tan^2 x + \sec^2 x = 7$; (2) $\cos 5\theta + \cos 3\theta = \sqrt{2}\,.\cos 4\theta$.

59. The lengths of the sides of a triangle are 3 feet, 5 feet, and 6 feet. What is its area?

60. Given that

$$\sin 38^0\,25' = 0{\cdot}6213757, \quad \sin 38^0\,26' = 0{\cdot}6216036,$$

find $\sin^{-1}(0{\cdot}6215000)$.

---

61. Which is greater, 70^g or the angle whose circular measure is 1·2?

62. Determine geometrically $\cos 30^0$ and $\cos 45^0$.

If $\sin A$ be the arithmetic mean between $\sin B$ and $\cos B$, then

$$\cos 2A = \cos^2(B + 45^0).$$

63. Establish the following relations:

(1) $\tan^2 A - \sin^2 A = \tan^2 A \sin^2 A$;

(2) $\cot A - \cot 2A = \operatorname{cosec} 2A$;

(3) $$\frac{\sin(x+3y) + \sin(3x+y)}{\sin 2x + \sin 2y} = 2\cos(x+y).$$

64. Show that for certain values of the angles

$$2\sin \tfrac{1}{2}A = \sqrt{1+\sin A} - \sqrt{1-\sin A}.$$

Is the formula true when $A = 240^0$? If not, how must it be modified?

65. Prove that $\sin(A+B) = \sin A \cos B + \cos A \sin B$, and deduce the expression for $\cos(A+B)$.

Show that

$$\sin A \cos(B+C) - \sin B \cos(A+C) = \sin(A-B)\cos C.$$

66. One side of a triangular lawn is 102 feet long, its inclinations to the other sides being $70^0\,30'$, $78^0\,10'$ respectively. Determine the other sides and the area. $L\sin 70^0\,30' = 9{\cdot}974$, $\log 102 = 2{\cdot}009$, $L\sin 78^0\,10' = 9{\cdot}990$, $\log 185 = 2{\cdot}267$, $L\sin 31^0\,20' = 9{\cdot}716$, $\log 192 = 2{\cdot}283$, $\log 2 = {\cdot}301$, $\log 9234 = 3{\cdot}965$.

---

67. Which is greater, 126^0 or the angle whose circular measure is 2·3?

68. Establish the following relations:

(1) $\cot^2 A - \cos^2 A = \cot^2 A \cos^2 A$;

(2) $\tan A + \cot 2A = \operatorname{cosec} 2A$;

(3) $$\frac{\cos(x-3y) - \cos(3x-y)}{\sin 2x + \sin 2y} = 2\sin(x-y).$$

69. Show that for certain values of the angles

$$2\cos \tfrac{1}{2}A = \sqrt{1+\sin A} + \sqrt{1-\sin A}.$$

Is the formula true when $A = 300^0$? If not, how must it be modified?

70. Prove that $\sin 30^0 + \sin 120^0 = \sqrt{2}\cos 15^0$.

71. Establish the identities:

(1) $1 + \cos A + \sin A = \sqrt{2(1+\cos A)(1+\sin A)}$;

(2) $$\operatorname{cosec} 2A = \frac{\operatorname{cosec}^2 A}{2\sqrt{\operatorname{cosec}^2 A - 1}};$$

(3) $\sin\frac{2\pi}{7}+\sin\frac{4\pi}{7}-\sin\frac{6\pi}{7}=4\sin\frac{\pi}{7}\sin\frac{3\pi}{7}\sin\frac{5\pi}{7}$.

72. The sides of a triangular lawn are 102, 185, and 192 feet in length, the smallest angle being approximately $31^\circ 20'$. Find its other angles and its area.

$\log 102=2{\cdot}009$, $\quad L\sin 31^\circ 20'=9{\cdot}716$,
$\log 185=2{\cdot}267$, $\quad L\sin 70^\circ 30'=9{\cdot}974$,
$\log 192=2{\cdot}283$, $\quad L\sin 78^\circ 10'=9{\cdot}990$,
$\log 2=\cdot 301$, $\log 9234=3{\cdot}965$.

---

73. If the circumference of a circle be divided into five parts in arithmetical progression, the greatest part being six times the least, express in radians the angle each subtends at the centre.

74. Define the sine of an angle, wording your definition so as to include angles of any magnitude.

Prove that $\sin(90^\circ+A)=\cos A$,
and $\cos(90^\circ+A)=-\sin A$,
and by means of these deduce the formulæ
$\sin(180^\circ+A)=-\sin A$, $\cos(180^\circ+A)=-\cos A$.

75. Prove the formulæ:
(1) $\cot^2 A=\operatorname{cosec}^2 A-1$;
(2) $\cot^4 A+\cot^2 A=\operatorname{cosec}^4 A-\operatorname{cosec}^2 A$.
Verify (2) when $A=30^\circ$.

76. Show that all angles satisfying the equation
$$\cos\theta=\cos\alpha$$
are included in the formula $\theta=2n\pi\pm\alpha$.

Solve completely the equation $2\cos^2\theta+\sin^2\theta-1=0$.

77. If $\sin B$ be the geometric mean between $\sin A$ and $\cos A$, then
$$\cos 2B=2\cos^2(A+45^\circ).$$

78. The lengths of two of the sides of a triangle are 1 foot and $\sqrt{2}$ feet respectively, the angle opposite the shorter side is 30°. Prove that there are two triangles which satisfy these conditions; find their angles, and show that their areas are in the ratio $\sqrt{3}+1 : \sqrt{3}-1$.

---

79. If the circumference of a circle be divided into six parts in arithmetical progression, the greatest being six times the least, express in radians the angle each subtends at the centre.

80. Define the tangent of an angle, wording your definition so as to include angles of any magnitude.

Prove that $\tan(90^\circ+A)=-\cot A$, and by means of this formula deduce the formula $\tan(180^\circ+A)=\tan A$.

81. Show that all angles satisfying the equation

$$\tan\theta = \tan\alpha$$

are included in the formula $\theta = n\pi + \alpha$.

Solve completely the equation $\sec^3\theta - 2\tan^2\theta = 2$.

82. Prove that $\cos(A+B) = \cos A\cos B - \sin A\sin B$, and deduce the expression for $\sin(A+B)$.

Show that

$$\cos A\cos(B+C) - \cos B\cos(A+C) = \sin(A-B)\sin C.$$

83. Establish the identities:

(1) $1 + \cos A - \sin A = \sqrt{2(1+\cos A)(1-\sin A)}$;

(2) $\sec 2A = \dfrac{\sec^2 A}{2 - \sec^2 A}$;

(3) $\cos\dfrac{2\pi}{7} + \cos\dfrac{4\pi}{7} + \cos\dfrac{6\pi}{7} + 4\cos\dfrac{\pi}{7}\cos\dfrac{3\pi}{7}\cos\dfrac{5\pi}{7} + 1 = 0.$

84. Two adjacent sides of a parallelogram 5 in. and 3 in. long respectively, include an angle of 60^0. Find the lengths of the two diagonals and the area of the figure.

---

85. Investigate the following formulæ:

(1) $\sin\dfrac{3A}{2} = (1 + 2\cos A)\sin\tfrac{1}{2}A$;

(2) $\sin(\theta+\delta) - \sin\theta = \cos\theta\sin\delta\,(1 - \tan\theta\tan\tfrac{1}{2}\delta)$.

86. Prove that

(1) $\sin 10^0 + \sin 50^0 = \sin 70^0$;

(2) $\sqrt{3} + \tan 40^0 + \tan 80^0 = \sqrt{3}\tan 40^0\tan 80^0$;

(3) if $A + B + C = 180^0$,

$$\frac{\sin A - \sin B\cos C}{\cos B} = \frac{\sin B - \sin A\cos C}{\cos A}.$$

87. Find the value of $\sin 18^0$, and deduce that

$$4\sin 18^0\cos 36^0 = 1.$$

88. The length of one side of a triangle is 1006·62 feet and the adjacent angles are 44^0 and 70^0. Solve the triangle, having given

$L\sin 44^0 = 9{\cdot}8417713$,	$L\sin 70^0 = 9{\cdot}9729858$,
$L\sin 66^0 = 9{\cdot}9607302$,	$\log 1006{\cdot}62 = 3{\cdot}0028656$,
$\log 7654321 = 6{\cdot}8839067$,	$\log\ 103543 = 5{\cdot}0151212$.

---

89. Find the length of the arc of a circle whose radius is 8 feet which subtends at the centre an angle of 50^0, having given

$$\pi = 3{\cdot}1416.$$

90. Prove that $\sin A = -\sin(A - 180^0)$.
Find the sines of 30^0 and 2010^0.

91. Investigate an expression for all angles which have a given cosecant.
Write down the general value of $\operatorname{cosec}^{-1}(-\sqrt{2})$.

92. Prove that

(1) $\cos^2 A + \cos^2 B - 2\cos A \cos B \cos(A+B) = \sin^2(A+B)$;
(2) $\cos^2 A + \sin^2 A \cos 2B = \cos^2 B + \sin^2 B \cos 2A$.

93. Prove that in any triangle

$$a^2 \cos 2B + b^2 \cos 2A = a^2 + b^2 - 4ab \sin A \sin B.$$

94. If $a = 123$, $B = 29^0 17'$, $C = 135^0$, find c, having given

$\log 123 = 2{\cdot}0899051$, $\log 2 = {\cdot}3010300$,
$\log 3211 = 3{\cdot}5066403$, diff. for $1 = 1352$.
$L \sin 15^0 43' = 9{\cdot}4327777$.

---

95. Define the unit of circular measure, and prove that it is an invariable angle.
If an arc of 12 feet subtend at the centre of a circle an angle of 50^0, what is the radius of the circle, π being equal to $3{\cdot}1416$?

96. Express the cosine and cotangent in terms of the cosecant.
If $\cot A + \operatorname{cosec} A = 5$, find $\cos A$.

97. Investigate an expression for all angles which have a given secant.
Write down the general value of $\sec^{-1}(-2)$.

98. Prove that

(1) $\sin^2 A + \sin^2 B + 2 \sin A \sin B \cos(A+B) = \sin^2(A+B)$;
(2) $\sin^2 A - \cos^2 A \cos 2B = \sin^2 B - \cos^2 B \cos 2A$.

99. Prove that $2\cos \tfrac{1}{2}A = \pm\sqrt{1+\sin A} \pm \sqrt{1-\sin A}$, and determine the proper signs of the roots when $A = 2400^0$.

100. Prove that in any triangle

$$\frac{\cos 2A}{a^2} - \frac{\cos 2B}{b^2} = \frac{1}{a^2} - \frac{1}{b^2}.$$

---

101. Prove (i) $2\cot 2A = \cot A - \tan A$,
(ii) $\sin^{-1}\frac{3}{5} - \sin^{-1}\frac{5}{13} = \sin^{-1}\frac{16}{65}$,
(iii) $\cot(A+15^0) - \tan(A-15^0) = \dfrac{4\cos 2A}{2\sin 2A + 1}$.

102. Solve the equations

$$\cos(2x+3y)=\tfrac{1}{2},\ \cos(3x+2y)=\tfrac{1}{2}\sqrt{3}.$$

103. Trace the changes in the sign and magnitude of $\sin(\pi \sin x)$ as x increases from 0 to 2π.

104. Solve the equation

$$\tan^{-1}x+\tan^{-1}(1-x)=2\tan^{-1}\sqrt{x-x^2}.$$

105. A ship sailing due north observes two lighthouses bearing respectively N.E. and N.N.E.; after sailing 20 miles the lighthouses are seen to be in a line due east; find the distance between the lighthouses in miles accurately to four places of decimals; having given,

$\log 2=\cdot3010300$, $L\tan 22^\circ30'=9\cdot6172243$, $\log 11\cdot715=1\cdot0687423$,

and $\log 11\cdot716=1\cdot0687794$.

---

106. Prove (i) $\tan\frac{\theta}{2}=\sqrt{\left(\frac{1-\cos\theta}{1+\cos\theta}\right)}$,

(ii) $\frac{\tan 5\theta+\tan 3\theta}{\tan 5\theta-\tan 3\theta}=4\cos 2\theta\cos 4\theta$,

(iii) $\sin^{-1}\frac{8}{5}+\sin^{-1}\frac{8}{17}=\sin^{-1}\frac{77}{85}$.

107. Solve the equation $2\sin^2\theta-(1+\sqrt{3})\sin 2\theta+2\sqrt{3}\cos^2\theta=0$.

108. Show that in any triangle

(i) $a\sin(B-C)+b\sin(C-A)+c\sin(A-B)=0$,

(ii) $c(\cos A+\cos B)=2(a+b)\sin^2\frac{1}{2}C$.

109. Prove that if D, E, F are the feet of the perpendiculars from A, B, C upon the opposite sides of the triangle ABC the diameters of the circumscribing circles of the triangles AEF, BDF, CDE are $a\cot A$, $b\cot B$, $c\cot C$ respectively.

110. A man who is walking on a horizontal plane towards a tower observes that at a certain point the elevation of the top of the tower is 10° and after going 50 yds. nearer to the tower the elevation is 15°; find the height of the tower in yards to 4 places of decimals, having given $L\sin 15^\circ=9\cdot4129962$, $L\cos 5^\circ=9\cdot9983442$,

$\log 25\cdot783=1\cdot4113334$, $\log 25\cdot784=1\cdot4113503$.

---

111. Prove that when $\sin A$ is a geometric mean between $\sin B$ and $\cos B$, then $\cos 2A=2\sin(45^\circ-B)\cos(45^\circ+B)$.

112. Prove (i) $\sin A(\cos 2A+\cos 4A+\cos 6A)=\sin 3A\cos 4A$,

(ii) $2\sin^{-1}\frac{1}{2}=\cos^{-1}\frac{1}{2}$.

113. In any quadrilateral $ABCD$ prove that

$$AB\cos A-BC\cos(A+B)+CD\cos D=AD.$$

114. Prove that in any triangle

$$\tan 2A + \tan 2B + \tan 2C = \tan 2A \tan 2B \tan 2C,$$

hence show that if xyz are numbers such that

$$x + y + z = xyz,$$

then $x(1-y^2)(1-z^2) + y(1-z^2)(1-x^2) + z(1-x^2)(1-y^2) = 4xyz.$

115. From each of two ships a mile apart the angle is observed which is subtended by the other ship and a beacon on shore, these angles are $52^0\,25'\,15''$ and $75^0\,9'\,30''$; find the distances of the beacon from each of the ships, having given $L \sin 75^0\,9'\,30'' = 9{\cdot}9852635$, $L \sin 52^0\,25'\,15'' = 9{\cdot}8990055$, $\log 1{\cdot}2197 = {\cdot}0862530$, $\log 1{\cdot}2198 = {\cdot}0862886$.

---

116. The cosines of two angles of a triangle are $\frac{3}{5}$ and $\frac{12}{13}$ respectively; find all the trigonometrical ratios of the third angle.

117. Prove (i) $\dfrac{\sin A + 2\sin 3A + \sin 5A}{\cos A - 2\cos 3A + \cos 5A} = \dfrac{4\sin A - 3\operatorname{cosec} A}{4\cos A - 3\sec A}$,

(ii) $\cot^{-1}\frac{1}{3} = \cot^{-1} 3 + \cot^{-1}\frac{3}{4}$.

118. The lengths of the side of a triangle are 242, 1212 and 1450 yards; show that the area is 6 acres.

119. Prove that $ad_1^2 + bd_2^2 + cd_3^2 = abc$, where d_1, d_2, d_3 are the distances of the centre of the inscribed circle from A, B, C.

120. A man observes that when he has walked c feet up an inclined plane, the angular depression of an object in the horizontal plane through the foot of the slope is α; and that when he has walked a further distance c feet the angular depression of the same object is β; show that the inclination of the slope to the horizon is

$$\cot^{-1}(2\cot\beta - \cot\alpha).$$

---

121. Prove $\tan^{-1}(\frac{1}{2}\tan 2A) + \tan^{-1}(\cot A) + \tan^{-1}(\cot^3 A) = 0$.

122. The general solution of $\tan(45 - \frac{1}{2}x) + \cot(45^0 - \frac{1}{2}x) = 4$ is $(6n \pm 1)\,60^0$, n being any integer.

123. Prove that $\sqrt{(1+\sin A)} = 1 + 2\sin\frac{1}{4}A\sqrt{(1-\sin\frac{1}{2}A)}$, when A lies between -90^0 and 180^0.

124. If $a\sin A + b\sin B + c\sin C = 0$,

and $a\cos A + b\cos B + c\cos C = 0$,

then $a : b : c = \sin(B-C) : \sin(C-A) : \sin(A-B)$.

125. A man ascends a mountain by a direct course, the inclination of his path to the horizon being at first α and afterwards changing suddenly to β, which continues to the summit; the mountain being a feet high and the angle of elevation of the summit from the starting point being γ, show that the length of the ascent is

$$a\frac{\cos\{\frac{1}{2}(\alpha+\beta)-\gamma\}}{\sin\gamma\cos\frac{1}{2}(\beta-\alpha)}.$$

---

126. $\operatorname{cosec}A\operatorname{cosec}2A+\operatorname{cosec}2A\operatorname{cosec}3A=\operatorname{cosec}A(\cot A-\cot 3A)$.

127. Solve the equation

$$\cos 3\theta+\cos 5\theta+\sqrt{2}(\cos\theta+\sin\theta)\cos\theta=0.$$

128. In any triangle

$$4\sin A\sin B\sin^2\tfrac{1}{2}C=(\sin B+\sin C-\sin A)(\sin C+\sin A-\sin B).$$

129. Find (without tables) to 3 places of decimals the numbers of which $1\cdot5$, $\cdot\dot{3}$ and $1\cdot\dot{3}$ are the common logarithms.

130. Two chords diverging from the same point on the circumference of a circle are to each other as the sines of the angles they respectively make with the tangent at that point.

---

131. Prove that there are eleven, and only eleven, pairs of regular polygons which are such that the number of degrees in an angle of one of them = the number of grades in an angle of the other; and that there are only four pairs when the number is an integer.

132. $\cos 11A+3\cos 9A+3\cos 7A+\cos 5A$
$$=16\cos^3 A\cos(4A+\tfrac{1}{4}\pi)\cos(4A-\tfrac{1}{4}\pi).$$

133. If $\sin^{-1}x+\sin^{-1}\frac{1}{2}x=\frac{1}{4}\pi$, then $x=\sqrt{\{\frac{2}{17}(5-2\sqrt{2})\}}$.

134. Transform $\sqrt{(2\sec A)}=\frac{1}{6}\sqrt[3]{(\cos^2 B\operatorname{cosec}C)}$ into an equation between tabular logarithms.

135. The distances of A one of the angular points of a regular octagon $ABCDEFGH$ from the sides BC, CD, DE respectively are as

$$1:1+\sqrt{2}:2+\sqrt{2}.$$

---

136. $\sin(2\alpha+\beta+\gamma)\sin(\beta-\gamma)+\sin(\alpha+2\beta+\gamma)\sin(\gamma-\alpha)$
$$+\sin(\alpha+\beta+2\gamma)\sin(\alpha-\beta)=0.$$

137. If $\log 10\frac{2}{3}\frac{5}{4}=\alpha$ and $\log 2=\beta$, then $\log 4100=\alpha+12\beta$.

138. $\sin 81^0-\sin 39^0-\sin 21^0+\sin 99^0=\sin 90^s$.

139. Solve $\sin(n+1)\theta+\sin(n-1)\theta=\sin 2\theta$.

140. $ABCD$ is a quadrilateral and AB, DC produced meet in E; the angle $DAC=DBC=\alpha$; the angle $CAB=\beta$; the angle $CBE=\gamma$; prove that $$BE=AB\,\frac{\sin\beta\sin(\alpha+\beta)}{\sin(\gamma-\beta)\sin(\alpha+\beta+\gamma)}.$$

---

141. $\cos 2A+\sin 2B=2\sin\{\frac{1}{4}\pi-(A-B)\}\cos\{\frac{1}{4}\pi-(A+B)\}$
$\cos 2A-\sin 2B=2\sin\{\frac{1}{4}\pi-(A+B)\}\cos\{\frac{1}{4}\pi-(A-B)\}$.

142. (i) $\cos 55^0+\cos 65^0+\cos 175^0=0$,
(ii) $\sin^2 24^0-\sin^2 6^0=\frac{1}{8}(\sqrt{5}-1)$.

143. If $A+B+C+D=180^0$, then
$\cos 2A-\cos 2B+\cos 2C-\cos 2D=4\sin(B+C)\cos(C+A)\sin(A+B)$.

144. Given $\log 35=a$, $\log 325=b$, $\log 245=c$,
find $\log 5$, $\log 7$ and $\log 13$.

145. A train is going due East at the rate of 24 miles an hour; when will it be 18 miles distant from a town which is on the N.E. of the train at a distance of 24 miles?

---

146. If $4\sin A\tan(A-B)+3\sec A=4\sin^2 A\sec A$,
then $\tan A\tan B=3$.

147. If $\tan(\pi\cot\theta)=\cot(\pi\tan\theta)$,
then $4\tan\theta=2n+1\pm\sqrt{(4n^2+4n-15)}$.

148. If $x=\cos 2\alpha+\cos\alpha$, $y=\sin 2\alpha+\sin\alpha$,
then $2x=(x^2+y^2)^2-3(x^2+y^2)$.

149. In any triangle $2R\sin C=b\cos A+\sqrt{(a^2-b^2\sin^2 A)}$.

150. In a triangle right-angled at C
$$\tan^{-1}\frac{a}{b+c}+\tan^{-1}\frac{b}{a+c}=\frac{\pi}{4}.$$

---

151. One angle of a quadrilateral is 60^0, another 50^g, another $\frac{2}{5}\pi$; express all four angles in degrees.

152. If $\sin^{-1}m+\sin^{-1}n=\frac{1}{2}\pi$, prove that $\sin^{-1}m=\cos^{-1}n$.

153. If $\cos\alpha=\dfrac{\cos\beta-e}{1-e\cos\beta}$, then $\tan\frac{1}{2}\alpha=\sqrt{\left(\dfrac{1+e}{1-e}\right)}\tan\frac{1}{2}\beta$.

154. $(\cos\theta+\sin\theta)(\cos 2\theta+\sin 2\theta)=\cos\theta+\cos(3\theta-\frac{1}{2}\pi)$.

155. In any triangle $a^2b^2c^2-4b^2S^2$; $a^2b^2c^2\cos A-4bcS^2$ and $a^2b^2c^2-4c^2S^2$ are in G. P.

---

156. Simplify $\cos^2(A+B)+\cos^2(A-B)-\cos 2A\cos 2B$.

157. With two units differing by 10^0 the measure of an angle is as 3 is to 4; find the units.

158. $\cos 7^0 30' = \frac{1}{4}(-1+\sqrt{2}+\sqrt{3})\sqrt{(2+\sqrt{2})}$.

159. In any triangle

$$\frac{\sin\frac{1}{2}A+\cos\frac{1}{2}B-\sin\frac{1}{2}C}{\sin\frac{1}{2}A+\cos\frac{1}{2}C-\sin\frac{1}{2}B}=\frac{1+\tan\frac{1}{2}B}{1+\tan\frac{1}{2}C}.$$

160. In any triangle a^2, b^2-c^2, $b^2+c^2-2bc\cos(B-C)$, are in G. P.

---

161. Solve $\cos 3\theta+\sin 3\theta=\cos\theta+\sin\theta$.

162. If $x=3\cos\phi+\cos 3\phi$, $y=3\sin\phi-\sin 3\phi$, then

$$x^{\frac{2}{3}}+y^{\frac{2}{3}}=4^{\frac{2}{3}}.$$

163. $\cos^2 18^0\sin^2 36^0-\cos 36^0\sin 18^0=\frac{1}{16}$.

164. Given that $\cos\theta=-\frac{1}{2}$ is one solution of the equation $\cos\theta+\cos 3\theta=\frac{1}{2}$, find the others.

165. If the diagonal BD of a quadrilateral inscribed in a circle passes through the centre its area $=(s-a)(s-d)=(s-b)(s-c)$.

---

166. If $\tan(A+B)=3\tan A$,

then $\sin(2A+2B)+\sin 2A=2\sin 2B$.

167. $\sin 18^0+\cos 18^0=\sqrt{2}\cos 27^0$.

168. Solve $\tan 3\theta+\tan 2\theta+\tan\theta=0$.

169. If $A+B+C=90^0$, then $\cos^3 2A+\cos^3 2B+\cos^3 2C$
$=3\sin A\sin B\sin C-\sin 3A\sin 3B\sin 3C+1$.

170. In any triangle $bc\cos^2\frac{1}{2}A+ca\cos^2\frac{1}{2}B+ab\cos^2\frac{1}{2}C=(s)^2$.

---

171. Prove that $7\theta=\pi$ satisfies the equation

$$8\cos\theta\cos 2\theta\cos 3\theta=1.$$

172. If $A+B+C=90^0$, then $\cos A\cos B\cos C$ cannot be greater than $\frac{8}{9}\sqrt{3}$.

173. $\sin^{-1}\frac{1}{5}\sqrt{5}+\cot^{-1}3=\frac{1}{4}\pi$.

174. In any triangle

$$\sin 3A+\sin 3B+\sin 3C+4\cos\tfrac{3}{2}A\cos\tfrac{3}{2}B\cos\tfrac{3}{2}C=0.$$

175. If the circle circumscribing an isosceles triangle is equal to the escribed circle touching one of the equal sides the triangle is right angled.

---

176. If $\sin x + \sin^2 x = 1$, then $\cos^2 x + \cos^4 x = 1$.

177. $\sin 10^0 \sin 50^0 \sin 70^0 = \frac{1}{8}$.

178. Simplify $\tan^{-1}\left(\frac{x\cos\theta}{1 - x\sin\theta}\right) - \cot^{-1}\left(\frac{\cos\theta}{x - \sin\theta}\right)$.

179. A, B, C, D are the angles of a quadrilateral; prove that the sum of the products of the sines of $\frac{1}{2}A$, $\frac{1}{2}B$, $\frac{1}{2}C$, $\frac{1}{2}D$ taken two together is equal to the sum of the products of their cosines taken two together.

180. At each end of a horizontal line of length $2a$ the altitude of a certain tower is α and at the middle point of the line it is β; show that the vertical height of the tower above the horizontal plane is

$$a \sin\alpha \sin\beta \sqrt{\{\operatorname{cosec}(\beta + \alpha) \operatorname{cosec}(\beta - \alpha)\}}.$$

---

181. Prove that

$$\sin(\alpha - \beta)\cos 2\beta + \cos(\alpha - \beta)\sin 2\beta = \sin(\beta - \alpha)\cos 2\alpha + \cos(\beta - \alpha)\sin 2\alpha.$$

182. Find the values of $\sin 2\theta \cos \frac{1}{2}\theta$ and $\tan\theta$ when $\theta = \sin^{-1}\frac{5}{13}$.

183. Prove geometrically that $\sin 2\theta$ and $2\sin\theta\cos\theta$ are always of the same sign.

184. If D is the middle point of the side BC, then

$$4AD^2 = b^2 + c^2 + 2bc\cos A.$$

185. If l, m, n are the distances of the centre of the inscribed circle from ABC, then

$$\frac{lmn}{r} = \frac{2abc}{a+b+c}.$$

---

186. If $a\sin\theta = b\sin\phi$, $c = a\cos\theta + b\cos\phi$, then

$$\tan(\theta + \phi)\,(c^2 - a^2 - b^2) = \sqrt{\{(a+b+c)\,(-a+b+c)\,(a-b+c)\,(a+b-c)\}}.$$

187. If $\sin^{-1}\alpha + \sin^{-1}\beta + \sin^{-1}\gamma = \pi$, then

$$\alpha\sqrt{(1-\alpha^2)} + \beta\sqrt{(1-\beta^2)} + \gamma\sqrt{(1-\gamma^2)} = 2\alpha\beta\gamma.$$

188.
$$\frac{\sin(\theta-\beta)\sin(\theta-\gamma)}{\sin(\alpha-\beta)\sin(\alpha-\gamma)} + \frac{\sin(\theta-\gamma)\sin(\theta-\alpha)}{\sin(\beta-\gamma)\sin(\beta-\alpha)} + \frac{\sin(\theta-\alpha)\sin(\theta-\beta)}{\sin(\gamma-\alpha)\sin(\gamma-\beta)} = 1.$$

189. If $A + B + C = 90$, then

$\sin^2 A + \sin^2 B + \sin^2 C$ cannot be less than $\frac{3}{4}$.

190. The perimeter of a regular circumscribing polygon of a circle is double that of the inscribed polygon having the same number of sides; what is that number?

---

191. Eliminate θ from the equations

$$m \sin 2\theta = n \sin \theta, \quad p \cos 2\theta = q \cos \theta.$$

192. Reduce $1 - \cos^2 \theta - \cos^2 \phi - \cos^2 \psi + 2 \cos \theta \cos \phi \cos \psi$ to the product of four cosines.

193. $$\tan 5A = \frac{5 \tan A - 10 \tan^3 A + \tan^5 A}{1 - 10 \tan^2 A + 5 \tan^4 A}.$$

194. Prove (i) $4 \sin 18^0 \cos 36^0 = 1$,

(ii) $\cos 36^0 - \sin 18^0 = \frac{1}{2}$.

195. $r = s \tan \frac{1}{2}A \tan \frac{1}{2}B \tan \frac{1}{2}C$.

---

196. Solve the equation

$$\cos \theta - \sin \theta = \cos \alpha - \sin \alpha.$$

197. Prove that when

$$\cos(x+2A) + \cos(x+2B) + \cos(x+2C) + \cos(x+2D) = 4 \cos(A+B) \cos(A+C) \cos(A+D),$$

then

$$\sin(x+2A) + \sin(x+2B) + \sin(x+2C) + \sin(x+2D) = 4 \sin(A+B) \sin(A+C) \sin(A+D),$$

where A, B, C, D are the angles of a quadrilateral and x is not zero.

198. Express in four factors

$$\sin^2 A + \sin^2 B + \sin^2 C - 2 \sin A \sin B \sin C - 1.$$

199. In any triangle

$$2R \{\cos^2 \tfrac{1}{2}A + \cos^2 \tfrac{1}{2}B + \cos^2 \tfrac{1}{2}C\} = 4R + r.$$

200. The diagonals of a four-sided figure are h and k, and the area is C; show that the area of its circumscribing square is

$$\frac{h^2 k^2 - 4C^2}{h^2 + k^2 - 4C}.$$

**MISCELLANEOUS EXAMPLES. LXXVI.

(1) If $2\cos\theta - \cos 2\theta = a$ and $2\sin\theta - \sin 2\theta = b$, prove that $(a^2+b^2-3)^2 = 12 - 8a$.

(2) If $h\cos\theta + k\sin\theta = 1$ and $l\cos\theta + m\sin\theta = 1$, prove that $(l-h)^2 + (m-k)^2 = (lh-mk)^2$.

(3) The diagonals of a rhombus are $2a$ and $2b$, prove that the cosines of its angles are $\pm\frac{a^2-b^2}{a^2+b^2}$.

(4) One of the values of

$$\sqrt{[2+\sqrt{\{2+\sqrt{(2+\ldots+\sqrt{2+2\cos A})}\}}]} \text{ is } 2\cos\frac{A}{2^n}.$$

(5) If $8x = \log_e 3$, prove that the angle whose tangent is $\frac{e^{2x}-e^{-2x}}{e^{2x}+e^{-2x}}$ is 15^0.

(6) If the number of degrees (A) in an angle is the same as the number of radians in another angle, and if the tangents of these angles are equal, prove that A is some multiple of $\frac{180\pi}{180-\pi}$.

(7) If α, β, γ are in A.P., then

$$\sin\alpha + \sin\gamma = 2\sin\beta\,.\cos(\beta-\alpha).$$

(8) Two parallel chords of a circle, lying on the same side of the centre, subtend respectively 72^0 and 144^0 at the centre. Prove that the distance between the chords is half the radius of the circle.

(9) Prove that $4\sin(\theta-\alpha)\,.\sin(m\theta-\alpha)\,.\cos(\theta-m\theta)$
$$= 1+\cos(2\theta-2m\theta) - \cos(2\theta-2\alpha) - \cos(2m\theta-2\alpha).$$

(10) Express $x^4+y^4+z^4-2y^2z^2-2z^2x^2-2x^2y^2$ in a form fitted for logarithmic calculation.

(11) Solve the equations

(i) $\sin 3\theta + \sqrt{3}\cos 3\theta = 1$. (ii) $\sin m\theta = \cos n\theta$.

(iii) $\dfrac{\cos(\beta+x)}{\cos(\alpha-x)} = \dfrac{m\sin\beta}{n\sin\alpha}$. (iv) $\tan m\theta = \cot n\theta$.

(v) $\tan\theta + \tan 2\theta + \tan 3\theta = 0$.

(vi) $\cos 8\theta - \cos 5\theta + \cos 3\theta = 1$.

(vii) $\cos\theta \,.\, \cos 3\theta = \cos 5\theta \,.\, \cos 7\theta$.

(12) In any triangle $\tan\dfrac{A-B}{2} = \tan\left(\phi - \dfrac{\pi}{4}\right)\cot\dfrac{C}{2}$, where $\phi = \tan^{-1}\dfrac{a}{b}$.

(13) Prove that

$$b^2 + c^2 - 2bc\cos(60^0 + A) = c^2 + a^2 - 2ca\cos(60^0 + B).$$

Hence prove that if equilateral triangles are described on the sides of the triangle ABC, the centres of circles inscribed in them will be the angular points of an equilateral triangle.

(14) A round tower stands on an island in a lake. A, B are two points on the land such that AB is a feet and points directly to the middle of the tower. At A and B the base of the tower subtends the angles 2α and 2β respectively. Prove that the diameter of the tower is $2\dfrac{a\sin\alpha\,.\,\sin\beta}{\sin\beta - \sin\alpha}$.

(15) At P the top of a tower of height h, the angles of depression of two objects on the horizontal plane on which the tower stands are $\frac{1}{4}\pi - \alpha$ and $\frac{1}{4}\pi + \alpha$. Prove that the angle $APB = 2\alpha$, and that $AB = 2h\tan 2\alpha$.

(16) On the side of a hill there are two places B, C inaccessible to each other, but known to be at the same distance (a) from a certain station A also on the hill. At the lower place C the *horizontal* angle θ between A and B is observed as well as the altitudes λ, μ of A and B. Prove that the distance between B and C is $2a\left\{\cos(\lambda-\mu)\cos^2\dfrac{\theta}{2} - \cos(\lambda+\mu)\sin^2\dfrac{\theta}{2}\right\}$.

(17) In the ambiguous case, given B, c, b, if a_1, a_2 are the two values of a, prove the following statements:

(i) $a_1 + a_2 = 2c \cos B$. (ii) $a_1 a_2 = c^2 - b^2$.

(iii) $a_1^2 - 2a_1 a_2 \cos 2B + a_2^2 = 4b^2 \cos^2 B$.

(iv) The distance between the centres of the circumscribing circles of the two triangles is $\dfrac{a_1 - a_2}{2 \sin B}$.

(v) The circumscribing circles of the two triangles are equal.

(vi) If $B = 45^0$, the angle between the two positions of b is

$$\cos^{-1} \frac{2a_1 a_2}{a_1^2 + a_2^2}.$$

(18) Prove that if I_1, I_2, I_3 are the centres of the escribed circles, then $I_2 A I_3$ is a straight line, and $I_1 A$, $I_2 B$, $I_3 C$ are perpendicular to $I_2 I_3$, $I_3 I_1$, $I_1 I_2$ respectively. Prove also that

(i) $I_2 I_3 = a \operatorname{cosec} \dfrac{A}{2}$. (ii) The angle $I_2 I_1 I_3 = \dfrac{\pi}{2} - \dfrac{A}{2}$.

(iii) The area of $I_1 I_2 I_3 = \dfrac{sa}{\sin A} = \dfrac{\frac{1}{2}S}{\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}}$.

(iv) The radius of the circle circumscribing $I_1 I_2 I_3 = 2R$.

(19) If D, E, F are the feet of the perpendiculars from A, B, C on the opposite sides, the triangle DEF is called the 'pedal triangle' of the triangle ABC*; prove that

(i) $EF = a \cos A$. (ii) The angle $EDF = \pi - 2A$.

(iii) $\dfrac{1}{r} = \dfrac{1}{AD} + \dfrac{1}{BE} + \dfrac{1}{CF}$. (iv) $\dfrac{AD^2}{BE \cdot CF} = \dfrac{bc}{a^2}$.

(v) The radius of the circle circumscribing $AEF = R \cos A$.

(vi) The radius of the circle circumscribing $DEF = \frac{1}{2}R$.

* The student should notice that ABC is the pedal triangle of $I_1 I_2 I_3$ in Question (18).

(20) Prove that in any triangle ABC

$$a^2=(b-c)^2\left\{1+\frac{4bc}{(b-c)^2}\sin^2\frac{A}{2}\right\}.$$

Hence prove that if $\tan\theta=\dfrac{2\sqrt{bc}}{b-c}\sin\dfrac{A}{2}$, then $a=(b-c)\sec\theta$.

(21) Prove that in any triangle ABC

$$a=(b+c)\cos\beta, \text{ where } \sin\beta=\frac{2\sqrt{bc}}{b+c}\cos\frac{A}{2}.$$

Use this method, or that of Example (20), to find a by the aid of the Tables, when $b=347$, $c=293$, $A=39^0\ 42'$. [See Art. 260.]

(22) Prove that in any triangle

$$\log a=\log(b-c)+L\cos\frac{A}{2}-L\cos\phi\ ;\ \text{where } \phi=\tan^{-1}\frac{b+c}{b-c}\tan\frac{A}{2}.$$

(23) If D, E, F are the middle points of BC, CA, AB, then

$$\sin BAD=\frac{b\sin A}{\sqrt{b^2+2bc\cos A+c^2}}.$$ [See Ex (11) (15) page 205.]

(24) If $x=II_1$, $y=II_2$, $z=II_3$, $d=2R$, prove that

$$xyz+d(x^2+y^2+z^2)=4d^3.$$

(25) The circle escribed to the side BC of the triangle ABC will be the inscribed circle of the triangle whose sides are

$$\frac{sa}{s-a},\ \frac{sb}{s-a},\ \frac{sc}{s-a}.$$

(26) If a point Q be taken within ABC such that the angles BQC, CQA, AQB are each 120^0, prove that

$$OA=\tfrac{1}{2}\cdot\frac{4\sqrt{2}\Delta+\sqrt{6}(b^2+c^2-a^2)}{\{12\sqrt{3}\Delta+3(a^2+b^2+c^2)\}^{\frac{1}{2}}}.$$

(27) If $\sin A$ and $\cos A$ are both given, prove that in general n different values and no more can be assigned to $\sin\dfrac{A}{n}$.

(28) If from a point on a circular arc (of radius r) which subtends an angle α at the centre, perpendiculars are drawn to its bounding radii, then the distance between the feet of the perpendiculars is $r \sin \alpha$.

(29) If two circles, radii a and b, touch externally, the angle between their common tangents $= 2 \tan^{-1} \frac{1}{2} \left(\sqrt{\frac{a}{b}} - \sqrt{\frac{b}{a}} \right)$.

(30) If the inscribed circle touches the sides in DEF, prove that $EF : FD : DE = \cos \frac{A}{2} : \cos \frac{B}{2} : \cos \frac{C}{2}$.

(31) If the bisectors of the angles A, B, C meet the opposite sides in D, E, F then (i) $AD = \frac{2bc}{b+c} \cos \frac{A}{2}$. (ii) $CD = \frac{a \sin B}{\sin B + \sin C}$.

(iii) The area of DEF

$$= S \frac{2abc}{(b+c)(c+a)(a+b)} = S \cdot \frac{2 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}}{\cos \frac{B-C}{2} \cdot \cos \frac{C-A}{2} \cdot \cos \frac{A-B}{2}}.$$

(32) If α, β, γ, δ are the angles of a quadrilateral inscribed in a circle, then

$$\cos(\alpha+\beta) \cdot \cos(\beta+\gamma) \cdot \cos(\gamma+\delta) \cdot \cos(\delta+\alpha) = (1 - \cos^2 \alpha - \cos^2 \beta)^2.$$

(33) If θ_1 and θ_2 be two values of θ found from the equation $a \cos \theta + b \sin \theta = c$, then

$$\frac{1}{a} \cdot \cos \frac{\theta_1 + \theta_2}{2} = \frac{1}{b} \cdot \sin \frac{\theta_1 + \theta_2}{2} = \frac{1}{c} \cdot \cos \frac{\theta_1 - \theta_2}{2}.$$

(34) If
$$\left.\begin{aligned} a &= b \cos \chi + c \cos \phi \\ b &= c \cos \theta + a \cos \chi \\ c &= a \cos \phi + b \cos \theta \end{aligned}\right\}$$

prove that $\cos 2\theta + \cos 2\phi + \cos 2\chi + 4 \cos \theta \cdot \cos \phi \cdot \cos \chi + 1 = 0$,

hence [see Ex. (20), p. 143], prove that

$$(\theta \pm \phi \pm \chi) = (2n+1)\pi.$$

(35) A circle is drawn touching the inscribed circle and the sides AB, AC (not produced) of the triangle ABC. Another circle is drawn touching the circle and the same side. Prove that the radius of the n^{th} circle thus drawn is $r\left(\frac{1-\sin\frac{A}{2}}{1+\sin\frac{A}{2}}\right)^n$ and that the sum of the radii of all the circles, when $n=\infty$, is

$$\frac{r}{2}\left(\operatorname{cosec}\frac{A}{2}-1\right).$$

(36) If AO, BO, CO pass through the centre of the circumscribing circle and meet the opposite sides on DEF, then

$$\frac{1}{AD}+\frac{1}{BE}+\frac{1}{CF}=\frac{2}{R}.$$

(37) If h and k be the lengths of the diagonals of a quadrilateral and θ the angle between them, prove that the area of the quadrilateral is $\frac{1}{2}hk\sin\theta$.

(38) If $\frac{x}{\sin X}=\frac{y}{\sin Y}=\frac{z}{\sin Z}$ and $X+Y+Z=180^0$, prove that $x=y\cos Z+z\cos Y$.

(39) Two regular polygons of n and $2n$ sides are described such that the circle inscribed in the first circumscribes the second; also the radius of the circle inscribed in the second is to that of the circle circumscribing the first as $3+\sqrt{3}$ is to $4\sqrt{2}$. Prove that $n=6$.

(40) Prove by the aid of a figure like that of Art. 282 that $I_1O^2=R^2+2Rr_1$.

(41) If $yz+zx+xy=1$, prove by Trigonometry that

$$\frac{x}{1-x^2}+\frac{y}{1-y^2}+\frac{z}{1-z^2}=\frac{4xyz}{(1-x^2)(1-y^2)(1-z^2)}.$$

[See Ex. (33), (32), p. 194.]

EXAMINATION PAPERS.

†I. PREVIOUS EXAMINATION.

December, 1886.

PAPER I.

I. DEFINE the secant and cotangent of an angle, and prove that

(i) $\sec\theta \operatorname{cosec}\theta = \tan\theta + \cot\theta$,

(ii) $\cot^2\theta \dfrac{\sec\theta - 1}{1+\sin\theta} + \sec^2\theta \dfrac{\sin\theta - 1}{1+\sec\theta} = 0.$

PAPER II.

1. From the definitions of the Trigonometrical Functions, prove that

$$\sin^2 A = 1 - \cos^2 A \quad \text{and} \quad \sin A \tan A = \sec A - \cos A.$$

Prove that $\frac{1}{8}(\cos^6 A + \sin^6 A) - \frac{1}{4}(\cos^2 A - \sin^2 A)^2 = \frac{1}{17}$.

2. Investigate the values of $\tan 45^0$ and $\sin 60^0$.

Two adjacent sides of a parallelogram are of lengths 15 and 24, and the angle between them is 60^0; find the lengths of both diagonals.

II. PREVIOUS EXAMINATION, CAMBRIDGE.

June, 1887.

PAPER I.

1. Define the sine and tangent of an angle, and shew how to find the sine and tangent whose cosine (m) is given.

If $\sin A = \tan B$, prove

$$\cos^2 A \cos^2 B = (\cos B + \sin B)(\cos B - \sin B).$$

† The Additional Subjects in the Cambridge Previous Examination, which are required of Candidates for any Tripos, are now (1887) *either* Mathematics, *or* French, *or* German.

The Mathematical subjects are (i) The Trigonometry of one Angle, (ii) Elementary Dynamics [see Lock's *Dynamics for Beginners*], (iii) Elementary Statics. The Trigonometry is set in the first one or two questions in each of the two papers. The questions quoted above were set in December, 1886, and in June, 1887.

2. Trace the changes in the tangent of an angle as the angle changes from 180^0 to 270^0.

If $\sin\theta = -\frac{3}{5}$, find $\tan\theta$; and explain by means of a figure the reason why there are two answers to the question.

PAPER II.

1. Explain the mode of measuring angles in degrees, minutes and seconds.

Find the number of seconds of angle through which the earth revolves about its axis in a second of time.

2. Find the value of $\sec 60^0$, and of $\sec 45^0$.

Solve the equation

$$2 \operatorname{cosec} A = 2 \sin A + \cot A.$$

III. CAMBRIDGE LOCAL EXAMINATIONS.

Dec. 1886.

JUNIOR CANDIDATES.

1. Prove that the angle subtended at the centre of a circle by an arc equal to the radius is the same for all circles.

Express the angle as a fraction of a right angle.

2. Define the sine, cosine and tangent of an angle. Prove that these trigonometrical ratios are always the same for the same angle.

Find these ratios for an angle of 45^0.

3. Prove that

$$\cos(A+B) = \cos A \cos B - \sin A \sin B.$$

Prove that the sum of the cosines of two angles is equal to twice the cosine of half their sum multiplied by the cosine of half their difference.

4. Prove the following relations:

(i) $\tan(A+B) = \dfrac{\tan A + \tan B}{1 - \tan A \tan B}$.

(ii) $(1 + \sin A + \cos A)^2 = 2(1 + \sin A)(1 + \cos A)$.

(iii) $\dfrac{\sin 3A + \sin 5A}{\cos 3A - \cos 5A} = \cot A$.

5. Prove that the logarithm of the product of two numbers is equal to the sum of the logarithms of the numbers.

Having given $\log 2 = \cdot 3010300$, $\log 7 = \cdot 8450980$, find the logarithms of $(1\cdot75)^{\frac{1}{2}}$, $(24\cdot5)^{-\frac{1}{3}}$.

6. Prove that in any triangle ABC

(i) $a = b\cos C + c\cos B$.

(ii) $2bc\cos A = b^2 + c^2 - a^2$.

7. Shew how to solve a triangle when one side and two angles are known.

Find the side b in the triangle ABC from the following data:

$$a = 156{\cdot}22,\quad B = 57^\circ 25',\quad C = 63^\circ 42',$$

$$\log 15{\cdot}622 = 1{\cdot}1937366,\quad L\sin 57^\circ 25' = 9{\cdot}9256261,$$

$$\log 15{\cdot}37552 = 1{\cdot}1868297,\quad L\sin 58^\circ 53' = 9{\cdot}9325330.$$

8. The angles of elevation of the top of a tower on a horizontal plane observed at two points distant a feet and b feet respectively from the base and in the same straight line with it are found to be complementary. Shew that the height of the tower is $\sqrt{ab}$ feet. If θ be the angle subtended at the top of the tower by the line joining the two points, prove that

$$\sin\theta = \frac{a \sim b}{a + b}.$$

IV. OXFORD LOCAL EXAMINATIONS.

July, 1887.

JUNIOR CANDIDATES.

1. What is π?

What is 'the angle whose circular measure is π'?

In a triangle ABC the angle A is x degrees, the angle B x grades, and the circular measure of C is $\frac{\pi x}{9}$; find the number of degrees in each of the angles.

2. Prove that

$$2\sin\frac{A}{2} = \pm\sqrt{1+\sin A} \pm \sqrt{1-\sin A},$$

and determine which are the correct signs when

$$270^\circ > A > 180^\circ.$$

3. Obtain the following formulæ:

(1) $\cos(A+B) = \cos A\cos B - \sin A\sin B$;

(2) $\tan A + \tan B = \sin(A+B)\sec A\sec B$;

(3) $\dfrac{\sin(p+q) - 2\sin p + \sin(p-q)}{\cos(p+q) - 2\cos p + \cos(p-q)} = \tan p$;

(4) $\sin 2\theta\sin 2\phi = \sin^2(\theta+\phi) - \sin^2(\theta-\phi)$.

4. Given $\log 2 = \cdot 3010300$,
$\log 3 = \cdot 4771213$, $\log 24668 = 4 \cdot 3921339$,
$\log 11 = 1 \cdot 0413927$, $\log 24669 = 4 \cdot 3921515$:
find the logarithm of $30 \cdot 25$ and calculate the value of

$$\{165 \times (30)^9 \times \sqrt[3]{24}\} \div (121)^7.$$

5. In a triangle ABC, a, b, c are the sides, s is the semi-perimeter, Δ the area, R, r the radii of the circumscribed and inscribed circles: prove that

(1) $\tan \frac{B-C}{2} = \frac{b-c}{b+c} \cot \frac{A}{2}$;

(2) $\sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}}$;

(3) $2R = a \operatorname{cosec} A = b \operatorname{cosec} B = c \operatorname{cosec} C$;

(4) $2\Delta = ca \sin B$.

6. In a plane triangle the sides a, b and the angle A are known; shew that in general two values of c can be found, and that the difference of these values is

$$2\sqrt{a^2 - b^2 \sin^2 A}.$$

7. A ladder placed at an angle of 75^0 just reaches the sill of a window 27 feet above the ground on one side of a street. On turning the ladder over without moving its foot, it is found that, when it rests against a wall on the other side of the street, it is at an angle of 15^0. Find the breadth of the street.

8. If $B = 36^0\,46'$, $b = 311 \cdot 8785$, $c = 521 \cdot 05$, find C.
[$L \sin B = 9 \cdot 7771060$, $\log 31187 = 4 \cdot 4939736$,
$\log 521 \cdot 05 = 2 \cdot 7168794$, $\log 31188 = 4 \cdot 4939875$.]

V. CAMBRIDGE LOCAL EXAMINATIONS.

December, 1886.

SENIOR STUDENTS.

1. Explain the method of measuring angles in circular measure, and find the circular measure of the angle of a regular pentagon.
Prove that

$$\frac{\tan^3 A}{1+\tan^2 A} + \frac{\cot^3 A}{1+\cot^2 A} = \frac{1 - 2\sin^2 A \cos^2 A}{\sin A \cos A}.$$

2. Establish the identities:

(1) $\tan^4 A = \frac{3 - 4\cos 2A + \cos 4A}{3 + 4\cos 2A + \cos 4A}$.

(2) $\sin A - \sin B = 2 \sin \frac{A-B}{2} \cos \frac{A+B}{2}$.

If $A+B+C=180^0$, shew that

$$(\sin B-\sin C)\cot\frac{A}{2}+(\sin C-\sin A)\cot\frac{B}{2}+(\sin A-\sin B)\cot\frac{C}{2}=0.$$

3. Find an expression for all the angles which have a given sine.

Solve the equations:

(1) $\cos\theta+\tan\theta=\sec\theta$.

(2) $\sin\theta-2\sin 2\theta\cos\theta+\cos 3\theta=\cos 2\theta$.

4. Prove that in any triangle if a, b, c be the sides opposite to the angles A, B, C

(1) $$\frac{\sin A}{a}=\frac{\sin B}{b}=\frac{\sin C}{c},$$

(2) $$\cos A+\cos B+\cos C=1+\frac{r}{R},$$

where r and R are the radii of the inscribed and circumscribed circles respectively.

If the line joining A to the centre of the inscribed circle meets the opposite side in D, prove that

$$\tan ADB=\frac{b+c}{b-c}\tan\frac{A}{2}.$$

*5. If θ is the circular measure of a positive angle less than a right angle, shew that $\sin\theta$ lies between θ and $\theta-\frac{\theta^3}{6}$.

Find the limiting value of $\left(\frac{n}{\theta}\sin\frac{\theta}{n}\right)^n$ when n is indefinitely increased.

*6. Prove that, if θ lie between $\frac{\pi}{4}$ and $-\frac{\pi}{4}$,

$$\theta=\tan\theta-\tfrac{1}{3}\tan^3\theta+\tfrac{1}{5}\tan^5\theta-\ldots\ldots.$$

Shew that

$$\log(a+b\sqrt{-1})=\tfrac{1}{2}\log(a^2+b^2)+\sqrt{-1}\tan^{-1}\frac{b}{a}.$$

* See *Higher Trigonometry*.

VI. OXFORD LOCAL EXAMINATIONS.

July, 1887.

SENIOR CANDIDATES.

1. Prove geometrically that the sine of 90^0+A is equal to the cosine of A for all values of A.

Find all the values of B, less than 180^0, for which $\sin 5B=\frac{1}{2}\sqrt{2}$.

2. (1) Prove that $2\sin A = \pm\sqrt{1+\sin 2A} \pm \sqrt{1-\sin 2A}$; and determine the signs of the roots when $2A$ is greater than 180^0 and less than 270^0.

(2) Verify the identities:

(α) $\sin 4A = 4\cos A(\sin A - 2\sin^3 A)$;

(β) $(\tan 4A + \tan 2A)(1 - \tan^2 3A \tan^2 A) = 2\tan 3A \sec^2 A$.

3. Prove the following properties of a triangle ABC, of which the sides are a, b, c, and the semi-perimeter is s:

(1) $\text{area} = \sqrt{s(s-a)(s-b)(s-c)}$;

(2) $$\frac{bc\cos A + ca\cos B + ab\cos C}{bc+ca+ab} = \frac{a\sin A + b\sin B + c\sin C}{(b+c)\sin A + (c+a)\sin B + (a+b)\sin C}.$$

4. ABC is an isosceles triangle, having the vertical angle $A = 56^0\,30'$; on AB as base an isosceles triangle DAB is described, having its vertical angle $D = 38^0\,40'$: the perpendicular from A upon $BC = 22{\cdot}75$ inches: find the length of the perpendicular from D upon AB to four places of decimals.

$L\cot 19^0\,20' = 10{\cdot}4548807$; $\quad L\cos 28^0\,15' = 9{\cdot}9449220$;

$\log 11375 = 4{\cdot}0553514$;

$\log 36754 = 4{\cdot}5653046$, diff. for $1 = 119$.

5. (1) If in a triangle the radius of the circumscribed circle is double of the radius of the inscribed circle, the triangle is equilateral.

(2) Compare the areas of two equilateral and equiangular polygons, each of n sides, one inscribed in and the other circumscribed about the same circle.

VII. WOOLWICH.

December, 1886.

[N.B.—*Great importance will be attached to accuracy.*]

1. Explain what is meant by the *circular measure* of an angle, and shew that the circular measure of an acute angle is intermediate in value between the sine and the tangent of the angle.

The perimeter of a certain sector of a circle is equal to the length of the arc of a semicircle having the same radius. Express the angle of the sector in degrees, minutes, and seconds.

2. Prove that the secant of any angle will be either greater than $+1$ or less than -1.

Shew that $\sec A = \pm\sqrt{1+\tan^2 A}$, and explain the appearance of the double sign.

3. Express $\sin(A-B)$ and $\cos(A-B)$ in terms of the sines and cosines of A and B.

Find the values of $\sin 15^0$, $\sin 105^0$, $\cos 165^0$, and $\tan 195^0$.

4. Obtain a formula including all angles which have a given tangent.

If $\tan A + \tan 2A = \tan 3A$, prove that A must be a multiple of 60^0 or 90^0.

5. A ring, 10 inches in diameter, is suspended from a point 1 foot above its centre by six equal strings attached to its circumference at equal intervals. Find the cosine of the angle between two consecutive strings.

6. Prove that

(1) $\sin 2A = \dfrac{2\tan A}{1+\tan^2 A}$.

(2) $\cos^2 A - \cos A \cos(60^0 + A) + \sin^2(30 - A) = \frac{3}{4}$.

7. State and prove the rule for finding, by inspection, the characteristic of the logarithm of a fraction.

Find from the tables supplied an approximate value of the seventh root of ·000026751.

8. Express the cosine of any angle of a triangle in terms of the three sides.

Prove that

$$\sin(B - C) : \sin(B + C) :: b^2 - c^2 : a^2.$$

9. The sides of a triangle are 17, 20, and 27. Find from the tables supplied all the angles.

10. A man travelling due west along a straight road observes that when he is due south of a certain windmill the straight line drawn to a distant church makes an angle of 30^0 with the direction of the road. A mile further on the bearings of the windmill and tower are N.E. and N.W. respectively. Find the distances of the tower from the windmill, and from the nearest point of the road.

11. Find the area of a regular polygon of n sides inscribed in a circle of given radius.

If a regular pentagon and a regular decagon have the same perimeter, prove that their areas are as $2 : \sqrt{5}$.

VIII. SANDHURST FURTHER.

December, 1885.

1. Define the two common units of angular measure.

Find the circular measure of 42^0, and find the angle whose circular measure is $\frac{5}{8}$.

2. Assuming that

$$\left.\begin{aligned}\sin(A+B) &= \sin A \cos B + \cos A \sin B\\ \cos(A+B) &= \cos A \cos B - \sin A \sin B\end{aligned}\right\},$$

find in terms of the ratios of A, the values of

$$\sin 2A, \quad \cos 2A, \quad \tan 2A, \quad \sin \tfrac{1}{2}A, \quad \cos \tfrac{1}{2}A \text{ and } \tan \tfrac{1}{2}A.$$

3. Prove that

$$1-\tan^2 A\tan^2 B=\frac{\cos^2 B-\sin^2 A}{\cos^2 A\cos^2 B};$$

and from the equation $2\tan^2\theta=\sec^2\theta$ find a general expression for θ.

4. Prove that in any triangle

$$\cos A+\cos B=\frac{2(a+b)}{c}\sin^2\tfrac{1}{2}C.$$

A lighthouse appears to a man in a boat 300 yards from its foot, to subtend an angle of $6^\circ 20' 27{\cdot}7''$; find in feet the height of the lighthouse, having given

$L\tan 6^\circ 20' = 9{\cdot}0452836$, difference for $1 = {\cdot}0011507$; $\log 3 = {\cdot}4771213$.

5. Prove that in any triangle

$$\tan 2A+\tan 2B+\tan 2C=\tan 2A\tan 2B\tan 2C.$$

Hence shew that if x, y, z are three numbers such that

$$x+y+z=xyz,$$

then $x(1-y^2)(1-z^2)+y(1-z^2)(1-x^2)+z(1-x^2)(1-y)=4xyz$.

IX. SANDHURST FURTHER.

June, 1886.

1. Express the value of the tangent, secant, and cosecant in terms of the sine of the angle, and also in terms of the cosine of twice the angle.

Find the values of the tangent, secant, and cosecant of $22^\circ 30'$.

2. Prove the following:

$$(\sin 2A)^2=2\cos^2 A(1-\cos 2A),$$

$$\tan 6A=\frac{\cos 5A-\cos 7A}{\sin 7A-\sin 5A};$$

and if

$$\frac{a}{b}=\frac{\cos A}{\cos B},$$

prove that $a\tan A+b\tan B=(a+b)\tan\frac{1}{2}(A+B)$.

3. Shew that in any triangle

$$(b+c-a)\tan\tfrac{1}{2}A=(c+a-b)\tan\tfrac{1}{2}B=(a+b-c)\tan\tfrac{1}{2}C$$

$$=\left\{\frac{(b+c-a)(c+a-b)(a+b-c)}{a+b+c}\right\}^{\frac{1}{2}}.$$

If R, r_a, r_b, r_c are the radii respectively of the circumscribed and three escribed circles of a triangle, shew that

$$Rr_a(s-a)=Rr_b(s-b)=Rr_c(s-c)=\tfrac{1}{4}abc,$$

where S is the semi-perimeter of the triangle.

4. Find the area of a triangle in terms of (i) two sides and the angle between them, (ii) two angles and the side between them.

5. Define the characteristic and the mantissa of a logarithm. Find the logarithm of 5 when the base is 3, and the logarithm of $\frac{1}{3}$ when the base is 5;

$$\log_{10} 3 = \cdot 4771213, \quad \log_{10} 2 = \cdot 3010300.$$

6. Two sides of a triangle are 9 and 7, and the included angle is $38^0\ 56'\ 32{\cdot}8''$; find the base and remaining sides

$$L\tan 19^0 29' = 9{\cdot}5487471, \quad L\tan 19^0\ 28' = 9{\cdot}5483452.$$

X. COLLEGE OF PRECEPTORS, PUPILS' EXAMINATION.

July, 1886.

1. In a quadrilateral $ABCD$, the angle $A = 30^0$, $B = 60$ grades, $C = \frac{2}{3}\pi$; find the number of degrees in D. Find what number of degrees must be taken from the angle D and added to A so that the figure may be inscribed in a circle.

2. If $\sec A = m$, what will $\cot A$ be? Shew from your result that when $A = 0$ or 180^0 the cotangent will be infinite in both cases.

3. Prove $\sin(A+B) = \sin A \cos B + \cos A \sin B$.

Shew that if $A + B + C = 180^0$,

$$\sin A + \sin B + \sin C = 4 \cos \tfrac{1}{2}A \cos \tfrac{1}{2}B \cos \tfrac{1}{2}C;$$

and that this is true numerically if A, B, C the angle of a triangle be 30, 60, 90 degrees respectively.

4. Solve $\cot^2 A + \tan^2 A = \frac{10}{3}$; giving the general value of A.

5. If $\log_a 243 = 5$, find a.

Given $\log_{10} 30 = 1{\cdot}4771213$, $\log_{10} 20 = 1{\cdot}3010300$, find $\log_{10} 72$ and $\log_{10} 1{\cdot}44$.

6. In any triangle, prove the formula

$$\tan \frac{A}{2} = \pm \sqrt{\left\{ \frac{(s-b)(s-c)}{s(s-a)} \right\}},$$

and shew from it that if $a = 5$, $b = 3$, $c = 4$, then A will be a right angle. Can you give any explanation of the double sign?

7. Find the area of a regular polygon of n sides, each side being a.

8. Two angles of a triangle A and B, and their included side C being given, find an expression for the area; shew from the result that if A and B each remain constant while the side C varies, Euclid's statement is true; 'that the areas of similar rectilineal figures vary in the duplicate ratio of their homologous sides.'

9. Standing at a certain point, I observe the elevation of a house to be $15^0\ 15'$, and that the sill of one of its windows, known to be 20 feet above the ground, subtends an angle of 20^0 at the same point; shew that the height of the house is nearly 22 yards;

$$\log_{10} 200 = 2{\cdot}3010300, \quad L\tan 50^0\ 15' = 10{\cdot}0800379,$$
$$L\tan 20^0 = 9{\cdot}5610659, \quad \log_{10} 6{\cdot}606 = 0{\cdot}8200020.$$

XI. OXFORD AND CAMBRIDGE SCHOOL EXAMINATIONS, 1884.

1. Define a degree, and the unit of circular measure; and find the ratio of the first to the second.

In a triangle ABC, the circular measure of one angle (A) is a, and the number of degrees in another is B. If the ratio of the number of degrees in the third angle (C) to the circular measure of B be $B : a$, prove that A, B, C are in continued proportion.

2. Define the secant of an angle, and trace its changes in sign and magnitude as the angle increases from 90^0 to 270^0.

Simplify

$$\left\{\frac{1}{\cos\theta + \tan^2\theta \sin\theta} - \frac{1}{\cos\theta \cot^2\theta + \sin\theta}\right\} \times \left\{\frac{\sec\theta \operatorname{cosec}\theta - 1}{\operatorname{cosec}\theta - \sec\theta}\right\}.$$

3. Prove that if two angles have the same tangent, their difference is a multiple of two right angles.

Solve the equation

$$\sec\theta\,(\sec^2\theta + 2)\,(\operatorname{cosec}\theta - \sin\theta) = 4.$$

4. Prove geometrically that when $A + B$ is less than 90^0,

$$\cos(A+B) = \cos A \cos B - \sin A \sin B.$$

Prove that

$$\sin^{-1}\frac{1}{3} + \sin^{-1}\frac{1}{3\sqrt{11}} + \sin^{-1}\frac{3}{\sqrt{11}} = \frac{\pi}{2}.$$

5. A hill is inclined at a angle 36^0 to the horizon. An observer walks 100 yards away from the foot of the hill, and then finds that the elevation of a point halfway up the hill is 18^0. Find the height of the hill; and find the ratio of an error in measuring the distance walked to the consequent error in the height of the hill.

6. In any triangle prove that

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} = \frac{1}{2R},$$

where R is the radius of the circle described round ABC.

Find b, c, R in the triangle for which $B = 45^0$, $C = 60^0$, $a = \sqrt{2}$.

7. Find the area of a regular polygon of n sides; and deduce the area of a circle.

8. Find the cosines of the half angles of a triangle in terms of the sides.

If the bisectors of the angles A, B, C of a triangle meet the sides in D, E, F respectively, prove that

$$\frac{(s-b)\,DF^2}{(c+a)^2} - \frac{(s-c)\,DE^2}{(a+b)^2} = (c-b)\left\{\frac{abc}{(b+c)(c+a)(a+b)}\right\}^2.$$

Questions 9, 10 *in Higher Trigonometry.*

XII. OXFORD AND CAMBRIDGE SCHOOL EXAMINATIONS, 1885.

1. What is the unit of circular measure? Why is this unit used?

The angles of a triangle are in A.P.; prove that the mean angle is 60^0. If the number of right angles in the greatest is equal to the circular measure of the least, express the angles in degrees

$$\left(\pi = \frac{22}{7}\right).$$

2. Define the tangent of an angle less than 90^0, and extend the definition to angles unlimited in magnitude.

Find $\tan 30^0$, $\tan 120^0$, $\tan 240^0$.

3. Find the general solution of the equation $\sin\theta = \sin\alpha$.

Solve the equations:

(i) $2\cos^2\theta + \sqrt{2}\sin\theta = 2$.

(ii) $(1+\tan\theta)(1-\sin 2\theta) = 1-\tan\theta$.

4. Prove geometrically:

(i) $\sin(A-B) = \sin A\cos B - \cos A\sin B$.

(ii) $\dfrac{\cos(A+B)}{\sin(A-B)} = \dfrac{1-\tan A\tan B}{\tan A-\tan B}$.

Find the relation between α, β and γ, in order that

$$\cot\alpha\cot\beta\cot\gamma - \cot\alpha - \cot\beta - \cot\gamma \text{ should vanish.}$$

5. Prove the following statements:

(i) $\dfrac{1+\sin 2x+\cos 2x}{1+\sin 2x-\cos 2x} = \cot x$.

(ii) $3\cos^2\alpha = 4\cos^6\dfrac{\alpha}{2} + 4\sin^6\dfrac{\alpha}{2} - 1$.

(iii) $\cos 3\alpha\cos 2\alpha + \sin 4\alpha\sin\alpha = \cos\alpha\cos 2\alpha$.

6. Prove that in any triangle

$$\cos\left(\frac{A}{2}\right) = \sqrt{\left(\frac{s(s-a)}{bc}\right)}.$$

Having given that

$a = 12270$ ft., $b = 11550$ ft., $c = 11433$ ft.,

$\log 1{\cdot}76265 = {\cdot}2461661$, $\log 5{\cdot}3565 = {\cdot}7288811$,

$\log 1{\cdot}155 = {\cdot}0625820$, $\log 1{\cdot}1433 = {\cdot}0581602$,

$L \cos 32^\circ 15' = 9{\cdot}9272306$ diff. for $1' = {\cdot}0000797$;

find the angle A.

7. Find an expression for the radius of one of the described circles of a triangle in terms of the sides.

Prove that

$$\frac{bc}{r_1} + \frac{ca}{r_2} + \frac{ab}{r_3} = 2R\left\{\frac{b}{a} + \frac{c}{a} + \frac{c}{b} + \frac{a}{b} + \frac{a}{c} + \frac{b}{c} - 3\right\}.$$

8. Prove that if θ be the circular measure of an angle, the limiting value of $\dfrac{\sin\theta}{\theta}$ as θ is diminished is unity.

What is the limiting value of $\dfrac{\sin n'}{n}$ when n is diminished?

Questions 9, 10, 11 *in Higher Trigonometry.*

XIII. TRINITY COLLEGE. *June* 6, 1885. 9—12.

1. Define the sine and tangent of an angle.

The base BC of a triangle ABC is trisected in Q and R. Prove that

$$\sin BAR \sin CAQ = 4 \sin BAQ \sin CAR.$$

2. Give a geometrical proof that

(i) $\cos(\theta+\phi) = \cos\theta\cos\phi - \sin\theta\sin\phi$,

(ii) $\cos\phi - \cos\theta = 2\sin\frac{1}{2}(\theta+\phi)\sin\frac{1}{2}(\theta-\phi)$,

where θ and ϕ are each less than a right angle.

Show that if

$$\cos(\theta+\phi) = \frac{\sin^2\theta\cos\phi - \cos^2\theta\sin\phi}{\cos\theta\tan\alpha} = \frac{\sin^2\phi\cos\theta - \cos^2\phi\sin\theta}{\cos\phi\tan\beta},$$

then will

$$\cos(\alpha+\beta) = \frac{\sin^2\alpha\cos\beta - \cos^2\alpha\sin\beta}{\cos\alpha\tan\theta} = \frac{\sin^2\beta\cos\alpha - \cos^2\beta\sin\alpha}{\cos\beta\tan\phi}.$$

3. Find by mathematical induction the value of

$$\tan(\theta_1+\theta_2+\ldots+\theta_n) \text{ in terms of } \tan\theta_1,\ \tan\theta_2,\ldots.$$

Show that in a triangle ABC

$$\tan\tfrac{3}{4}A+\tan\tfrac{3}{4}B+\tan\tfrac{3}{4}C=\cot\tfrac{3}{4}A+\cot\tfrac{3}{4}B+\cot\tfrac{3}{4}C,$$

provided

$$\tan\tfrac{3}{4}A\tan\tfrac{3}{4}B=\cot\tfrac{3}{4}C.$$

4. From the results

$$a\operatorname{cosec}A=b\operatorname{cosec}B=c\operatorname{cosec}A, \text{ and } A+B+C=\pi,$$

deduce the formulæ

(i) $a^2=b^2+c^2-2bc\cos A$;

(ii) $a=b\cos C+c\cos B$.

In the base BC (produced if necessary) of an isosceles triangle ABC a point D is taken such that the sum of AD and AB is n times CD. Show that their difference is $1/n$th of BD.

5. The base of a triangle is equal to its altitude, and the two other sides are known. Determine the remaining parts of the triangle by formulæ adapted to logarithms.

Show that if r be the ratio of the two given sides, r must lie between

$$\tfrac{1}{2}(\sqrt{5}-1) \text{ and } \tfrac{1}{2}(\sqrt{5}+1).$$

6. Find the radius of the circle inscribed in a triangle in terms of one side and two angles.

Circles are inscribed in the triangles $D_1E_1F_1$, $D_2E_2F_2$, $D_3E_3F_3$, where D_1, E_1, F_1 are the points of contact of the circles escribed to the side BC, &c. Show that if r_1, r_2, r_3 be the radii of these circles

$$\frac{1}{r_1}:\frac{1}{r_2}:\frac{1}{r_3}=1-\tan\tfrac{1}{4}A:1-\tan\tfrac{1}{4}B:1-\tan\tfrac{1}{4}C.$$

7. Prove that the ratio of the radii r and R of the circles inscribed and circumscribed about a triangle ABC is

$$4\sin\tfrac{1}{2}A\sin\tfrac{1}{2}B\sin\tfrac{1}{2}C:1.$$

Two circles of radii ρ_1 and ρ_2 are drawn through the centre of the circumscribing circle to touch the sides AB and AC. Show that

$$\frac{1}{\rho_1}+\frac{1}{\rho_2}-\frac{r}{\rho_1\rho_2}=\frac{r+2R}{R^2}.$$

8. Show that

$$\tan^{-1}x-\tan^{-1}y=\tan^{-1}\frac{1}{y}-\tan^{-1}\frac{1}{x}.$$

Eliminate a between

$$a\cot(\theta-a)=b\cot(\phi-a)=c\cot(\psi-a).$$

Questions 9, 10, 11, 12 *in Higher Trigonometry.*

XIV. JESUS, CHRIST'S, EMMANUEL AND SIDNEY SUSSEX COLLEGES. *June* 9, 1885. 1—4.

1. Define the *sine* and *tangent* of an angle.

A pyramid has for base a square of side a; its vertex lies on a line through the middle point of the base, perpendicular to it, and at a distance h from it; prove that the angle α between two lateral faces is given by

$$\sin\alpha = \frac{2h\sqrt{2a^2+4h^2}}{a^2+4h^2}.$$

2. Prove geometrically that

(i) $\cos B - \cos A = 2\sin\frac{A+B}{2}\sin\frac{A-B}{2}$.

(ii) $\tan(A-B) = \frac{\tan A - \tan B}{1+\tan A\tan B}$.

If α, β, γ are in Arithmetical Progression, prove that

$$2\left\{\frac{\tan(\alpha+\beta)}{\tan(\beta+\gamma)} - \frac{\tan(\alpha-\beta)}{\tan(\beta-\gamma)}\right\}^{-1} + 1 = \frac{\sin 4\beta}{\sin(\alpha-\gamma)}.$$

3. Investigate the value of cosine 72^0.

Solve the equation

$$32\sin^5 x + 16\cos^4 x + 12\sin 2x\cos x + 20\sin^2 x - 22\sin x = 15.$$

4. Find the value of $\tan 22\frac{1}{2}^0$.

Trace the changes in $\frac{\tan 3\alpha}{\tan^3\alpha}$ from 0 to $\frac{\pi}{2}$, showing that $17+12\sqrt{2}$ is a minimum, and $17-12\sqrt{2}$ is a maximum value.

5. Prove that

$$\sin^3\alpha\sin(\beta-\gamma) + \sin^3\beta\sin(\gamma-\alpha) + \sin^3\gamma\sin(\alpha-\beta) + \sin(\alpha-\beta)\sin(\beta-\gamma)\sin(\gamma-\alpha)\sin(\alpha+\beta+\gamma) = 0.$$

Also, if α, β, γ be the angles of any triangle, prove that

$$\cos 4\alpha(\cos 2\beta - \cos 2\gamma) + \cos 4\beta(\cos 2\gamma - \cos 2\alpha) + \cos 4\gamma(\cos 2\alpha - \cos 2\beta) = 16\sin(\beta-\gamma)\sin(\gamma-\alpha)\sin(\alpha-\beta)\sin\alpha\sin\beta\sin\gamma.$$

6. Prove, with the usual notation, that the area of a triangle is equal to

$$\sqrt{s(s-a)(s-b)(s-c)}.$$

Show that the sum of the areas of the two equilateral triangles, each of which has its vertices at three given distances from a fixed point, is equal to the sum of the equilateral triangles on these distances, and the difference of the aforesaid areas is three times the area of the triangle whose sides are the given distances.

7. If Δ be the area of any quadrilateral whose semiperimeter is s, and sides a, b, c, d, prove

$$\Delta^2=(s-a)(s-b)(s-c)(s-d)-abcd\cos^2\frac{\alpha+\gamma}{2},$$

where α, γ are opposite angles of the figure.

8. Two persons are stationed on two different floors of a house, one vertically above the other, and observe the sun sinking behind a distant tower. From the higher station the sun's lower limb seems to graze the top of the tower. From the lower station the higher limb seems to graze it. Determine the height of the tower in terms of its horizontal distance from the stations, the distance between the stations, and the angular diameter of the sun.

9. Prove that the distance between the centres of the inscribed and circumscribed circles of a triangle is $\sqrt{R^2-2Rr}$, where r, R are radii of these circles.

If the sum of the pairs of radii of the escribed circles of a triangle taken in order round the triangle be denoted by s_1, s_2, s_3, and the corresponding differences by d_1, d_2, d_3, prove that

$$d_1d_2d_3+d_1s_2s_3+d_2s_3s_1+d_3s_1s_2=0.$$

10. Find an expression for the area of a segment of a circle.

With each corner of a square of side a as centre, and a side as radius, a circle is described, show that the area of the curvilinear quadrilateral formed by the intersection of these four circles has an area

$$a^2\left(1-\sqrt{3}+\frac{\pi}{3}\right).$$

Questions 11, 12, 13 *in Higher Trigonometry.*

XV. MATHEMATICAL TRIPOS. PART I. *May* 21st, 1885. 2—5.

Questions 1—6 *Algebra.*

vii. Prove the following formulæ geometrically,

(i) $\sin A+\sin B=2\sin\frac{1}{2}(A+B)\cos\frac{1}{2}(A-B)$,

(ii) $\frac{\pi}{4}=\tan^{-1}\left(\frac{m}{n}\right)-\tan^{-1}\frac{m-n}{m+n}$,

(iii) $\sin^2\alpha+\sin^2\beta=\sin^2(\alpha+\beta)-2\sin\alpha\sin\beta\cos(\alpha+\beta)$.

viii. If $\cot^{-1}x-\cot^{-1}(x+2)=15^0$, find x.

If $$\tan\left(\frac{\pi}{4}+\frac{\psi}{2}\right)=\tan^5\left(\frac{\pi}{4}+\frac{\phi}{2}\right),$$

prove that

$$\sin\psi = 5\sin\phi\,\frac{(1+\alpha^2\sin^2\phi)(1+\beta^2\sin^2\phi)}{\left(1+\frac{1}{\alpha^2}\sin^2\phi\right)\left(1+\frac{1}{\beta^2}\sin^2\phi\right)},$$

and find α and β.

ix. If $A+B+C=\pi$, prove the formulæ,

(i) $$\frac{\tan A}{\tan B\tan C}+\frac{\tan B}{\tan C\tan A}+\frac{\tan C}{\tan A\tan B}$$
$$=\tan A+\tan B+\tan C-2(\cot A+\cot B+\cot C),$$

(ii) $$\sin\tfrac{1}{2}A+\sin\tfrac{1}{2}B+\sin\tfrac{1}{2}C-1$$
$$=4\sin\tfrac{1}{4}(\pi-A)\sin\tfrac{1}{4}(\pi-B)\sin\tfrac{1}{4}(\pi-C),$$

and if $A+B+C=n\pi$,

(iii) $$\sin 2A+\sin 2B+\sin 2C=\pm 4\sin A\sin B\sin C,$$

(iv) $$\cos 2A+\cos 2B+\cos 2C+1=\mp 4\cos A\cos B\cos C,$$

and determine when the upper sign is to be used.

x. Express the area of a triangle in terms of the sides.

Show how to construct the right-angled triangle of minimum area which has its vertices on three parallel lines; and if a, b are the distances of the middle line from the other two, show that the hypotenuse makes with the parallel lines an angle

$$\cot^{-1}\frac{a-b}{a+b}.$$

If the given angle of the triangle instead of being a right angle is equal to α, find the angle which the side opposite to it makes with the parallel lines when the area is a minimum.

xi. Show how to solve a triangle when two sides and the included angle are given.

If two sides of a triangle are 71 and 25 feet and the contained angle $69^0\ 32'$ calculate the remaining angles and side, and show that if a small error has been made in the measurement of the smaller side it will affect the calculated value of the third side very slightly.

xii. Find the radii of the inscribed and escribed circles of a triangle; and if these are r, r_1, r_2, r_3 and that of the circumscribed circle is R, prove that $r_1+r_2+r_3-r=4R$.

If D, E, F are the centres of the escribed circles and O that of the inscribed circle, prove that

$$\frac{EF^2}{r_2r_3}+\frac{FD^2}{r_3r_1}+\frac{DE^2}{r_1r_2}=\frac{OD^2}{rr_1}+\frac{OE^2}{rr_2}+\frac{OF^2}{rr_3}=8\frac{R}{r}.$$

ANSWERS TO THE EXAMPLES.

I. Page 2.

(1) 80. (2) 10. (3) 16. (4) $109\frac{11}{21}$.

(5) 5 acres. (6) $\frac{1760a}{b}$. (7) $\frac{a \cdot c}{3}$ yds.

(8) A shilling and a three-penny piece.

II. Pages 4, 5.

(1) 77440. (2) 6 ft. (3) $\frac{1}{1}\frac{0}{0}\frac{5}{5}$. (4) $6\frac{3}{4}$, $\frac{27}{4}$, $\frac{27}{4}$.

(5) 15·625, $15\frac{5}{8}$, $\frac{1}{8}\frac{2}{2}\frac{5}{5}$, $\frac{1}{8}\frac{2}{2}\frac{5}{5}$. (6) $9\frac{9}{10}$, 9 shillings.

(7) $\frac{80a}{b}$, $\frac{80a}{c}$, $\frac{80a}{d}$, $\frac{80a}{k}$. (8) 21 shillings.

III. Pages 7, 8.

(1) 10 ft. (2) 80 yds. (3) 20 ft. (4) 50 ft.

(5) 90 ft. (6) $20\frac{10}{29}$ nearly. (7) $5a$ feet.

(8) $12a$ yards. (10) $\frac{\sqrt{2}}{2}a$ yards. (12) $\frac{2\sqrt{3}}{3}a$ feet.

(13) $1 : \sqrt{2}$. (14) $\sqrt{84}$ ft. (15) $2\sqrt{9a^2 - b^2}$ ft.

IV. Page 13.

(1) 9·899 ft. (2) 2489 yds. (3) 58·36 in.

(4) 1138 in. (5) 933·4 ft. (6) 2504 in.

(7) 8·66 ft. (8) 4·607 ft. (9) 48 in.

(10) 85 ft. 11 in.

V. Pages 21, 22.

(1) $3\frac{1}{7}$ yds. (2) $25\frac{1}{7}$ ft. (3) $150\frac{6}{7}$ in. (4) $3\frac{2}{11}$ ft.

(5) $7\frac{7}{11}$ ft. (6) 560. (7) $15\frac{1}{2}$ nearly. (8) 33600.

(9) $32\frac{1}{7}$. (10) 7 ft. (11) $553\frac{1}{7}$, 13·8 in. (12) $339\frac{3}{7}$ ft.

(13) 443 in. (14) 235 in. (15) 203 in.

(16) 274 in. (17) 1886 in.

VII. Page 28.

(1) ·632118 of a right angle. (3) ·021827 of a right angle.

(2) 1·0426991 ,, (4) ·03294894 ,,

(5) ·006241 of a right angle. (9) ·6900071 of a right angle.
(6) 10·000812 ,, (10) 1·19030045 ,,
(7) ·3204052 ,, (11) 10·061801 ,,
(8) ·0102034 ,, (12) ·0226048 ,,

(13) $36^g\ 78'\ 91''$. (14) $104^g\ 30'\ 21''$. (15) $1^g\ 20'\ 3''$.
(16) $10'\ 20''$. (17) $6^g\ 25'$. (18) $302^g\ 12'\ 50''$.
(19) $100^g\ 10'$. (20) $1^g\ 1'\ \cdot 1''$. (21) $645^g\ 10'$.
(22) $2^g\ 30'$. (23) $1'\ 10''$. (24) $10''$.

VIII. Page 30.

(1) ·09175 of a right angle $= 9^g\ 17'\ 50''$.
(2) ·0675 ,, $= 6^g\ 75'$.
(3) 1·07875 ,, $= 107^g\ 87'\ 50''$.
(4) $\cdot 1804290\dot{1}2345679\dot{9}$ $= 18^g\ 4'\ 29''$, etc.
(5) $1\cdot 46\dot{7}$,, $= 146^g\ 77'\ 77\cdot\dot{7}''$.
(6) $\cdot 5\dot{4}$,, $= 54^g\ 44'\ 44\cdot\dot{4}''$.
(7) $1^0\ 14'\ 15''$. (8) $7^0\ 52'\ 30''$. (9) $153^0\ 24'\ 29\cdot 34''$.
(10) $21^0\ 36'\ 8\cdot 1''$. (11) $16^0\ 12'\ 37\cdot 26''$. (12) $31^0\ 30'$.

IX. Page 35.

1. (1) 2 right angles or 180^0. (2) $\frac{3}{2}$ of a right angle.
(3) $\frac{2}{\pi}$ right angles. (4) $\frac{6}{\pi}$ right angles.
(5) 2 right angles. (6) $\frac{4}{\pi^2}$ right angles.
(7) $\frac{2\theta}{\pi}$ right angles. (8) ·002 of a right angle.
(9) 20 right angles.

2. (1) π. (2) 2π. (3) $\frac{\pi}{3}$. (4) $\frac{\pi}{8}$.
(5) $\frac{\pi}{180}$. (6) 1^c. (7) $\frac{n}{180}\pi$. (8) $\frac{1}{2}^c$. (9) $\frac{A\pi}{180}$.

3. (1) $\frac{\pi}{6}$. (2) $\frac{\pi}{4}$. (3) $\frac{\pi}{12}$. (4) $\frac{\pi}{200}$.
(5) $\frac{\pi}{20000}$. (6) $\frac{\pi}{200000}$. (7) $\frac{n\pi}{200}$. (8) 1^c. (9) 5π.

4. (1) $\frac{1}{8}$. (2) $\frac{1}{9}^0$. (3) 1. (4) $\frac{3\pi}{50}$. (5) $\frac{8}{2}\frac{3}{0}$. (6) $\frac{\pi}{180}$

X. Pages 37—39.

(1) $\frac{3}{5}$. (2) 90. (3) $4\frac{4}{5}$. (4) $\frac{70}{11}$.
(5) $112\frac{1}{2}$ ft. (6) $5\frac{37}{55}$ ft. (7) $9\frac{3}{7}$ ft.
(8) 838000 miles. (9) $\frac{1}{9}$ radian $= 6\frac{4}{11}$ degrees.
(10) $21\frac{21}{44}$ degrees. (11) $51\frac{6}{11}''$. (12) about 34 yds.
(13) 1 : 3·1416. (14) 3·1416. (15) 3·1416.
(16) 400 : 1. (17) ·0000484.... (18) $49\frac{1}{11}$ in.
(19) $\frac{\pi}{2}$ i.e. a right angle. (21) 473 : 489.
(22) (i) $k=1$, (ii) $k=\frac{180}{\pi}$.

XI. Pages 40—43.

(1) $\frac{\pi}{10}$. (2) 33°, 15°. (3) 90°, 45°. (4) 18°, $22\frac{1}{2}$°.
(5) 38°, 18°. (7) 45°, 60°, 75°. (8) 39°, 60°, 81°.
(9) 33·3̇ᵍ, 66·6̇ᵍ, 100ᵍ. (14) 25°, 8° 6′. (15) $\frac{2}{25}$.
(16) $\frac{n\pi}{10800}$. (17) (i) 120°, 133·3̇ᵍ, $\frac{2\pi}{3}$, (ii) 135°, 150ᵍ, $\frac{3\pi}{4}$,
(iii) 156°, 173·3̇ᵍ, $\frac{13\pi}{15}$. (20) 450′. (21) (i) $3\frac{3}{4}$, (ii) $\frac{15}{2\pi}$.
(22) $1\frac{9}{10}$°. (23) a right angle. (24) $\frac{10ac}{9b}$.
(25) $\frac{9a+10b}{10c}$ degrees. (26) $\frac{5\pi}{9}$. (27) $\frac{5\pi}{19}$, $\frac{9\pi}{38}$, $\frac{\pi}{2}$.
(28) $\frac{1800\pi}{19\pi+1800}$. (29) $\frac{\pi^2}{10\pi+18000}$. (30) 9 or 16.

XII. Pages 50—52.

(1) (i) DA, BD. (ii) DB, AD. (iii) DA, CD. (iv) DC, AD.
(2) (i) $\frac{DB}{AB}$. (ii) $\frac{DC}{CA}$. (iii) $\frac{CD}{AD}$. (iv) $\frac{DA}{BA}$. (v) $\frac{DB}{AD}$.
(vi) $\frac{DC}{AC}$. (vii) $\frac{CD}{CA}$. (viii) $\frac{DA}{CD}$. (ix) $\frac{BD}{BA}$. (x) $\frac{DA}{CA}$.
(3) (i) $\frac{DB}{CB}$, $\frac{BA}{CA}$. (ii) $\frac{CD}{CB}$, $\frac{CB}{CA}$. (iii) $\frac{DB}{CD}$, $\frac{BA}{CB}$.
(iv) $\frac{DB}{AB}$, $\frac{BC}{AC}$. (v) $\frac{AD}{AB}$, $\frac{AB}{AC}$. (vi) $\frac{DB}{AD}$, $\frac{BC}{AB}$.

(4) (i) $\frac{DA}{BA}$. (ii) $\frac{BA}{EA}$ or $\frac{AC}{EC}$. (iii) $\frac{DC}{BC}$. (iv) $\frac{AB}{AE}$.
(v) $\frac{AD}{AB}$ or $\frac{AB}{AC}$. (vi) $\frac{BD}{BC}$. (vii) $\frac{DB}{CD}$, or $\frac{BA}{CB}$, or $\frac{AE}{CA}$.
(viii) $\frac{DA}{BD}$. (ix) $\frac{BA}{EB}$ or $\frac{AC}{EA}$. (x) $\frac{DC}{BD}$. (xi) $\frac{DB}{AB}$ or $\frac{BC}{AC}$.
(xii) $\frac{BE}{AE}$.

(5) $\sin A=\frac{3}{5}$, $\cos A=\frac{4}{5}$, $\tan A=\frac{3}{4}$; $\sin B=\frac{4}{5}$, $\cos B=\frac{3}{5}$, $\tan B=\frac{4}{3}$.
(7) Of the smaller angle, the sine $=\frac{5}{13}$, cosine $=\frac{12}{13}$, tangent $=\frac{5}{12}$.
Of the larger angle, the sine $=\frac{12}{13}$, cosine $=\frac{5}{13}$, tangent $=\frac{12}{5}$.

(8) Of the smaller angle, the sine $=\frac{1}{2}$, cosine $=\frac{\sqrt{3}}{2}$, tangent $=\frac{1}{\sqrt{3}}$.
Of the larger angle, the sine $=\frac{\sqrt{3}}{2}$, cosine $=\frac{1}{2}$, tangent $=\sqrt{3}$.

XIV. Pages 63—65.

(1) 179 ft. (2) 346 ft. (3) 86·6 ft. (4) 138·5 ft.
(5) $7\frac{1}{2}$ ft. (6) 60^0, 173 ft. (7) 63·17 yds. (8) 34·15 ft.
(9) 73·2 ft. (10) 86·6 ft. (11) ·866 miles $=1524$ yds.
(12) 173·2 yds. (14) 373 ft. (15) 3733 ft.
(16) $\frac{\sqrt{6}}{2}$ miles $=6465$ ft. (17) $\frac{\sqrt{3}.a}{3b}$. (18) 30^0.
(19) about 523·6 miles.

XVI. Page 74.

(1) $\sin A=\sqrt{1-\cos^2 A}$, $\tan A=\frac{\sqrt{1-\cos^2 A}}{\cos A}$, $\cot A=\frac{\cos A}{\sqrt{1-\cos^2 A}}$,
$\sec A=\frac{1}{\cos A}$, $\operatorname{cosec} A=\frac{1}{\sqrt{1-\cos^2 A}}$.

(2) $\sin A=\frac{1}{\sqrt{1+\cot^2 A}}$, $\cos A=\frac{\cot A}{\sqrt{1+\cot^2 A}}$, $\tan A=\frac{1}{\cot A}$,
$\sec A=\frac{\sqrt{1+\cot^2 A}}{\cot A}$, $\operatorname{cosec} A=\sqrt{1+\cot^2 A}$.

(3) $\sin A=\frac{\sqrt{\sec^2 A-1}}{\sec A}$, $\cos A=\frac{1}{\sec A}$, $\tan A=\sqrt{\sec^2 A-1}$,
$\cot A=\frac{1}{\sqrt{\sec^2 A-1}}$, $\operatorname{cosec} A=\frac{\sec A}{\sqrt{\sec^2 A-1}}$.

(4) $\sin A = \dfrac{1}{\operatorname{cosec} A}$, $\cos A = \dfrac{\sqrt{\operatorname{cosec}^2 A - 1}}{\operatorname{cosec} A}$, $\tan A = \dfrac{1}{\sqrt{\operatorname{cosec}^2 A - 1}}$,

$\cot A = \sqrt{\operatorname{cosec}^2 A - 1}$, $\sec A = \dfrac{\operatorname{cosec} A}{\sqrt{\operatorname{cosec}^2 A - 1}}$.

(5) $\cos A = \sqrt{1 - \sin^2 A}$, $\tan A = \dfrac{\sin A}{\sqrt{1 - \sin^2 A}}$, $\cot A = \dfrac{\sqrt{1 - \sin^2 A}}{\sin A}$,

$\sec A = \dfrac{1}{\sqrt{1 - \sin^2 A}}$, $\operatorname{cosec} A = \dfrac{1}{\sin A}$.

(6) $\sin A = \dfrac{\tan A}{\sqrt{1 + \tan^2 A}}$, $\cos A = \dfrac{1}{\sqrt{1 + \tan^2 A}}$, $\cot A = \dfrac{1}{\tan A}$,

$\sec A = \sqrt{1 + \tan^2 A}$, $\operatorname{cosec} A = \dfrac{\sqrt{1 + \tan^2 A}}{\tan A}$.

XVII. Page 75.

(1) $\frac{3}{4}$, $\frac{5}{3}$. (2) $\dfrac{2\sqrt{2}}{3}$, $\dfrac{1}{2\sqrt{2}}$. (3) $\frac{4}{5}$, $\frac{5}{3}$.

(4) $\dfrac{1}{\sqrt{15}}$, $\dfrac{\sqrt{15}}{4}$. (5) $\dfrac{\sqrt{3}}{2}$, $\frac{1}{2}$. (6) $\dfrac{\sqrt{5}}{3}$, $\frac{3}{2}$.

(7) $\dfrac{b}{\sqrt{c^2 - b^2}}$. (8) $\dfrac{a}{\sqrt{a^2 + b^2}}$, $\dfrac{b}{\sqrt{a^2 + b^2}}$.

(9) $\dfrac{\sqrt{a^2 - 1}}{a}$, $\dfrac{1}{\sqrt{a^2 - 1}}$. (11) $h^2(1 + k^2) = 1$.

XVIII. Page 77.

(2) $\sec\theta$ increases continuously from 1 to ∞.

(3) $\sin A$ diminishes continuously from 1 to 0.

(4) $\cot\theta$ diminishes continuously from ∞ to 0.

XIX. Page 82.

(4) Yes. (5) No. (6) Yes.

(7) (i) 60^0. (ii) -100^0. (iii) 0^0. (iv) -260^0. (v) 115^0.

(vi) 410^0. (vii) $-\dfrac{\pi}{4}$. (viii) $\dfrac{2\pi}{3}$.

XX. Page 84.

(1) 45^0. (2) 30^0. (3) 45^0. (4) 60^0.

(5) 30^0. (6) 30^0. (7) 30^0. (8) 0^0, or 45^0.

(9) 90^0, or 60^0. (10) 60^0. (11) 45^0. (12) 45^0.
(13) 90^0, or 45^0. (14) 18^0. (15) 45^0. (16) 45^0.
(17) 30^0. (18) 30^0.

XXI. Page 85.

(3) the value 3 is inadmissible. (4) $\frac{2\pm\sqrt{2}}{4}$.

(5) $\frac{3}{4}$, or $\frac{1}{2}$. (6) $\frac{3}{4}$, or $\frac{1}{3}$.

(7) the value $-\frac{7\sqrt{3}}{6}$ is inadmissible. (9) $1-\sin^4\theta$.

(10) $1-3\sin^2\theta+3\sin^4\theta$. (11) $\frac{1-2\cos^2\theta+2\cos^4\theta}{\cos^4\theta}$.

(13) $\frac{1-\sin A}{1+\sin A}$.

(14) $\operatorname{cosec}\theta$ decreases continuously from ∞ to 1.

(15) $\cot\theta$ increases continuously from 0 to ∞.

(16) $\theta=\frac{\pi}{4}$, $\phi=\frac{\pi}{12}$.

XXII. Page 89.

(1) +6. (2) 0. (3) +2. (4) +3.
(5) +10. (6) 0. (7) +7. (8) +7.

XXIV. Page 94.

(1) The second. (2) The fourth. (3) The second.
(4) The third. (5) The fourth. (6) The first.
(7) The second. (8) The first. (9) The first.
(10) The fourth. (11) The fourth.
(12) The first, if n be even, the third, if n be odd.

XXV. Page 98.

(1) +, +, +. (2) +, −, −. (3) −, −, +.
(4) −, +, −. (5) −, +, −. (6) −, −, +.
(7) +, −, −. (8) +, +, +. (9) +, +, +.
(10) +, +, +. (11) +, −, −. (12) −, +, −.

XXVI. Page 100.

(1) $+\frac{1}{2}$, $-\frac{\sqrt{3}}{2}$, $-\frac{1}{\sqrt{3}}$. (2) $+\frac{1}{\sqrt{2}}$, $-\frac{1}{\sqrt{2}}$, -1.

(3) $+\frac{\sqrt{3}}{2}$, $-\frac{1}{2}$, $-\sqrt{3}$. (4) $-\frac{1}{2}$, $+\frac{\sqrt{3}}{2}$, $-\frac{1}{\sqrt{3}}$.

(5) $-\frac{1}{\sqrt{2}}, +\frac{1}{\sqrt{2}}, -1.$ (6) $+\frac{\sqrt{3}}{2}, +\frac{1}{2}, +\sqrt{3}.$

(7) $-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, +1.$ (8) $-\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}, +1.$

(9) $+\frac{1}{2}, +\frac{\sqrt{3}}{2}, +\frac{1}{\sqrt{3}}.$ (10) $+\frac{1}{2}, +\frac{\sqrt{3}}{2}, +\frac{1}{\sqrt{3}}.$

(11) $-\frac{\sqrt{3}}{2}, -\frac{1}{2}, +\sqrt{3}.$ (12) $-\frac{\sqrt{3}}{2}, +\frac{1}{2}, -\sqrt{3}.$

(13) $+\frac{1}{\sqrt{2}}, +\frac{1}{\sqrt{2}}, +1.$ (14) $\frac{\sqrt{3}}{2}, -\frac{1}{2}, -\sqrt{3}.$

(15) $-\frac{1}{2}, -\frac{\sqrt{3}}{2}, +\frac{1}{\sqrt{3}}.$

XXVII. Page 100.

Each of these expressions changes *continuously* as the angle A increases from 0^0 to 360^0, and their values are *repeated* at each complete revolution of the revolving line.

Their values are given below when they are zero and when they cease to increase and begin to decrease, and vice versa. The first table gives also the sign of each between the values.

	0^0		90^0		180^0		270^0	
(1) $\cos A$	$+1$	$+$	0	$-$	-1	$-$	0	$+$
(2) $\tan A$	0	$+$	∞	$-$	0	$+$	∞	$-$
(3) $\cot A$	∞	$+$	0	$-$	∞	$+$	0	$-$
(4) $\sec A$	$+1$	$+$	∞	$-$	-1	$-$	∞	$+$
(5) $\operatorname{cosec} A$	∞	$+$	$+1$	$+$	∞	$-$	-1	$-$
(6) $1-\sin A$	$+1$	$+$	0	$+$	$+1$	$+$	2	$+$
(7) $\sin^2 A$	0	$+$	$+1$	$+$	0	$+$	$+1$	$+$

	0^0	45^0	90^0	135^0	180^0	225^0	270^0	315^0
(8) $\sin A \cdot \cos A$	0	$+\frac{1}{2}$	0	$-\frac{1}{2}$	0	$+\frac{1}{2}$	0	$-\frac{1}{2}$
(9) $\sin A + \cos A$	$+1$	$+\sqrt{2}$	$+1$	0	-1	$-\sqrt{2}$	-1	0
(10) $\tan A + \cot A$	∞	$+2$	∞	-2	∞	$+2$	∞	-2
(11) $\sin A - \cos A$	-1	0	$+1$	$+\sqrt{2}$	$+1$	0	-1	$-\sqrt{2}$

The following figures exhibit the changes in the sign and magnitude of $\sin\theta$ (fig. i.), $\cos\theta$ (fig. ii.), and $\tan\theta$ (fig. iii.). The measure of the distance from 0 along the line 0X = the circular measure of the angle; the vertical distance from 0X measures the Ratio.

Fig. i. The Curve of the Sine.



Fig. ii. The Curve of the Cosine.



Fig. iii. The Curve of the Tangent.



XXXI. Page 112.

(1) (i) $30^0, 150^0, -210^0, -330^0$. (ii) $45^0, 135^0, -225^0, -315^0$.
(iii) $60^0, 120^0, -240^0, -300^0$. (iv) $-30^0, -150^0, 210^0, 330^0$.

(2) (i) $20^0, 160^0, 380^0, 520^0$. (ii) $\frac{5\pi}{4}, \frac{7\pi}{4}, \frac{13\pi}{4}, \frac{15\pi}{4}$.

(iii) $\frac{8\pi}{7}, \frac{13\pi}{7}, \frac{22\pi}{7}, \frac{27\pi}{7}$.

(3) (i) $\theta = n\pi + (-1)^n\left(-\frac{\pi}{6}\right)$. (ii) $\theta = n\pi + (-1)^n\frac{\pi}{6}$.

(iii) $n\pi \pm \frac{\pi}{4}$.

XXXII. Page 115.

(1) (i) $\theta = 2n\pi \pm \frac{1}{3}\pi$. (ii) $\theta = n\pi + \frac{1}{4}\pi$. (iii) $\theta = n\pi - \frac{1}{4}\pi$.
(iv) $\theta = n\pi - \frac{1}{3}\pi$. (v) $\theta = 2n\pi \pm \frac{4}{5}\pi$. (vi) $\theta = n\pi + \frac{3}{4}\pi$.
(3) The tangent. (4) No.
(5) (i) 60^0, -60^0, 300^0, -300^0. (ii) 45^0, 225^0, 405^0, -135^0.
(iii) -45^0, -225^0, 135^0, 315^0. (iv) -60^0, -240^0, 120^0, 300^0.
(v) 144^0, -144^0, 216^0, -216^0. (vi) 135^0, -45^0, -225^0, 315^0.

XXXIV. Page 120.

(5) 1, $\frac{24}{25}$. (6) $\frac{16}{65}$, $\frac{33}{65}$.

XXXVIII. Page 131.

(1) $\sin(\theta+\phi)+\sin(\theta-\phi)$. (2) $\cos(\alpha-\beta)+\cos(\alpha+\beta)$.
(3) $\sin(2\alpha+3\beta)+\sin(2\alpha-3\beta)$. (4) $\cos 2\alpha+\cos 2\beta$.
(5) $\sin 8\theta - \sin 2\theta$. (6) $\cos\theta+\cos 2\theta$. (7) $\frac{1}{2}(\cos 3\theta - \cos 5\theta)$.
(8) $\frac{1}{2}(\sin 4\theta - \sin\theta)$. (9) $\sin 60^0 + \sin 40^0$.
(10) $\frac{1}{2}(\sin 60^0 - \sin 30^0)$. (11) $2\cos 3\theta \cos 2\theta$.
(12) $-\cos 4\theta \sin 2\theta$. (13) $4\cos^2\frac{\theta}{2}\sin 2\theta$.

XXXIX. Page 132.

(4) $\tan 2\alpha$. (5) $\tan x$. (7) $-2\cot 2x$. (8) $\tan^2 A$.
(12) $\sin 2\theta = 1$, $\therefore\ \theta = 45^0$; $\sin 2\phi = \frac{1}{2}$, $\therefore\ \phi = 15^0$.

XLV. Page 147.

(2) $\sin 2A = \frac{4}{5}$, $\sin 3A = -\frac{11}{5}\sqrt{5}$.
(3) $\sin 2\theta = -\frac{1}{8}\sqrt{15}$, $\sin 3\theta = \frac{3}{16}\sqrt{15}$, $\cos 3\theta = -\frac{11}{16}$; in the second.

XLVI. Page 149.

(3) $\sin\frac{A}{2} = +\sqrt{\frac{1-\cos A}{2}}$. (5) $\sin\frac{A}{2} = +\sqrt{\frac{1-\cos A}{2}}$.

XLVII. Page 154.

(1) (i) +, −. (ii) −, +. (iii) −, −. (iv) +, −. (v) +, −.
(vi) +, +. (vii) −, −. (viii) −, −.
(3) $\sin 9^0 = \frac{1}{4}\{\sqrt{(3+\sqrt{5})} - \sqrt{(5-\sqrt{5})}\}$,
$\cos 9^0 = \frac{1}{4}\{\sqrt{(3+\sqrt{5})} + \sqrt{(5-\sqrt{5})}\}$,
$\sin 81^0 = \cos 9^0$, $\cos 189^0 = -\cos 9^0$, $\tan 202\frac{1}{2}^0 = \sqrt{2}-1$,
$\tan 97\frac{1}{2}^0 = -(\sqrt{3}+\sqrt{2})(\sqrt{2}+1)$.
(7) Between $(2n \times 180^0 - 45^0)$ and $(2n \times 180^0 + 45^0)$.

XLVIII. Page 157.

(1) $\theta = -\frac{\pi}{4} + n\pi + (-1)^n \frac{\pi}{6}$. (2) $\theta = -\frac{\pi}{3} + n\pi + (-1)^n \frac{\pi}{6}$.

(3) $\theta = -\frac{\pi}{4} + n\pi + (-1)^n \frac{\pi}{3}$. (4) $\theta = +\frac{\pi}{4} + n\pi + (-1)^n \frac{\pi}{4}$.

(5) $\theta = +\frac{\pi}{4} + 2n\pi \pm \frac{\pi}{4}$. (6) $\theta = \frac{\pi}{6} + n\pi \pm \frac{\pi}{4}$.

(7) $x = -68^0\,12' + n \times 180^0 + (-1)^n\,(21^0\,48')$.

(8) $x = -69^0\,26'\,30'' + 2n \times 180^0 \pm (69^0\,26'\,30'')$.

(9) $x = 75^0\,4' + n \times 180^0 + (-1)^n\,(14^0\,56')$.

(10) $x = -\alpha - \frac{\pi}{4} + 2n\pi \pm \beta$.

L. Pages 159, 160.

(7) $\pm(\sqrt{3}+1)$. (8) $\pm\sqrt{2}$. (9) ± 1.

(13) $\frac{1}{6}(2n+1)\pi$; or, $\frac{1}{3}(3n \pm 1)\pi$. (14) $n\pi \pm \frac{\pi}{3}$; or $\frac{1}{6}(2n+1)\pi$.

(15) $\frac{1}{8}(2n+1)\pi$; or $\frac{n\pi}{3} + (-1)^n \frac{\pi}{12}$.

(16) $(2n+1)\pi$; or, $\cos^{-1}\frac{a-b}{a}$. (17) $\frac{n\pi}{m}$; or $\frac{2r+1}{m \pm n}.\pi$.

(18) Both equations are satisfied if $(x^2 - y^2) =$ an odd integer; or, if $(x+y)^2 = 2n$, and $(x-y)^2 = 4m - 2n$, m and n being integers.

(19) (i) $2\sin\theta.\cos\theta = \sin 2\theta$, and therefore it goes through the same changes as $\sin\theta$, while θ changes from 0 to 2π. (ii) $\cos^2\theta - \sin^2\theta = \cos 2\theta$. (iii) $\sin 3\theta$ goes through the same changes as $\sin\theta$, while θ goes from 0 to 3π. (iv) $\cot 2A$; compare with $\cot A$. (v) $\sin(\theta+\alpha)$ goes through the same changes while θ goes from 0 to π, as $\sin\theta$ goes through while θ goes from α to $\pi+\alpha$. (vi) $\cos(2\theta - \alpha)$ goes through the same changes while θ goes from 0 to π, as $\cos 2\theta$ goes through as θ changes from $-\frac{\alpha}{2}$ to $\pi - \frac{\alpha}{2}$. (See Ans. to XXVII.)

(20) They are the solutions of the equations $\sin\theta = \sin\alpha$, and $\cos\left(\frac{\pi}{2} - \theta\right) = \cos\left(\frac{\pi}{2} - \alpha\right)$; and we know that $\cos\left(\frac{\pi}{2} - \theta\right) = \sin\theta$.

(21) They are the solutions of the equations $\sin\left(\theta + \frac{\pi}{4}\right) = \sin\frac{\pi}{6}$, and $\cos\left(\theta - \frac{\pi}{4}\right) = \cos\frac{\pi}{3}$; also $\cos\left(\theta - \frac{\pi}{4}\right) = \sin\left(\theta + \frac{\pi}{4}\right)$, $\sin\frac{\pi}{6} = \cos\frac{\pi}{3}$.

LI. Page 162.

(1) (i) a^{2h+3k}, (ii) a^{4h-5k}, (iii) $a^{\frac{4h}{3}+\frac{5k}{3}}$. (iv) $a^{\frac{5h}{2}+\frac{3k}{2}}$.
(2) (i) 5·4690116. (ii) 10·6243928. (iii) 13·7509366.
(iv) ·8853661. (v) 1·7968680. (vi) 8·9699598.
(vii) 2·7345058.
(3) 2^3, 2^5, 2^{-1}, 2^{-4}, 2^{-3}, 2^7. (4) 3^2, 3^4, 3^{-1}, 3^{-3}, 3^{-2}, 3^{-4}.

LII. Page 163.

(1) ·60206, ·9542426, ·90309, ·7781513, 1·20412, 1·690196.
(2) 1·146128, 1·20412, 1·2552726, 1·3802113, 1·4313639, 1·6232493.
(3) 1, ·69897, 1·1760913, 1·39794, 1·4771213, 1·5440680.
(4) 1·5563026, 1·60206, 1·6812413, 1·69897, 2·30103, 3.
(5) 7·201593, 3·858708. (6) ·7545579, 2·989843.
(7) 1·4532. (8) 2408·6. (9) (i) 4·5868. (ii) ·93646.
(10) 3·9549. (11) 40975·3 sq. ft. (12) 34·925 in.
(13) 3·2617 in. (14) 110115 cub. yds.

LIII. Page 165.

(1) 3, $\frac{10}{3}$, $\frac{1}{4}$, $\frac{2}{3}$, $-\frac{5}{2}$. (2) 3, 6, −1, −3, −6, 2.
(3) 2, 4, −1, −3, −2, −4. (4) $\frac{3}{2}$, $\frac{2}{3}$, $-\frac{1}{4}$, −1.
(5) 3, −1, 5, −2, 3, −3. (6) $\frac{4}{3}$, $\frac{2}{3}$, $\frac{1}{3}$, $\frac{1}{3}$, $\frac{1}{2}$.
(7) ·7781513, 1·6232493, 1·20412.
(8) 1·6901960, 1·5563026, 1·7993406.
(9) 2·30103, 2·7781513, 1·845098.
(10) ·69897, 5228787, 1·69897.
(11) 1·544068, 2·1760913, −1+·30103.
(12) ·5440680, ·8627278, −2+·9084852.

LIV. Pages 168, 169.

(1) 4, 2, 0, 5, 1. (2) −2, −5, −1, −3.
(3) 3, −1, 0, 1, 0, −7. (4) 4, 1, 6, 3.
(5) the second decimal place, the first dec. pl., the sixth dec. pl.
(6) ten thousands, units, hundreds, third dec. pl., first dec. pl., units.
(7) 10, 4, 25, 31. (8) 9, 11, 85, 4, 9, 6.
(9) units, fourth dec. pl., thousands, seventh dec. pl., second dec. pl.
(10) tenth integral pl., twelfth dec. pl., fifth dec. pl., units, twelfth dec. pl., first dec. pl.

LV. Pages 171, 172.

(1) 2·8901023, ·8901023, 4·8901023, 5·8901023.

(2) 6·7714552, ·7714552, $\bar{4}$·7714552, 2·7714552, 3·7714552.

(3) ·27724... (4) ·00001638... (5) ·77448... (6) ·005968...

LVI. Page 174.

(1) Divide each log by 3. (2) Multiply each log by 2.

(3) Divide each log by ·30103.

(4) Multiply each log by ·4771213.

(5) Divide each log by ·4771213. (6) 3·32190...

(7) 1·183... (8) 1·10730... ·66438...

LVII. Pages 175, 176.

(1) 3, 0, $\frac{1}{5}$, 0, $\frac{7}{6}$. (4) $\bar{1}$·8121177, 55.

(5) ·51375. (6) 7, 4, 3, 3.

(7) (i) $x=\dfrac{2\log 7}{\log 2+4\log 3}$. (ii) $x=\dfrac{7\log 2+4\log 7}{2\log 3+\log 7}$.

(iii) $x=\dfrac{2\log 7}{2\log 2+\log 3}$. (iv) $x=\dfrac{4(\log 3+\log 7)}{8\log 2+3(\log 3+\log 7)}$.

(8) $2+\dfrac{1}{\log_{10}7}$. (9) $\frac{3}{2}+\dfrac{1}{2\log_{10}3}$. (10) $\dfrac{1}{1-\log_{10}2}$.

(11) 0, $\dfrac{1}{b+1}$, $\dfrac{3a}{2b+2}$, $\dfrac{2}{b+1}$, $\dfrac{b}{b+1}$, $\dfrac{3a+2}{2b+2}$, $\dfrac{bc}{b+1}$.

(12) 63 − 31 = 32. (13) $(a^{11}-a^{10})$ integers. (14) 1·9485 nearly.

(19) 2·53855. (20) 4·59999. (21) 167 years.

LVIII. Pages 181, 182.

(1) ·8839066. (2) 2·7513738. (3) $\bar{4}$·9413333.

(4) 6·8086920. (5) ·5710750. (6) 3·70404.

(7) 45740·26. (8) 2492837. (9) ·000439658. (10) 5·689158.

LIX. Pages 184, 185.

(1) ·6737652. (2) ·6737652 (3) ·9306572.

(4) 41° 48′ 37″. (5) 70° 31′ 43·6″. (6) 75° 31′ 21″.

(7) 9·8515594. (8) 9·7114477. (9) 10·1338768.

(10) 35° 4′ 23″. (11) 28° 16′ 27·5″. (12) 21° 56′ 41″.

LX. Pages 188, 189.

(1) 34° 19′ 31·8″. (2) 1498·2 ft. (3) 45° 36′ 56″.

(4) 5293·4 ft., 6982·3 ft. (5) 576·2 chains.

(6) 4729 chains. (7) 3666·8 feet. (8) 42° 15′, 11444 chains.

LXI. Page 190.

(1) 3842·9 ft. (2) 281·74 ft. (3) 115·3 ft. (4) 285·6 ft.

(5) 58° 17′ 24″, 31° 42′ 36″. (6) 656·1 chains, 41° 17′ 12″.

(7) 81 ft. (8) 1942 ft. (9) 646·7 miles. (10) 1000 ft.

LXVI. Pages 208, 209.

(1) 41° 16′ 51·5″. (2) 73° 32′ 12″, 62° 46′ 18″.

(3) 29° 17′ 16″, 31° 55′ 31″. (4) 64° 31′ 58″.

(5) 73°, 23′ 54·4″. (6) 41° 24′ 34·6″. (7) 82° 49′ 9″.

(8) 75°, 60°, 45°. (9) 135°, 30°, 15°.

LXVII. Page 211.

(1) 313·46 yds. (2) 28·87 inches, 31·43 inches.

(3) 1192·55 yds. (4) 22·415 ft.

(5) 24·995 = 25 ft. nearly, 17·559 ft., 65° 59′ 42″.

LXVIII. Pages 213, 214.

(1) 108° 36′ 30″, 31° 23′ 30″. (2) 93° 11′ 49″, 36° 48′ 11″.

(3) 57″ 27′ 25·4″, 62° 32′ 34·6″. (4) 64° 26′ 47″, 37°, 7′ 13″.

(5) 72° 12′ 59″. (6) 20·5 chains. (7) 122·7.

(8) 74° 13′ 50″, 35° 16′ 10″.

LXIX. Pages 218, 219.

(1) $A = 91° 18′ 21″$, $C = 88° 41′ 39″$; or $A = 128° 41′ 39″$, $C = 11° 18′ 21″$.

(2) $B = 70° 0′ 56″$, $C = 59° 59′ 4″$; or, $B = 109° 59′ 4″$, $C = 20° 0′ 56″$.

(3) $B = 38° 38′ 24″$, $C = 91° 21′ 36″$, $c = 155·3$. (4) 61° 16′ 10″.

(5) $A = 72° 4′ 48″$, $B = 41° 56′ 12″$; or, $A = 107° 55′ 12″$,
$B = 6° 5′ 48″$, $b = 17·56$. (6) β is ambiguous; 60·3893 ft.

LXX. Page 220.

The angles are given correct to the nearest second.

(1) $28^\circ 35' 39''$. (2) $104^\circ 44' 39''$. (3) $32^\circ 20' 48''$.
(4) $43^\circ 40'$. (5) $128^\circ 23' 13''$. (6) 106531 ft.
(7) 3437·6 yds. (8) 1728·2 chains. (9) 25376 yards.
(10) $A = 66^\circ 27' 48''$, $B = 12^\circ 55' 12''$. (11) $A = 92^\circ 12' 53''$, $B = 35^\circ 37' 7''$.
(12) $B = 29^\circ 1' 40''$, $C = 74^\circ 55' 50''$. (13) $B - 70^\circ 35' 24''$; or, $109^\circ 24' 36''$.
(14) $B = 51^\circ 56' 17''$; or, $128^\circ 3' 43''$. (15) $B = 62^\circ 6' 10''$; or, $117^\circ 53' 50''$.
(16) Very nearly 90°. (17) 1319·6 yds.

LXX b. P. 220 (i), (ii).

(1) $\cos A = \frac{1}{2}$, $\cos \frac{1}{2}A = \frac{1}{2}\sqrt{3}$. (2) 45°, 60°, 75°.
(3) 135°, 30°, 15°. (4) 3. (5) 14. (6) $1 + \sqrt{3}$. (7) 120°.
(8) 120°. (9) 120°. (10) 90°, $36^\circ 52'$. (11) $130^\circ 27'$. (12) $125^\circ 6'$.
(13) 120°. (14) $A = 54^\circ$ or 126°, $B = 108^\circ$ or 36°. (15) $a = 1$.
(16) $C = 30^\circ$, $a = \sqrt{3} + 1$, $b = 2$. (17) $A = 75^\circ$, $a = b = 2\sqrt{3} + 1$.
(18) $C = 60^\circ$ or 120°. (19) $100\sqrt{3}$. (20) no.
(22) $A = 105^\circ$, $C = 60^\circ$, $B = 15^\circ$. (23) $\frac{1}{2}\sqrt{3}(\sqrt{5} + 1)$.
(24) $A = 90^\circ$ or 60°, $C = 75^\circ$ or 105°, $a = 2\sqrt{2}$ or $\sqrt{6}$. (25) 30° or 150°. (26) $A = 45^\circ$ or 135°, $B = 30^\circ$ or 120°, $b = \sqrt{2}(1 + \sqrt{3})$ or $\sqrt{6}(1 + \sqrt{3})$. (27) 60°, 75°, 6 yds. (28) It is impossible.
(30) $15 : 8\sqrt{3} : 4\sqrt{3} + 9$.

LXXII. Page 229.

(15) To find the point E in an unlimited straight line CE at which a finite straight line AB subtends the greatest angle, a circle must be described passing through A and B, and touching the line CE in the point E. In (15) the centre of the circle lies vertically above E, and in the horizontal line through the middle point of AB.

LXXIII. Page 239.

(1) (i) 10 sq. ft. (ii) 43·3 sq. in. (iii) 148·13 sq. yds.
(iv) 84 sq. chains = 8·4 acres. (v) 100 sq. ft.
(vi) 151872 sq. yds.
(2) 4, $10\frac{1}{2}$, 12, 14 ft. (4) $8\frac{1}{3}$ ft.

LXXVI a. Pages 253—268.

(1) $\sin A = \frac{3}{5}$, $\cos A = \frac{4}{5}$. (3) $A = n \times 180^0$; or, $360^0 \pm 60^0$.

(6) 4227·47 feet.

(9) 30^0, 60^0, 90^0, 120^0, etc. have for sine $\frac{1}{2}$, $\sqrt{\frac{3}{2}}$, 1, $\sqrt{\frac{3}{2}}$, $\frac{1}{2}$, 0, $-\frac{1}{2}$, $-\sqrt{\frac{3}{2}}-1$, $-\sqrt{\frac{3}{2}}$, $-\frac{1}{2}$ respectively.

(12) The other sides are 765·4321 ft.; 1006·6 ft.

(15) 30^0, 60^0, 90^0, etc. have for tan $\frac{1}{3}\sqrt{3}$, $\sqrt{3}$, ∞, $-\sqrt{3}$, $-\frac{1}{3}\sqrt{3}$, 0, $\frac{1}{3}\sqrt{3}$, ∞, $-\sqrt{3}$, $-\frac{1}{3}\sqrt{3}$ respectively.

(17) $\frac{168}{193}$, $\frac{168}{195}$, $\frac{32592}{193 \times 195}$. (19) $\sec A = \frac{8}{4}\sqrt{2}$.

(21) $\theta = \frac{1}{4}(2n+1)\pi$. (22) $\cos 5\alpha = 16\cos^5\alpha - 20\cos^3\alpha + 5\cos\alpha$.

(23) Both signs negative. (24) $\frac{3}{5}\frac{1}{3}\frac{3}{8}$

(25) $\frac{11}{6^0 8^0}A$ radians; $5{\cdot}72956^0$.

(26) sine, $\frac{4}{5}$; tan, $\frac{4}{3}$; cot, $\frac{3}{4}$; cosec, $\frac{5}{4}$; sec, $\frac{5}{3}$.

(31) $\frac{8}{11^0}\theta$ deg.; $19{\cdot}09854^0$. (36) $C = 18^0$, $a = c = \dfrac{2}{\sqrt{(10+2\sqrt{5})}}$.

(37) -1320^0. (38) $-\frac{2}{3}\sqrt{2}$. (39) $(2n+1)\pi$; or, $2n\pi \pm \frac{1}{3}\pi$.

(41) 1 foot, 120^0, 30^0; or 2 feet, 60^0, 90^0. (42) $104^0\,28'\,39''$.

(43) -630^0. (44) $-\frac{1}{8}\sqrt{5}$. (46) $n\pi$; and $2n\pi \pm \frac{2}{3}\pi$.

(48) $\frac{1}{2}\{\sqrt{6} \pm \sqrt{2}\}$ and 15^0, 135^0; or, 105^0, 45^0.

(49) 9^0; $286^0 \,.\, 28' \,.\, 41{\cdot}16''$; $\frac{7}{5}$.

(52) (i) $n\pi \pm \frac{1}{3}\pi$. (ii) $\frac{1}{2}n\pi \pm \frac{1}{3}\pi$, or $n\pi + (-1)^n \frac{1}{2}\pi$.

(53) $37\frac{1}{2}$ sq. ft. (54) $40^0 \,.\, 29' \,.\, 19{\cdot}85''$.

(55) $\frac{1}{10^0 8^0 0}\pi$; $\frac{1}{1^0 0}\pi$; $\frac{5}{4}$. (56) $\tan\alpha = 4\sqrt{3}$, $\operatorname{cosec}\alpha = 1\frac{7}{2}\sqrt{3}$.

(58) (i) $n\pi \pm \frac{1}{4}\pi$. (ii) $\frac{1}{2}n\pi \pm \frac{1}{3}\pi$; or, $2n\pi \pm \frac{1}{4}\pi$.

(59) $2\sqrt{14}$ sq. ft. (60) $38^0 \,.\, 25' \,.\, 32{\cdot}725''$.

(61) 1·2 radians $= 76^0 39416^g$.

(64) The proper formula is $+\sqrt{(1+\sin A)} + \sqrt{(1-\sin A)}$.

(66) 192 ft., 185 ft. and 9234 sq. ft.

(67) 2·3 radians $= 131^0 779926^0$.

(69) The proper formula is $-\sqrt{(1+\sin A)} - \sqrt{(1-\sin A)}$.

(72) $78^0\,10'$, $70^0\,30'$, 9234 sq. ft.

(73) $\frac{4}{3^0 5^0}\pi$; $\frac{8}{3^0 5^0}\pi$; $\frac{14}{3^0 5^0}\pi$; $\frac{18}{3^0 5^0}\pi$; $\frac{24}{3^0 5^0}\pi$.

(76) $(2n \pm \frac{1}{2})\pi$, or $(2n \pm \frac{1}{3})\pi$. (78) 135^0, 15^0; or 45^0, 105^0.

(79) $\frac{2}{2^0 1^0}\pi$; $\frac{4}{2^0 1^0}\pi$; $\frac{6}{2^0 1^0}\pi$; $\frac{8}{2^0 1^0}\pi$; $\frac{1}{2}\frac{0}{1}\pi$; $\frac{1}{2}\frac{2}{1}\pi$. (81) $(2n \pm \frac{1}{3})\pi$.

(84) 7 ft.; $\sqrt{19}$ ft.; $1\frac{6}{2^0}\sqrt{3}$ sq. ft.

(88) 1035·43 ft.; 765·4321 ft.; 66^0. (89) 6·981 feet.

(90) $\frac{1}{2}$; $-\frac{1}{3}$. (91) $n\pi + (-1)^n \frac{5}{4}\pi$. (94) 321·0793.

(95) 13·751 ft. (96) $\frac{1}{1^0 3^0}$. (97) $2n\pi \pm \frac{2}{3}\pi$.

(99) $2\cos\frac{1}{2}A = +\sqrt{(1+\sin A)} - \sqrt{(1-\sin A)}$.

(102) $x = \frac{2}{5}r\pi \pm \frac{2}{15}\pi \pm \frac{1}{10}\pi$, $y = \frac{2}{5}r\pi \pm \frac{1}{5}\pi \pm \frac{1}{15}\pi$.

(104) $x = \frac{1}{2}$. (105) 11·7157 miles. (107) $n\pi + \frac{1}{3}\pi$; or, $n\pi + \frac{1}{4}\pi$.

(110) 25·7834 yds. (115) 1·219714 miles and 1 mile.

(116) $\sin = \frac{6}{65}$, $\cos = -\frac{16}{65}$, $\tan = -\frac{63}{16}$.

(127) $4\theta \pm \theta = 2n\pi \pm \frac{3}{4}\pi$; or, $\frac{1}{2}(2n+1)\pi$.

(134) $5 + \frac{1}{2}\log 2 + \frac{1}{2}L\sec A = 1 - \log 2 + \frac{2}{3}L\cos B + \frac{1}{3}L\operatorname{cosec}$.

(139) $m\pi$; or, $n\theta = r\pi + (-1)^r(\frac{1}{2}\pi - \theta)$.

(144) $\log 5 = 2a - c$, $\log 7 = c - a$, $\log 13 = b - 4a + 2c$.

(145) In $\frac{1}{4}(2\sqrt{2} \pm 1)$ hours. (151) 60^0, 45^0, 135^0, 120^0.

(156) 1. (157) 30^0 and 40^0.

(161) $\theta = n\pi$ or $2\theta = n\pi + \frac{1}{4}\pi$. (164) $\cos\theta = \frac{1}{4}(1 \pm \sqrt{5})$.

(168) $\sin 2\theta = 0$ or $\cos 2\theta = \frac{1}{3}$ or $-\frac{1}{2}$. (178) θ.

(182) $\dfrac{600}{13^2\sqrt{26}}$ and $\dfrac{5}{12}$. (191) $m^2 + p^2 = n^2 + q^2$.

(192) $-4\cos\dfrac{B+C-A}{2}\cos\dfrac{C+A-B}{2}\cos\dfrac{A+B-C}{2}\cos\dfrac{A+B+C}{2}$.

(196) $\frac{1}{4}\pi - \theta = n\pi + (-1)^n(\frac{1}{4}\pi - \alpha)$.

(198) $-4\cos\dfrac{90^0 - B - C + A}{2}\cos\dfrac{90^0 - C - A + B}{2}$
$\cos\dfrac{90^0 - A - B + C}{2}\sin\dfrac{A + B + C - 90^0}{2}$.

LXXVI b. Pages 269—275.

(10) $-(x+y+z)(y+z-x)(z+x-y)(x+y-z)$.

(11) (i) $\frac{1}{15}\{6n\pi - 2\pi + (-1)^n\pi\}$. (ii) $\dfrac{4r\pi + \pi}{2(m \pm n)}$.

(iii) $x = \dfrac{\alpha - \beta}{2} + \cot^{-1}\left\{\tan\dfrac{\alpha+\beta}{2}\cdot\dfrac{m\sin\beta + n\sin\alpha}{n\sin\alpha - m\sin\beta}\right\}$.

(iv) $\dfrac{2r\pi + \pi}{2(m+n)}$. (v) $\sin 2\theta(3\cos 2\theta - 1)(2\cos 2\theta + 1) = 0$.

(vi) $\sin 4\theta \cdot \cos\dfrac{5\theta}{2} \cdot \sin\dfrac{3\theta}{2} = 0$. (vii) $\sin 8\theta \cdot \sin 4\theta = 0$.

(21) 223·17.

CAMBRIDGE: PRINTED BY C. J. CLAY, M.A. AND SONS, AT THE UNIVERSITY PRESS.

WORKS BY THE SAME AUTHOR.

Fourth Edition Revised. Globe 8vo.

ARITHMETIC FOR SCHOOLS. Complete with Answers, 4*s.* 6*d.* Without Answers, 4*s.* 6*d.* Part I., with Answers, 2*s.*; Part II., with Answers, 3*s.* Fourth Edition with a new chapter on Commercial Discount.

KEY. Crown 8vo. 10*s.* 6*d.*

*** *The complete book and both parts can also be obtained* without answers *at the same price, though in different binding. But the edition* with answers *will always be supplied unless the other is specially asked for.*

ATHENÆUM:—"Mr Lock may certainly be congratulated on realizing the hope expressed in his preface, for his text-book is 'at once simple and scientific.'...Mr Lock shews that he is not only a proficient in arithmetic, but a master of the art of teaching it, and his work may be confidently recommended to both teachers and scholars."

CAMBRIDGE REVIEW:—"The best arithmetic for its purpose which we have seen."

LITERARY WORLD:—"It is the work of a thoroughly good teacher...Mr Lock's admirable *Trigonometry*, published last year, and the present work are, to our mind, models of what mathematical school-books should be."

Second Edition. Globe 8vo. 2*s.* 6*d.*

ARITHMETIC FOR BEGINNERS. A School-Class Book of Commercial Arithmetic.

KEY. By Rev. R. G. WATSON. Crown 8vo. 8*s.* 6*d.*

JOURNAL OF EDUCATION:—"This Arithmetic is founded on the author's larger work, such alterations having been made as were necessary to render it suitable for less advanced students, and in particular for those who are candidates for commercial certificates. The book is well adapted to the requirements of this new class of examinations."

PRACTICAL TEACHER:—"Will be useful in almost all schools, as it contains enough of theory and abundance of practice. The author is thoroughly master of his subject, he is a successful and experienced teacher, and he has the faculty of literary expression. Few writers of mathematical books combine these qualifications in an equal degree."

NATURE:—"A capital handbook."

18mo., Cloth, 1*s.* With answers, 1*s.* 6*d.*

A SHILLING CLASS-BOOK OF ARITHMETIC. Adapted for use in Elementary Schools.

GUARDIAN:—"This little book differs from most books of the class in that it is based on scientific principles."

SCHOOLMASTER:—"An excellent book of elementary arithmetic. The explanatory portion is pithy, but clear, and the examples plentiful, well graduated and fairly difficult."

WORKS BY THE SAME AUTHOR.

Just Published. Globe 8vo. *2s. 6d.*

THE TRIGONOMETRY OF ONE ANGLE.

*** *An easy introduction to Trigonometry containing all the Trigonometry necessary for the Previous Examination at Cambridge.*

Fourth Edition Revised and Enlarged. Globe 8vo. *2s. 6d.*

TRIGONOMETRY FOR BEGINNERS, as far as the Solution of Triangles.

KEY. Crown 8vo. *6s. 6d.*

SCHOOLMASTER:—"It is exactly the book to place in the hands of beginners.....Science teachers engaged in this particular branch of study will find the book most serviceable, while it will be equally useful to the private student."

NATURE:—"Mr Lock's great advantage is preceptorial skill in arrangement and exposition. On this score he deserves much credit indeed. There are few points in which it is possible to suggest improvement."

Fourth Edition. Globe 8vo. *4s. 6d.*

HIGHER TRIGONOMETRY. Both Parts in One Volume. Globe 8vo. *7s. 6d.*

*** *To this Edition a chapter on Antiparallels has been added.*

CAMBRIDGE REVIEW:—"This little book is obviously the work of one who has had considerable experience in teaching; it is written very clearly, the statements are definite and the proofs concise, and yet a teacher would not find it necessary to add much in the way of supplementary explanation; there is a copious supply of examples—both throughout the text and in the examination papers at the end—the solution is always so much more instructive than any amount of explanation. Many things generally taught here appear for the first time in a text-book, notably the ambiguity of logarithms (p. 29), and we welcome the appearance of the hyperbolic sine and cosine, whose introduction to common usage would save much labour, and make several systems of formulæ symmetrical and complete."

MACMILLAN AND CO., LONDON.

Messrs Macmillan & Co.'s Mathematical Books.

WORKS BY THE SAME AUTHOR.

Second Edition. Globe 8vo. 4*s.* 6*d.*

ELEMENTARY STATICS. A Companion Volume to "Dynamics for Beginners."

NATURE:—"Many treatises deal with elementary statics, but few can rival in clearness Mr Lock's work. The subject throughout is treated in the author's best style, and the book can be cordially recommended for the use of beginners....The working of examples, as every teacher knows, is the only way to obtain a thorough knowledge of any subject which requires mathematical treatment, and Mr Lock has fully recognized the importance of this. Typical examples, excellently selected, are worked out at full length, and numerous others are given as exercises. There is also a selection of papers from some of the Oxford and Cambridge examinations. A new departure is the introduction of a short chapter on graphic statics, which we highly approve of. The teaching of this subject has made rapid strides during the last few years, and the methods are so simple, and applicable in cases which would involve laborious calculations, that the introduction of the subject into text-books is very desirable. The whole subject is made interesting from beginning to end, and the proofs of the various propositions are very simple and clear. We have no doubt that the book will be appreciated by all who have an opportunity of judging of its merits."

EDUCATIONAL TIMES:—"It is marked by the same clearness and accuracy as the author's *Dynamics*, and affords further testimony of Mr Lock's ability both as a teacher and as a writer of text-books."

CAMBRIDGE REVIEW:—"We can, without hesitation, recommend this little work as a text-book for beginners."

IRON:—"This is a good little book, well printed, with clearly drawn illustrations, accurate in work, the demonstrations lucid and brief, and with a very large number of exercises...The book is likely to be a favourite school-book."

SCHOOLMASTER:—"This volume on statics has been as carefully prepared as any of the other scientific works bearing the name of Mr Lock. They all carry with them the stamp of thoroughness and experience, and this on statics is admirable, for its careful gradations and sensible arrangement and variety of problems to test one's knowledge of that subject."

GUARDIAN:—"In his 'Elementary Statics' Mr Lock has made a valuable addition to his useful series of mathematical class books...a special feature of that little book is an interesting chapter on 'Graphic Statics'."

Globe 8vo.

MECHANICS FOR BEGINNERS. Part I. THE MECHANICS OF SOLIDS. [*Ready in August* 1891.

MACMILLAN AND CO., LONDON.

Messrs Macmillan & Co.'s Mathematical Books.

WORKS BY THE SAME AUTHOR.

Fourth Edition. Globe 8vo. 4*s.* 6*d.*

ELEMENTARY DYNAMICS.

*** *To this Edition a chapter on the Pendulum has been added and many minor improvements effected.*

SATURDAY REVIEW:—"The parts that we have more carefully read we have found to be put with much freshness, and altogether the treatment is such as to make the subject interesting to an intelligent pupil."

ENGINEERING:—"This is beyond all doubt the most satisfactory treatise on Elementary Dynamics that has yet appeared."

NATURE:—"This book is an attempt to explain the Elementary principles of Dynamics in a manner suitable for school-work with boys of ordinary mathematical attainments. Accordingly it contains a great number of easy *numerical* examples, some worked out in illustration of the text, the others arranged in groups at frequent intervals. There is considerable freshness in these exercises, and they form altogether a very useful series...The exposition throughout is remarkable for clearness and precision of statement. The definitions of terms seem particularly well worded. The names *velo* and *celo* have been adopted for the units of velocity and acceleration, and are used systematically in both text and examples; we hope these terms may win their way to general acceptance, for the language of the subject gains both in simplicity and directness by their introduction... The debt of gratitude which many teachers and students already owe to Mr Lock will be considerably increased by this new class-book on a difficult subject, wherein it appears to us that the skill and experience of the author are displayed with great advantage."

Professor MINCHIN (*Professor of Applied Mathematics, Cooper's Hill*), says:—"I have used Mr Lock's Elementary Dynamics with those beginning the subject at Cooper's Hill this year, and I find the arrangement of the book—the order in which the different steps are placed before the student—to be exceedingly well adapted for the teaching of the subject. I have found also that the terms *velo* and *celo*, instead of exciting repugnance in the student's mind, were received (after provoking at first a little merriment) with sympathy, and, rapidly losing their strangeness, served as valuable and distinguishing marks of velocity and acceleration."

Globe 8vo.

EUCLID FOR BEGINNERS. Being an Introduction to existing Text-Books. [*In the Press.*

Globe 8vo.

ELEMENTARY HYDROSTATICS. A companion Volume to Elementary Statics and Elementary Dynamics.

[*In preparation.*

MACMILLAN AND CO., LONDON.

May 1891

A Catalogue

OF

Educational Books

PUBLISHED BY

Macmillan & Co.

BEDFORD STREET, STRAND, LONDON

CONTENTS

	PAGE
CLASSICS—	
Elementary Classics	2
Classical Series	4
Classical Library; Texts, Commentaries, Translations	6
Grammar, Composition, and Philology	9
Antiquities, Ancient History, and Philosophy	12
MODERN LANGUAGES AND LITERATURE—	
English	14
French	18
German	20
Modern Greek	21
Italian	21
Spanish	21
MATHEMATICS—	
Arithmetic	22
Book-Keeping	23
Algebra	23
Euclid and Pure Geometry	24
Geometrical Drawing	25
Mensuration	25
Trigonometry	25
Analytical Geometry	26
Problems and Questions in Mathematics	27
Higher Pure Mathematics	27
Mechanics	28
Physics	30
Astronomy	32
Historical	33
NATURAL SCIENCES—	
Chemistry	33
Physical Geography, Geology, and Mineralogy	35
Biology	36
Medicine	38
HUMAN SCIENCES—	
Mental and Moral Philosophy	39
Political Economy	41
Law and Politics	42
Anthropology	43
Education	43
TECHNICAL KNOWLEDGE—	
Civil and Mechanical Engineering	44
Military and Naval Science	44
Agriculture	45
Domestic Economy	46
Book-Keeping	46
GEOGRAPHY	46
HISTORY	47
ART	50
DIVINITY	51

CLASSICS.

Elementary Classics; Classical Series; Classical Library, (1) Texts, (2) Translations; Grammar, Composition, and Philology; Antiquities, Ancient History, and Philosophy.

*ELEMENTARY CLASSICS.

18mo, Eighteenpence each.

The following contain Introductions, Notes, **and Vocabularies**, and in some cases **Exercises.**

ACCIDENCE, LATIN, AND EXERCISES ARRANGED FOR BEGINNERS.—By W. WELCH, M.A., and C. G. DUFFIELD, M.A.

AESCHYLUS.—PROMETHEUS VINCTUS. By Rev. H. M. STEPHENSON, M.A.

ARRIAN.—SELECTIONS. With Exercises. By Rev. JOHN BOND, M.A., and Rev. A. S. WALPOLE, M.A.

AULUS GELLIUS, STORIES FROM.—Adapted for Beginners. With Exercises. By Rev. G. H. NALL, M.A., Assistant Master at Westminster.

CÆSAR.—THE HELVETIAN WAR. Being Selections from Book I. of The Gallic War. Adapted for Beginners. With Exercises. By W. WELCH, M.A., and C. G. DUFFIELD, M.A.

THE INVASION OF BRITAIN. Being Selections from Books IV. and V. of The Gallic War. Adapted for Beginners. With Exercises. By W. WELCH, M.A., and C. G. DUFFIELD, M.A.

SCENES FROM BOOKS V. AND VI. By C. COLBECK, M.A.

THE GALLIC WAR. BOOK I. By Rev. A. S. WALPOLE, M.A.

BOOKS II. AND III. By the Rev. W. G. RUTHERFORD, M.A., LL.D.

BOOK IV. By CLEMENT BRYANS, M.A., Assistant Master at Dulwich College.

BOOK V. By C. COLBECK, M.A., Assistant Master at Harrow.

BOOK VI. By the same Editor.

BOOK VII. By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.

THE CIVIL WAR. BOOK I. By M. MONTGOMERY, M.A. [*In preparation.*

CICERO.—DE SENECTUTE. By E. S. SHUCKBURGH, M.A.

DE AMICITIA. By the same Editor.

STORIES OF ROMAN HISTORY. Adapted for Beginners. With Exercises. By Rev. G. E. JEANS, M.A., and A. V. JONES, M.A.

EURIPIDES.—ALCESTIS. By Rev. M. A. BAYFIELD, M.A.

MEDEA. By A. W. VERRALL, Litt.D., and Rev. M. A. BAYFIELD, M.A. [*In the Press.*

HECUBA. By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.

EUTROPIUS.—Adapted for Beginners. With Exercises. By W. WELCH, M.A., and C. G. DUFFIELD, M.A.

HERODOTUS. TALES FROM HERODOTUS. Atticised by G. S. FARNELL, M.A. [*In the Press.*

HOMER.—ILIAD. BOOK I. By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.

BOOK XVIII. By S. R. JAMES, M.A., Assistant Master at Eton.

ODYSSEY. BOOK I. By Rev. J. BOND, M.A., and Rev. A. S. WALPOLE, M.A.

HORACE.—ODES. BOOKS I.-IV. By T. E. Page, M.A., Assistant Master at the Charterhouse. Each 1s. 6d.

LIVY.—BOOK I. By H. M. Stephenson, M.A.

BOOK XXI. Adapted from Mr. Capes's Edition. By J. E. Melhuish, M.A.

BOOK XXII. By the same.

THE HANNIBALIAN WAR. Being part of the XXI. and XXII. BOOKS OF LIVY adapted for Beginners. By G. C. Macaulay, M.A.

THE SIEGE OF SYRACUSE. Being part of the XXIV. and XXV. BOOKS OF LIVY, adapted for Beginners. With Exercises. By G. Richards, M.A., and Rev. A. S. Walpole, M.A.

LEGENDS OF ANCIENT ROME. Adapted for Beginners. With Exercises. By H. Wilkinson, M.A.

LUCIAN.—EXTRACTS FROM LUCIAN. With Exercises. By Rev. J. Bond, M.A., and Rev. A. S. Walpole, M.A.

NEPOS.—SELECTIONS ILLUSTRATIVE OF GREEK AND ROMAN HISTORY. With Exercises. By G. S. Farnell, M.A.

OVID.—SELECTIONS. By E. S. Shuckburgh, M.A.

EASY SELECTIONS FROM OVID IN ELEGIAC VERSE. With Exercises. By H. Wilkinson, M.A.

STORIES FROM THE METAMORPHOSES. With Exercises. By Rev. J. Bond, M.A., and Rev. A. S. Walpole, M.A.

PHÆDRUS.—SELECT FABLES. Adapted for Beginners. With Exercises. By Rev. A. S. Walpole, M.A.

THUCYDIDES.—THE RISE OF THE ATHENIAN EMPIRE. BOOK I. Chs. 89-117 and 228-238. With Exercises. By F. H. Colson, M.A.

VIRGIL.—SELECTIONS. By E. S. Shuckburgh, M.A.

BUCOLICS. By T. E. Page, M.A.

GEORGICS. BOOK I. By the same Editor.

BOOK II. By Rev. J. H. Skrine, M.A.

ÆNEID. BOOK I. By Rev. A. S. Walpole, M.A.

BOOK II. By T. E. Page, M.A.

BOOK III. By the same Editor.

BOOK IV. By Rev. H. M. Stephenson, M.A.

BOOK V. By Rev. A. Calvert, M.A.

BOOK VI. By T. E. Page, M.A.

BOOK VII. By Rev. A. Calvert, M.A.

BOOK VIII. By the same Editor.

BOOK IX. By Rev. H. M. Stephenson, M.A.

BOOK X. By S. G. Owen, M.A.

XENOPHON.—ANABASIS. Selections, adapted for Beginners. With Exercises. By W. Welch, M.A., and C. G. Duffield, M.A.

BOOK I. With Exercises. By E. A. Wells, M.A.

BOOK I. By Rev. A. S. Walpole, M.A.

BOOK II. By the same Editor.

BOOK III. By Rev. G. H. Nall, M.A.

BOOK IV. By Rev. E. D. Stone, M.A.

SELECTIONS FROM BOOK IV. With Exercises. By the same Editor.

SELECTIONS FROM THE CYROPÆDIA. With Exercises. By A. H. Cooke, M.A., Fellow and Lecturer of King's College, Cambridge.

The following contain Introductions and Notes, **but no Vocabulary :—**

CICERO.—SELECT LETTERS. By Rev. G. E. Jeans, M.A.

HERODOTUS.—SELECTIONS FROM BOOKS VII. and VIII. THE EXPEDITION OF XERXES. By A. H. Cooke, M.A.

HORACE.—SELECTIONS FROM THE SATIRES AND EPISTLES. By Rev. W. J. V. Baker, M.A.

SELECT EPODES AND ARS POETICA. By H. A. Dalton, M.A., Assistant Master at Winchester.

PLATO.—EUTHYPHRO AND MENEXENUS. By C. E. Graves, M.A.

TERENCE.—SCENES FROM THE ANDRIA. By F. W. Cornish, M.A., Assistant Master at Eton.

THE GREEK ELEGIAC POETS.—FROM CALLINUS TO CALLIMACHUS. Selected by Rev. Herbert Kynaston, D.D.

THUCYDIDES.—BOOK IV. Chs. 1-41. THE CAPTURE OF SPHACTERIA. By C. E. Graves, M.A.

CLASSICAL SERIES FOR COLLEGES AND SCHOOLS.

Fcap. 8vo.

ÆSCHINES.—IN CTESIPHONTA. By Rev. T. Gwatkin, M.A., and E. S. Shuckburgh, M.A. 5s.

ÆSCHYLUS.—PERSÆ. By A. O. Prickard, M.A., Fellow and Tutor of New College, Oxford. With Map. 2s. 6d.

SEVEN AGAINST THEBES. SCHOOL EDITION. By A. W. Verrall, Litt.D., Fellow of Trinity College, Cambridge, and M. A. Bayfield, M.A., Headmaster of Christ's College, Brecon. 2s. 6d.

ANDOCIDES.—DE MYSTERIIS. By W. J. Hickie, M.A. 2s. 6d.

ATTIC ORATORS.—Selections from ANTIPHON, ANDOCIDES, LYSIAS, ISOCRATES, and ISAEUS. By R. C. Jebb, Litt.D., Regius Professor of Greek in the University of Cambridge. 5s.

***CÆSAR.**—THE GALLIC WAR. By Rev. John Bond, M.A., and Rev. A. S. Walpole, M.A. With Maps. 4s. 6d.

CATULLUS.—SELECT POEMS. Edited by F. P. Simpson, B.A. 3s. 6d. The Text of this Edition is carefully expurgated for School use.

***CICERO.**—THE CATILINE ORATIONS. By A. S. Wilkins, Litt.D., Professor of Latin in the Owens College, Victoria University, Manchester. 2s. 6d.

PRO LEGE MANILIA. By Prof. A. S. Wilkins, Litt.D. 2s. 6d.

THE SECOND PHILIPPIC ORATION. By John E. B. Mayor, M.A., Professor of Latin in the University of Cambridge. 3s. 6d.

PRO ROSCIO AMERINO. By E. H. Donkin, M.A. 2s. 6d.

PRO P. SESTIO. By Rev. H. A. Holden, Litt.D. 3s. 6d.

SELECT LETTERS. Edited by R. Y. Tyrrell, M.A. [*In the Press.*

DEMOSTHENES.—DE CORONA. By B. Drake, M.A. 7th Edition, revised by E. S. Shuckburgh, M.A. 3s. 6d.

ADVERSUS LEPTINEM. By Rev. J. R. King, M.A., Fellow and Tutor of Oriel College, Oxford. 2s. 6d.

THE FIRST PHILIPPIC. By Rev. T. Gwatkin, M.A. 2s. 6d.

IN MIDIAM. By Prof. A. S. Wilkins, Litt.D., and Herman Hager, Ph.D., of the Owens College, Victoria University, Manchester. [*In preparation.*

EURIPIDES.—HIPPOLYTUS. By Rev. J. P. Mahaffy, D.D., Fellow of Trinity College, and Professor of Ancient History in the University of Dublin, and J. B. Bury, M.A., Fellow of Trinity College, Dublin. 2s. 6d.

MEDEA. By A. W. Verrall, Litt.D., Fellow of Trinity College, Cambridge. 2s. 6d.

IPHIGENIA IN TAURIS. By E. B. England, M.A. 3s.

ION. By M. A. Bayfield, M.A., Headmaster of Christ's College, Brecon. 2s. 6d.

BACCHAE. By R. Y. Tyrrell, M.A., Regius Professor of Greek in the University of Dublin. [*In preparation.*

HERODOTUS.—BOOK III. By G. C. Macaulay, M.A. 2s. 6d.

BOOK V. By J. Strachan, M.A., Professor of Greek in the Owens College, Victoria University, Manchester. [*In preparation.*

BOOK VI. By the same. 3s. 6d.

BOOK VII. By Mrs. A. F. Butler. 3s. 6d.

HESIOD.—THE WORKS AND DAYS. By W. T. Lendrum, M.A., Assistant Master at Dulwich College. [*In preparation.*

HOMER.—ILIAD. BOOKS I., IX., XI., XVI.-XXIV. THE STORY OF ACHILLES. By the late J. H. PRATT, M.A., and WALTER LEAF, Litt.D., Fellows of Trinity College, Cambridge. 5s.

ODYSSEY. BOOK IX. By Prof. JOHN E. B. MAYOR. 2s. 6d.

ODYSSEY. BOOKS XXI.-XXIV. THE TRIUMPH OF ODYSSEUS. By S. G. HAMILTON, B.A., Fellow of Hertford College, Oxford. 2s. 6d.

HORACE.—*THE ODES. By T. E. PAGE, M.A., Assistant Master at the Charterhouse. 5s. (BOOKS I., II., III., and IV. separately, 2s. each.)

THE SATIRES. By ARTHUR PALMER, M.A., Professor of Latin in the University of Dublin. 5s.

THE EPISTLES AND ARS POETICA. By A. S. WILKINS, Litt.D., Professor of Latin in the Owens College, Victoria University, Manchester. 5s.

ISAEOS.—THE ORATIONS. By WILLIAM RIDGEWAY, M.A., Professor of Greek in Queen's College, Cork. [*In preparation.*

JUVENAL.—*THIRTEEN SATIRES. By E. G. HARDY, M.A. 5s. The Text is carefully expurgated for School use.

SELECT SATIRES. By Prof. JOHN E. B. MAYOR. X. and XI. 3s. 6d. XII.-XVI. 4s. 6d.

LIVY.—*BOOKS II. and III. By Rev. H. M. STEPHENSON, M.A. 3s. 6d.

*BOOKS XXI. and XXII. By Rev. W. W. CAPES, M.A. With Maps. 4s. 6d.

*BOOKS XXIII. and XXIV. By G. C. MACAULAY, M.A. With Maps. 3s. 6d.

*THE LAST TWO KINGS OF MACEDON. EXTRACTS FROM THE FOURTH AND FIFTH DECADES OF LIVY. By F. H. RAWLINS, M.A., Assistant Master at Eton. With Maps. 2s. 6d.

THE SUBJUGATION OF ITALY. SELECTIONS FROM THE FIRST DECADE. By G. E. MARINDIN, M.A. [*In preparation.*

LUCRETIUS.—BOOKS I.-III. By J. H. WARBURTON LEE, M.A., Assistant Master at Rossall. 3s. 6d.

LYSIAS.—SELECT ORATIONS. By E. S. SHUCKBURGH, M.A. 5s.

MARTIAL.—SELECT EPIGRAMS. By Rev. H. M. STEPHENSON, M.A. 5s.

***OVID.**—FASTI. By G. H. HALLAM, M.A., Assistant Master at Harrow. With Maps. 3s. 6d.

*HEROIDUM EPISTULÆ XIII. By E. S. SHUCKBURGH, M.A. 3s. 6d.

METAMORPHOSES. BOOKS I.-III. By C. SIMMONS, M.A. [*In preparation.*

BOOKS XIII. and XIV. By the same Editor. 3s. 6d.

PLATO.—LACHES. By M. T. TATHAM, M.A. 2s. 6d.

THE REPUBLIC. BOOKS I.-V. By T. H. WARREN, M.A., President of Magdalen College, Oxford. 5s.

PLAUTUS.—MILES GLORIOSUS. By R. Y. TYRRELL, M.A., Regius Professor of Greek in the University of Dublin. 2d Ed., revised. 3s. 6d.

*AMPHITRUO. By ARTHUR PALMER, M.A., Professor of Latin in the University of Dublin. 3s. 6d.

CAPTIVI. By A. RHYS-SMITH, M.A. [*In the Press.*

PLINY.—LETTERS. BOOKS I. and II. By J. COWAN, M.A., Assistant Master at the Manchester Grammar School. 3s.

LETTERS. BOOK III. By Prof. JOHN E. B. MAYOR. With Life of Pliny by G. H. RENDALL, M.A. 3s. 6d.

PLUTARCH.—LIFE OF THEMISTOKLES. By Rev. H. A. HOLDEN, Litt.D. 3s. 6d.

LIVES OF GALBA AND OTHO. By E. G. HARDY, M.A. 5s.

POLYBIUS.—THE HISTORY OF THE ACHÆAN LEAGUE AS CONTAINED IN THE REMAINS OF POLYBIUS. By W. W. CAPES, M.A. 5s.

PROPERTIUS.—SELECT POEMS. By Prof. J. P. POSTGATE, Litt.D., Fellow of Trinity College, Cambridge. 2d Ed., revised. 5s.

SALLUST.—*CATILINA and JUGURTHA. By C. MERIVALE, D.D., Dean of Ely. 3s. 6d. Or separately, 2s. each.

*BELLUM CATULINÆ. By A. M. COOK, M.A., Assistant Master at St. Paul's School. 2s. 6d.

JUGURTHA. By the same Editor. [*In preparation.*

TACITUS.—THE ANNALS. BOOKS I. and II. By J. S. Reid, Litt.D. [*In prep.*
THE ANNALS. BOOK VI. By A. J. Church, M.A., and W. J. Brodribb, M.A. 2s.
THE HISTORIES. BOOKS I. and II. By A. D. Godley, M.A., Fellow of Magdalen College, Oxford. 3s. 6d. BOOKS III.-V. By the same. 3s. 6d.
AGRICOLA and GERMANIA. By A. J. Church, M.A., and W. J. Brodribb, M.A. 3s. 6d. Or separately, 2s. each.

TERENCE.—HAUTON TIMORUMENOS. By E. S. Shuckburgh, M.A. 2s. 6d. With Translation. 3s. 6d.
PHORMIO. By Rev. John Bond, M.A., and Rev. A. S. Walpole, M.A. 2s. 6d.

THUCYDIDES.—BOOK I. By C. Bryans, M.A. [*In preparation.*
BOOK II. By E. C. Marchant, M.A., Assistant Master at St. Paul's.
BOOK III. By C. Bryans, M.A. [*In preparation.*
BOOK IV. By C. E. Graves, M.A., Classical Lecturer at St. John's College, Cambridge. 3s. 6d.
BOOK V. By the same Editor. [*In the Press.*
BOOKS VI. and VII. THE SICILIAN EXPEDITION. By Rev. Percival Frost, M.A. With Map. 3s. 6d.
BOOK VIII. By Prof. T. G. Tucker, M.A. [*In preparation.*

TIBULLUS.—SELECT POEMS. By Prof. J. P. Postgate, Litt.D. [*In preparation.*

VIRGIL.—ÆNEID. BOOKS II. and III. THE NARRATIVE OF ÆNEAS. By E. W. Howson, M.A., Assistant Master at Harrow. 2s.

XENOPHON.—*THE ANABASIS. BOOKS I.-IV. By Profs. W. W. Goodwin and J. W. White. Adapted to Goodwin's Greek Grammar. With Map. 3s. 6d.
HELLENICA. BOOKS I. and II. By H. Hailstone, B.A. With Map. 2s. 6d.
CYROPÆDIA. BOOKS VII. and VIII. By A. Goodwin, M.A., Professor of Classics in University College, London. 2s. 6d.
MEMORABILIA SOCRATIS. By A. R. Cluer, B.A., Balliol College, Oxford. 5s.
HIERO. By Rev. H. A. Holden, Litt.D., LL.D. 2s. 6d.
OECONOMICUS. By the same. With Lexicon. 5s.

CLASSICAL LIBRARY.

Texts, Edited with **Introductions and Notes,** for the use of Advanced Students ; **Commentaries and Translations.**

ÆSCHYLUS.—THE SUPPLICES. A Revised Text, with Translation. By T. G. Tucker, M.A., Professor of Classical Philology in the University of Melbourne. 8vo. 10s. 6d.
THE SEVEN AGAINST THEBES. With Translation. By A. W. Verrall, Litt.D., Fellow of Trinity College, Cambridge. 8vo. 7s. 6d.
AGAMEMNON. With Translation. By A. W. Verrall, Litt.D. 8vo. 12s.
AGAMEMNON, CHOEPHORŒ, AND EUMENIDES. By A. O. Prickard, M.A., Fellow and Tutor of New College, Oxford. 8vo. [*In preparation.*
THE EUMENIDES. With Verse Translation. By Bernard Drake, M.A. 8vo. 5s.

ANTONINUS, MARCUS AURELIUS.—BOOK IV. OF THE MEDITATIONS. With Translation. By Hastings Crossley, M.A. 8vo. 6s.

ARISTOTLE.—THE METAPHYSICS. BOOK I. Translated by a Cambridge Graduate. 8vo. 5s.
THE POLITICS. By R. D. Hicks, M.A., Fellow of Trinity College, Cambridge. 8vo. [*In the Press.*
THE POLITICS. Translated by Rev. J. E. C. Welldon, M.A., Headmaster of Harrow. Cr. 8vo. 10s. 6d.
THE RHETORIC. Translated by the same. Cr. 8vo. 7s. 6d.
AN INTRODUCTION TO ARISTOTLE'S RHETORIC. With Analysis, Notes, and Appendices. By E. M. Cope, Fellow and late Tutor of Trinity College, Cambridge. 8vo. 14s.

THE ETHICS. Translated by Rev. J. E. C. Welldon, M.A. Cr. 8vo. [*In prep.*

THE SOPHISTICI ELENCHI. With Translation. By E. Poste, M.A., Fellow of Oriel College, Oxford. 8vo. 8s. 6d.

ON THE CONSTITUTION OF ATHENS. Edited by J. E. Sandys, Litt.D.

ARISTOPHANES.—THE BIRDS. Translated into English Verse. By B. H. Kennedy, D.D. Cr. 8vo. 6s. Help Notes to the Same, for the Use of Students. 1s. 6d.

ATTIC ORATORS.—FROM ANTIPHON TO ISAEOS. By R. C. Jebb, Litt.D., Regius Professor of Greek in the University of Cambridge. 2 vols. 8vo. 25s.

BABRIUS.—With Lexicon. By Rev. W. G. Rutherford, M.A., LL.D., Headmaster of Westminster. 8vo. 12s. 6d.

CICERO.—THE ACADEMICA. By J. S. Reid, Litt.D., Fellow of Caius College, Cambridge. 8vo. 15s.

THE ACADEMICS. Translated by the same. 8vo. 5s. 6d.

SELECT LETTERS. After the Edition of Albert Watson, M.A. Translated by G. E. Jeans, M.A., Fellow of Hertford College, Oxford. Cr. 8vo. 10s. 6d.

EURIPIDES.—MEDEA. Edited by A. W. Verrall, Litt.D. 8vo. 7s. 6d.

IPHIGENEIA IN AULIS. Edited by E. B. England, M.A. 8vo. [*In the Press.*

***INTRODUCTION TO THE STUDY OF EURIPIDES.** By Professor J. P. Mahaffy. Fcap. 8vo. 1s. 6d. (*Classical Writers.*)

HERODOTUS.—BOOKS I.–III. THE ANCIENT EMPIRES OF THE EAST. Edited by A. H. Sayce, Deputy-Professor of Comparative Philology, Oxford. 8vo. 16s.

BOOKS IV.–IX. Edited by R. W. Macan, M.A., Reader in Ancient History in the University of Oxford. 8vo. [*In preparation.*

THE HISTORY. Translated by G. C. Macaulay, M.A. 2 vols. Cr. 8vo. 18s.

HOMER.—THE ILIAD. By Walter Leaf, Litt.D. 8vo. Books I.–XII. 14s. Books XIII.–XXIV. 14s.

THE ILIAD. Translated into English Prose by Andrew Lang, M.A., Walter Leaf, Litt.D., and Ernest Myers, M.A. Cr. 8vo. 12s. 6d.

THE ODYSSEY. Done into English by S. H. Butcher, M.A., Professor of Greek in the University of Edinburgh, and Andrew Lang, M.A. Cr. 8vo. 6s.

***INTRODUCTION TO THE STUDY OF HOMER.** By the Right Hon. W. E. Gladstone. 18mo. 1s. (*Literature Primers.*)

HOMERIC DICTIONARY. Translated from the German of Dr. G. Autenrieth by R. P. Keep, Ph.D. Illustrated. Cr. 8vo. 6s.

HORACE.—Translated by J. Lonsdale, M.A., and S. Lee, M.A. Gl. 8vo. 3s. 6d.

STUDIES, LITERARY AND HISTORICAL, IN THE ODES OF HORACE. By A. W. Verrall, Litt.D. 8vo. 8s. 6d.

JUVENAL.—THIRTEEN SATIRES OF JUVENAL. By John E. B. Mayor, M.A., Professor of Latin in the University of Cambridge. Cr. 8vo. 2 vols. 10s. 6d. each. Vol. I. 10s. 6d. Vol. II. 10s. 6d.

THIRTEEN SATIRES. Translated by Alex. Leeper, M.A., LL.D., Warden of Trinity College, Melbourne. Cr. 8vo. 3s. 6d.

KTESIAS.—THE FRAGMENTS OF THE PERSIKA OF KTESIAS. By John Gilmore, M.A. 8vo. 8s. 6d.

LIVY.—BOOKS I.–IV. Translated by Rev. H. M. Stephenson, M.A. [*In prep.*

BOOKS XXI.–XXV. Translated by A. J. Church, M.A., and W. J. Brodribb, M.A. Cr. 8vo. 7s. 6d.

***INTRODUCTION TO THE STUDY OF LIVY.** By Rev. W. W. Capes, M.A. Fcap. 8vo. 1s. 6d. (*Classical Writers.*)

LONGINUS.—ON THE SUBLIME. Translated by H. L. Havell, B.A. With Introduction by Andrew Lang. Cr. 8vo. 4s. 6d.

MARTIAL.—BOOKS I. and II. OF THE EPIGRAMS. By Prof. John E. B. Mayor, M.A. 8vo. [*In the Press.*

MELEAGER.—FIFTY POEMS OF MELEAGER. Translated by Walter Headlam. Fcap. 4to. 7s. 6d.

PAUSANIAS.—DESCRIPTION OF GREECE. Translated with Commentary by J. G. FRAZER, M.A., Fellow of Trinity College, Cambridge. [*In prep.*

PHRYNICHUS.—THE NEW PHRYNICHUS; being a Revised Text of the Ecloga of the Grammarian Phrynichus. With Introduction and Commentary by Rev. W. G. RUTHERFORD, M.A., LL.D., Headmaster of Westminster. 8vo. 18s.

PINDAR.—THE EXTANT ODES OF PINDAR. Translated by ERNEST MYERS, M.A. Cr. 8vo. 5s.

THE OLYMPIAN AND PYTHIAN ODES. Edited, with an Introductory Essay, by BASIL GILDERSLEEVE, Professor of Greek in the Johns Hopkins University, U.S.A. Cr. 8vo. 7s. 6d.

THE NEMEAN ODES. By J. B. BURY, M.A., Fellow of Trinity College, Dublin. 8vo. 12s.

PLATO.—PHÆDO. By R. D. ARCHER-HIND, M.A., Fellow of Trinity College, Cambridge. 8vo. 8s. 6d.

PHÆDO. By W. D. GEDDES, LL.D., Principal of the University of Aberdeen. 8vo. 8s. 6d.

TIMAEUS. With Translation. By R. D. ARCHER-HIND, M.A. 8vo. 16s.

THE REPUBLIC OF PLATO. Translated by J. LL. DAVIES, M.A., and D. J. VAUGHAN, M.A. 18mo. 4s. 6d.

EUTHYPHRO, APOLOGY, CRITO, AND PHÆDO. Translated by F. J. CHURCH. 18mo. 4s. 6d.

PHÆDRUS, LYSIS, AND PROTAGORAS. Translated by J. WRIGHT, M.A. 18mo. 4s. 6d.

PLAUTUS.—THE MOSTELLARIA. By WILLIAM RAMSAY, M.A. Edited by G. G. RAMSAY, M.A., Professor of Humanity in the University of Glasgow. 8vo. 14s.

PLINY.—CORRESPONDENCE WITH TRAJAN. C. Plinii Caecilii Secundi Epistulæ ad Traianum Imperatorem cum Eiusdem Responsis. By E. G. HARDY, M.A. 8vo. 10s. 6d.

POLYBIUS.—THE HISTORIES OF POLYBIUS. Translated by E. S. SHUCKBURGH, M.A. 2 vols. Cr. 8vo. 24s.

SALLUST.—CATILINE AND JUGURTHA. Translated by A. W. POLLARD, B.A. Cr. 8vo. 6s. THE CATILINE (separately). 3s.

SOPHOCLES.—ŒDIPUS THE KING. Translated into English Verse by E. D. A. MORSHEAD, M.A., Assistant Master at Winchester. Fcap. 8vo. 3s. 6d.

TACITUS.—THE ANNALS. By G. O. HOLBROOKE, M.A., Professor of Latin in Trinity College, Hartford, U.S.A. With Maps. 8vo. 16s.

THE ANNALS. Translated by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Maps. Cr. 8vo. 7s. 6d.

THE HISTORIES. By Rev. W. A. SPOONER, M.A., Fellow and Tutor of New College, Oxford. 8vo. 16s.

THE HISTORY. Translated by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Map. Cr. 8vo. 6s.

THE AGRICOLA AND GERMANY, WITH THE DIALOGUE ON ORATORY. Translated by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Maps. Cr. 8vo. 4s. 6d.

*INTRODUCTION TO THE STUDY OF TACITUS. By A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. Fcap. 8vo. 1s. 6d. (*Classical Writers.*)

THEOCRITUS, BION, AND MOSCHUS. Translated by A. LANG, M.A. 18mo. 4s. 6d. Also an Edition on Large Paper. Cr. 8vo. 9s.

THUCYDIDES.—BOOK IV. A Revision of the Text, Illustrating the Principal Causes of Corruption in the Manuscripts of this Author. By Rev. W. G. RUTHERFORD, M.A., LL.D., Headmaster of Westminster. 8vo. 7s. 6d.

BOOK VIII. By H. C. GOODHART, M.A., Fellow of Trinity College, Cambridge. [*In the Press.*

VIRGIL.—Translated by J. LONSDALE, M.A., and S. LEE, M.A. Gl. 8vo. 3s. 6d.

THE ÆNEID. Translated by J. W. MACKAIL, M.A., Fellow of Balliol College, Oxford. Cr. 8vo. 7s. 6d.

XENOPHON.—Translated by H. G. Dakyns, M.A. In four vols. Cr. 8vo. Vol. I., containing "The Anabasis" and Books I. and II. of "The Hellenica." 10s. 6d. Vol. II. "Hellenica" III.–VII., and the two Polities—"Athenian" and "Laconian," the "Agesilaus," and the tract on "Revenues." With Maps and Plans. [*In the Press.*

GRAMMAR, COMPOSITION, & PHILOLOGY.

***BELCHER.**—SHORT EXERCISES IN LATIN PROSE COMPOSITION AND EXAMINATION PAPERS IN LATIN GRAMMAR. Part I. By Rev. H. Belcher, LL.D., Rector of the High School, Dunedin, N.Z. 18mo. 1s. 6d.
KEY, for Teachers only. 18mo. 3s. 6d.
*Part II., On the Syntax of Sentences, with an Appendix, including EXERCISES IN LATIN IDIOMS, etc. 18mo. 2s. KEY, for Teachers only. 18mo. 3s.

BLACKIE.—GREEK AND ENGLISH DIALOGUES FOR USE IN SCHOOLS AND COLLEGES. By John Stuart Blackie, Emeritus Professor of Greek in the University of Edinburgh. New Edition. Fcap. 8vo. 2s. 6d.
A GREEK PRIMER, COLLOQUIAL AND CONSTRUCTIVE. Globe 8vo.

***BRYANS.**—LATIN PROSE EXERCISES BASED UPON CÆSAR'S GALLIC WAR. With a Classification of Cæsar's Chief Phrases and Grammatical Notes on Cæsar's Usages. By Clement Bryans, M.A., Assistant Master at Dulwich College. Ex. fcap. 8vo. 2s. 6d. KEY, for Teachers only. 4s. 6d.
GREEK PROSE EXERCISES based upon Thucydides. By the same. [*In preparation.*

COOKSON.—A LATIN SYNTAX. By Christopher Cookson, M.A., Assistant Master at St. Paul's School. 8vo. [*In preparation.*

CORNELL UNIVERSITY STUDIES IN CLASSICAL PHILOLOGY. Edited by I. Flagg, W. G. Hale, and B. I. Wheeler. I. The *CUM*-Constructions: their History and Functions. By W. G. Hale. Part 1. Critical. 1s. 8d. net. Part 2. Constructive. 3s. 4d. net. II. Analogy and the Scope of its Application in Language. By B. I. Wheeler. 1s. 3d. net.

***EICKE.**—FIRST LESSONS IN LATIN. By K. M. Eicke, B.A., Assistant Master at Oundle School. Gl. 8vo. 2s. 6d.

***ENGLAND.**—EXERCISES ON LATIN SYNTAX AND IDIOM. ARRANGED WITH REFERENCE TO ROBY'S SCHOOL LATIN GRAMMAR. By E. B. England, Assistant Lecturer at the Owens College, Victoria University, Manchester. Cr. 8vo. 2s. 6d. KEY, for Teachers only. 2s. 6d.

GILES.—A SHORT MANUAL OF PHILOLOGY FOR CLASSICAL STUDENTS. By P. Giles, M.A., Fellow of Gonville and Caius College, Cambridge. Cr. 8vo. [*In the Press.*

GOODWIN.—Works by W. W. Goodwin, LL.D., D.C.L., Professor of Greek in Harvard University, U.S.A.
SYNTAX OF THE MOODS AND TENSES OF THE GREEK VERB. New Ed., revised and enlarged. 8vo. 14s.
*A GREEK GRAMMAR. Cr. 8vo. 6s.
*A GREEK GRAMMAR FOR SCHOOLS. Cr. 8vo. 3s. 6d.

GREENWOOD.—THE ELEMENTS OF GREEK GRAMMAR. Adapted to the System of Crude Forms. By J. G. Greenwood, sometime Principal of the Owens College, Manchester. Cr. 8vo. 5s. 6d.

HADLEY.—ESSAYS, PHILOLOGICAL AND CRITICAL. By James Hadley, late Professor in Yale College. 8vo. 16s.

HADLEY AND ALLEN.—A GREEK GRAMMAR FOR SCHOOLS AND COLLEGES. By James Hadley, late Professor in Yale College. Revised and in part rewritten by F. de F. Allen, Professor in Harvard College. Cr. 8vo. 6s.

HODGSON.—MYTHOLOGY FOR LATIN VERSIFICATION. A brief sketch of the Fables of the Ancients, prepared to be rendered into Latin Verse for Schools. By F. Hodgson, B.D., late Provost of Eton. New Ed., revised by F. C. Hodgson, M.A. 18mo. 3s.

***JACKSON.**—FIRST STEPS TO GREEK PROSE COMPOSITION. By Blomfield Jackson, M.A., Assistant Master at King's College School. 18mo. 1s. 6d.
KEY, for Teachers only. 18mo. 3s. 6d.

*SECOND STEPS TO GREEK PROSE COMPOSITION, with Miscellaneous Idioms, Aids to Accentuation, and Examination Papers in Greek Scholarship. By the same. 18mo. 2s. 6d. KEY, for Teachers only. 18mo. 3s. 6d.

KYNASTON.—EXERCISES IN THE COMPOSITION OF GREEK IAMBIC VERSE by Translations from English Dramatists. By Rev. H. KYNASTON, D.D., Professor of Classics in the University of Durham. With Vocabulary. Ex. fcap. 8vo. 5s.

KEY, for Teachers only. Ex. fcap. 8vo. 4s. 6d.

LUPTON.—*AN INTRODUCTION TO LATIN ELEGIAC VERSE COMPOSITION. By J. H. LUPTON, Sur-Master of St. Paul's School. Gl. 8vo. 2s. 6d. KEY TO PART II. (XXV.–C.) Gl. 8vo. 3s. 6d.

*AN INTRODUCTION TO LATIN LYRIC VERSE COMPOSITION. By the same. Gl. 8vo. 3s. KEY, for Teachers only. Gl. 8vo. 4s. 6d.

MACKIE.—PARALLEL PASSAGES FOR TRANSLATION INTO GREEK AND ENGLISH. With Indexes. By Rev. ELLIS C. MACKIE, M.A., Classical Master at Heversham Grammar School. Gl. 8vo. 4s. 6d.

*MACMILLAN.—FIRST LATIN GRAMMAR. By M. C. MACMILLAN, M.A Fcap. 8vo. 1s. 6d.

MACMILLAN'S GREEK COURSE.—Edited by Rev. W. G. RUTHERFORD, M.A., LL.D., Headmaster of Westminster. Gl. 8vo.

*FIRST GREEK GRAMMAR—ACCIDENCE. By the Editor. 2s.

*FIRST GREEK GRAMMAR—SYNTAX. By the same. 2s.

ACCIDENCE AND SYNTAX. In one volume. 3s. 6d.

*EASY EXERCISES IN GREEK ACCIDENCE. By H. G. UNDERHILL, M.A., Assistant Master at St. Paul's Preparatory School. 2s.

*A SECOND GREEK EXERCISE BOOK. By Rev. W. A. HEARD, M.A., Headmaster of Fettes College, Edinburgh. 2s. 6d.

EASY EXERCISES IN GREEK SYNTAX. By Rev. G. H. NALL, M.A., Assistant Master at Westminster School. [*In preparation.*

MANUAL OF GREEK ACCIDENCE. By the Editor. [*In preparation.*

MANUAL OF GREEK SYNTAX. By the Editor. [*In preparation.*

ELEMENTARY GREEK COMPOSITION. By the Editor. [*In preparation.*

*MACMILLAN'S GREEK READER.—STORIES AND LEGENDS. A First Greek Reader, with Notes, Vocabulary, and Exercises. By F. H. COLSON, M.A., Headmaster of Plymouth College. Gl. 8vo. 3s.

MACMILLAN'S LATIN COURSE.—By A. M. COOK, M.A., Assistant Master at St. Paul's School.

*FIRST PART. Gl. 8vo. 3s. 6d.

*SECOND PART. 2s. 6d. [*Third Part in preparation.*

*MACMILLAN'S SHORTER LATIN COURSE.—By A. M. COOK, M.A. Being an abridgment of "Macmillan's Latin Course," First Part. Gl. 8vo. 1s. 6d.

*MACMILLAN'S LATIN READER.—A LATIN READER FOR THE LOWER FORMS IN SCHOOLS. By H. J. HARDY, M.A., Assistant Master at Winchester. Gl. 8vo. 2s. 6d.

*MARSHALL.—A TABLE OF IRREGULAR GREEK VERBS, classified according to the arrangement of Curtius's Greek Grammar. By J. M. MARSHALL, M.A., Headmaster of the Grammar School, Durham. 8vo. 1s.

MAYOR.—FIRST GREEK READER. By Prof. JOHN E. B. MAYOR, M.A., Fellow of St. John's College, Cambridge. Fcap. 8vo. 4s. 6d.

MAYOR.—GREEK FOR BEGINNERS. By Rev. J. B. MAYOR, M.A., late Professor of Classical Literature in King's College, London. Part I., with Vocabulary, 1s. 6d. Parts II. and III., with Vocabulary and Index. Fcap. 8vo. 3s. 6d. Complete in one Vol. 4s. 6d.

NIXON.—PARALLEL EXTRACTS, Arranged for Translation into English and Latin, with Notes on Idioms. By J. E. NIXON, M.A., Fellow and Classical Lecturer, King's College, Cambridge. Part I.—Historical and Epistolary. Cr. 8vo. 3s. 6d.

PROSE EXTRACTS, Arranged for Translation into English and Latin, with General and Special Prefaces on Style and Idiom. By the same. I. Oratorical. II. Historical. III. Philosophical. IV. Anecdotes and Letters. 2d Ed., enlarged to 280 pp. Cr. 8vo. 4s. 6d. SELECTIONS FROM THE SAME. 3s. Translations of about 70 Extracts can be supplied to Schoolmasters (2s. 6d.), on application to the Author: and about 40 similarly of "Parallel Extracts," 1s. 6d. post free.

*PANTIN.—A FIRST LATIN VERSE BOOK. By W. E. P. Pantin, M.A., Assistant Master at St. Paul's School. Gl. 8vo. 1s. 6d.

*PEILE.—A PRIMER OF PHILOLOGY. By J. Peile, Litt.D., Master of Christ's College, Cambridge. 18mo. 1s.

*POSTGATE.—SERMO LATINUS. A short Guide to Latin Prose Composition. By Prof. J. P. Postgate, Litt.D., Fellow of Trinity College, Cambridge. Gl. 8vo. 2s. 6d. KEY to "Selected Passages." Gl. 8vo. 3s. 6d.

POSTGATE and VINCE.—A DICTIONARY OF LATIN ETYMOLOGY. By J. P. Postgate and C. A. Vince. [*In preparation.*

POTTS.—*HINTS TOWARDS LATIN PROSE COMPOSITION. By A. W. Potts, M.A., LL.D., late Fellow of St. John's College, Cambridge. Ex. fcap. 8vo. 3s.

*PASSAGES FOR TRANSLATION INTO LATIN PROSE. Edited with Notes and References to the above. Ex. fcap. 8vo. 2s. 6d. KEY, for Teachers only. 2s. 6d.

*PRESTON.—EXERCISES IN LATIN VERSE OF VARIOUS KINDS. By Rev. G. Preston. Gl. 8vo. 2s. 6d. KEY, for Teachers only. Gl. 8vo. 5s.

REID.—A GRAMMAR OF TACITUS. By J. S. Reid, Litt.D., Fellow of Caius College, Cambridge. [*In the Press.*

A GRAMMAR OF VIRGIL. By the same. [*In preparation.*

ROBY.—Works by H. J. Roby, M.A., late Fellow of St. John's College, Cambridge.

A GRAMMAR OF THE LATIN LANGUAGE, from Plautus to Suetonius. Part I. Sounds, Inflexions, Word-formation, Appendices. Cr. 8vo. 9s. Part II. Syntax, Prepositions, etc. 10s. 6d.

*SCHOOL LATIN GRAMMAR. Cr. 8vo. 5s.

AN ELEMENTARY LATIN GRAMMAR. [*In the Press.*

*RUSH.—SYNTHETIC LATIN DELECTUS. With Notes and Vocabulary. By E. Rush, B.A. Ex. fcap. 8vo. 2s. 6d.

*RUST.—FIRST STEPS TO LATIN PROSE COMPOSITION. By Rev. G. Rust, M.A. 18mo. 1s. 6d. KEY, for Teachers only. By W. M. Yates. 18mo. 3s. 6d.

RUTHERFORD.—Works by the Rev. W. G. Rutherford, M.A., LL.D., Headmaster of Westminster.

REX LEX. A Short Digest of the principal Relations between the Latin, Greek, and Anglo-Saxon Sounds. 8vo. [*In preparation.*

THE NEW PHRYNICHUS; being a Revised Text of the Ecloga of the Grammarian Phrynichus. With Introduction and Commentary. 8vo. 18s. (See also *Macmillan's Greek Course.*)

SHUCKBURGH.—PASSAGES FROM LATIN AUTHORS FOR TRANSLATION INTO ENGLISH. Selected with a view to the needs of Candidates for the Cambridge Local, and Public Schools' Examinations. By E. S. Shuckburgh, M.A. Cr. 8vo. 2s.

*SIMPSON.—LATIN PROSE AFTER THE BEST AUTHORS: Cæsarian Prose. By F. P. Simpson, B.A. Ex. fcap. 8vo. 2s. 6d. KEY, for Teachers only. Ex. fcap. 8vo. 5s.

STRACHAN and WILKINS.—ANALECTA. Selected Passages for Translation. By J. S. Strachan, M.A., Professor of Greek, and A. S. Wilkins, Litt.D., Professor of Latin in the Owens College, Manchester. Cr. 8vo. 5s. KEY to Latin Passages. Cr. 8vo. 6d.

THRING.—Works by the Rev. E. Thring, M.A., late Headmaster of Uppingham.

A LATIN GRADUAL. A First Latin Construing Book for Beginners. With Coloured Sentence Maps. Fcap. 8vo. 2s. 6d.

A MANUAL OF MOOD CONSTRUCTIONS. Fcap. 8vo. 1s. 6d.

*WELCH and DUFFIELD.—LATIN ACCIDENCE AND EXERCISES ARRANGED FOR BEGINNERS. By W. Welch and C. G. Duffield, Assistant Masters at Cranleigh School. 18mo. 1s. 6d.

WHITE.—FIRST LESSONS IN GREEK. Adapted to GOODWIN'S GREEK GRAMMAR, and designed as an introduction to the ANABASIS OF XENOPHON. By JOHN WILLIAMS WHITE, Assistant Professor of Greek in Harvard University, U.S.A. Cr. 8vo. 3s. 6d.

WRIGHT.—Works by J. WRIGHT, M.A., late Headmaster of Sutton Coldfield School.

A HELP TO LATIN GRAMMAR; or, the Form and Use of Words in Latin, with Progressive Exercises. Cr. 8vo. 4s. 6d.

THE SEVEN KINGS OF ROME. An Easy Narrative, abridged from the First Book of Livy by the omission of Difficult Passages; being a First Latin Reading Book, with Grammatical Notes and Vocabulary. Fcap. 8vo. 3s. 6d.

FIRST LATIN STEPS; OR, AN INTRODUCTION BY A SERIES OF EXAMPLES TO THE STUDY OF THE LATIN LANGUAGE. Cr. 8vo. 3s.

ATTIC PRIMER. Arranged for the Use of Beginners. Ex. fcap. 8vo. 2s. 6d.

A COMPLETE LATIN COURSE, comprising Rules with Examples, Exercises, both Latin and English, on each Rule, and Vocabularies. Cr. 8vo. 2s. 6d.

ANTIQUITIES, ANCIENT HISTORY, AND PHILOSOPHY.

ARNOLD.—A HANDBOOK OF LATIN EPIGRAPHY. By W. T. ARNOLD, M.A. [*In preparation.*

THE ROMAN SYSTEM OF PROVINCIAL ADMINISTRATION TO THE ACCESSION OF CONSTANTINE THE GREAT. By the same. Cr. 8vo. 6s.

ARNOLD.—THE SECOND PUNIC WAR. Being Chapters from THE HISTORY OF ROME by the late THOMAS ARNOLD, D.D., Headmaster of Rugby. Edited, with Notes, by W. T. ARNOLD, M.A. With 8 Maps. Cr. 8vo. 5s.

***BEESLY.**—STORIES FROM THE HISTORY OF ROME. By Mrs. BEESLY. Fcap. 8vo. 2s. 6d.

BLACKIE.—HORÆ HELLENICÆ. By JOHN STUART BLACKIE, Emeritus Professor of Greek in the University of Edinburgh. 8vo. 12s.

BURN.—ROMAN LITERATURE IN RELATION TO ROMAN ART. By Rev. ROBERT BURN, M.A., late Fellow of Trinity College, Cambridge. Illustrated. Ex. cr. 8vo. 14s.

BURY.—A HISTORY OF THE LATER ROMAN EMPIRE FROM ARCADIUS TO IRENE, A.D. 395-800. By J. B. BURY, M.A., Fellow of Trinity College, Dublin. 2 vols. 8vo. 32s.

***CLASSICAL WRITERS.**—Edited by JOHN RICHARD GREEN, M.A., LL.D. Fcap. 8vo. 1s. 6d. each.

SOPHOCLES. By Prof. L. CAMPBELL, M.A.

EURIPIDES. By Prof. MAHAFFY, D.D.

DEMOSTHENES. By Prof. S. H. BUTCHER, M.A.

VIRGIL. By Prof. NETTLESHIP, M.A.

LIVY By Rev. W. W. CAPES, M.A.

TACITUS. By Prof. A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A.

MILTON. By Rev. STOPFORD A. BROOKE, M.A.

FREEMAN.—Works by EDWARD A. FREEMAN, D.C.L., LL.D., Regius Professor of Modern History in the University of Oxford.

HISTORY OF ROME. (*Historical Course for Schools.*) 18mo. [*In preparation.*

HISTORY OF GREECE. (*Historical Course for Schools.*) 18mo. [*In preparation.*

A SCHOOL HISTORY OF ROME. Cr. 8vo. [*In preparation.*

HISTORICAL ESSAYS. Second Series. [Greek and Roman History.] 8vo. 10s. 6d.

FYFFE.—A SCHOOL HISTORY OF GREECE. By C. A. FYFFE, M.A. Cr. 8vo. [*In preparation.*

GARDNER.—SAMOS AND SAMIAN COINS. An Essay. By PERCY GARDNER, Litt.D., Professor of Archæology in the University of Oxford. With Illustrations. 8vo. 7s. 6d.

GEDDES.—THE PROBLEM OF THE HOMERIC POEMS. By W. D. GEDDES, Principal of the University of Aberdeen. 8vo. 14s.

GLADSTONE.—Works by the Rt. Hon. W. E. GLADSTONE, M.P.

THE TIME AND PLACE OF HOMER. Cr. 8vo. 6s. 6d.

LANDMARKS OF HOMERIC STUDY. Cr. 8vo. 2s. 6d.

*A PRIMER OF HOMER. 18mo. 1s.

GOW.—A COMPANION TO SCHOOL CLASSICS. By JAMES GOW, Litt.D., Master of the High School, Nottingham. With Illustrations. 2d Ed., revised. Cr. 8vo. 6s.

HARRISON and VERRALL.—MYTHOLOGY AND MONUMENTS OF ANCIENT ATHENS. Translation of a portion of the "Attica" of Pausanias. By MARGARET DE G. VERRALL. With Introductory Essay and Archæological Commentary by JANE E. HARRISON. With Illustrations and Plans. Cr. 8vo. 16s.

JEBB.—Works by R. C. JEBB, Litt.D., Professor of Greek in the University of Cambridge.

THE ATTIC ORATORS FROM ANTIPHON TO ISAEOS. 2 vols. 8vo. 25s.

*A PRIMER OF GREEK LITERATURE. 18mo. 1s.

(See also *Classical Series*.)

KIEPERT.—MANUAL OF ANCIENT GEOGRAPHY. By Dr. H. KIEPERT. Cr. 8vo. 5s.

LANCIANI.—ANCIENT ROME IN THE LIGHT OF RECENT DISCOVERIES. By RODOLFO LANCIANI, Professor of Archæology in the University of Rome. Illustrated. 4to. 24s.

MAHAFFY.—Works by J. P. MAHAFFY, D.D., Fellow of Trinity College, Dublin, and Professor of Ancient History in the University of Dublin.

SOCIAL LIFE IN GREECE; from Homer to Menander. Cr. 8vo. 9s.

GREEK LIFE AND THOUGHT; from the Age of Alexander to the Roman Conquest. Cr. 8vo. 12s. 6d.

THE GREEK WORLD UNDER ROMAN SWAY. From Plutarch to Polybius. Cr. 8vo. 10s. 6d.

RAMBLES AND STUDIES IN GREECE. With Illustrations. With Map. Cr. 8vo. 10s. 6d.

A HISTORY OF CLASSICAL GREEK LITERATURE. Cr. 8vo. Vol. I. The Poets, with an Appendix on Homer by Prof. SAYCE. In two parts. Vol. II. The Prose Writers. In two parts. Part I. Herodotus to Plato. Part II. Isocrates to Aristotle. 4s. 6d. each.

*A PRIMER OF GREEK ANTIQUITIES. With Illustrations. 18mo. 1s.

*EURIPIDES. 18mo. 1s. 6d. (*Classical Writers.*)

MAYOR.—BIBLIOGRAPHICAL CLUE TO LATIN LITERATURE. Edited after HÜBNER. By Prof. JOHN E. B. MAYOR. Cr. 8vo. 10s. 6d.

NEWTON.—ESSAYS ON ART AND ARCHÆOLOGY. By Sir CHARLES NEWTON, K.C.B., D.C.L. 8vo. 12s. 6d.

PHILOLOGY.—THE JOURNAL OF PHILOLOGY. Edited by W. A. WRIGHT, M.A., J. BYWATER, M.A., and H. JACKSON, Litt.D. 4s. 6d. each (half-yearly).

RENDALL.—THE CRADLE OF THE ARYANS. By Prof. G. H. RENDALL. 8vo. 3s.

SAYCE.—THE ANCIENT EMPIRES OF THE EAST. By A. H. SAYCE, M.A., Deputy-Professor of Comparative Philology, Oxford. Cr. 8vo. 6s.

SCHMIDT and WHITE. AN INTRODUCTION TO THE RHYTHMIC AND METRIC OF THE CLASSICAL LANGUAGES. By Dr. J. H. HEINRICH SCHMIDT. Translated by JOHN WILLIAMS WHITE, Ph.D. 8vo. 10s. 6d.

SHUCHHARDT.—DR. SCHLIEMANN'S EXCAVATIONS AT TROY, TIRYNS, MYCENÆ, ORCHOMENOS, ITHACA, presented in the light of recent knowledge. By Dr. CARL SHUCHHARDT. Translated by EUGENIE SELLERS. Introduction by WALTER LEAF, Litt.D. Illustrated. 8vo. [*In the Press.*

SHUCKBURGH.—A SCHOOL HISTORY OF ROME. By E. S. SHUCKBURGH, M.A. Cr. 8vo. [*In preparation.*

***STEWART.**—THE TALE OF TROY. Done into English by AUBREY STEWART Gl. 8vo. 3s. 6d.

***TOZER.**—A PRIMER OF CLASSICAL GEOGRAPHY. By H. F. TOZER, M.A. 18mo. 1s.

WALDSTEIN.—CATALOGUE OF CASTS IN THE MUSEUM OF CLASSICAL ARCHÆOLOGY, CAMBRIDGE. By CHARLES WALDSTEIN, University Reader in Classical Archæology. Cr. 8vo. 1s. 6d.
*** Also an Edition on Large Paper, small 4to. 5s.

WILKINS.—Works by Prof. WILKINS, Litt.D., LL.D.
*A PRIMER OF ROMAN ANTIQUITIES. Illustrated. 18mo. 1s.
*A PRIMER OF ROMAN LITERATURE. 18mo. 1s.

WILKINS and ARNOLD.—A MANUAL OF ROMAN ANTIQUITIES. By Prof. A. S. WILKINS, Litt.D., and W. T. ARNOLD, M.A. Cr. 8vo. Illustrated. [*In preparation.*

MODERN LANGUAGES AND LITERATURE.

English; French; German; Modern Greek; Italian; Spanish.

ENGLISH.

***ABBOTT.**—A SHAKESPEARIAN GRAMMAR. An Attempt to Illustrate some of the Differences between Elizabethan and Modern English. By the Rev. E. A. ABBOTT, D.D., formerly Headmaster of the City of London School. Ex. fcap. 8vo. 6s.

***BACON.**—ESSAYS. With Introduction and Notes, by F. G. SELBY, M.A., Professor of Logic and Moral Philosophy, Deccan College, Poona. Gl. 8vo. 3s.; sewed, 2s. 6d.

***BURKE.**—REFLECTIONS ON THE FRENCH REVOLUTION. By the same. Gl. 8vo. 5s.

BROOKE.—*PRIMER OF ENGLISH LITERATURE. By Rev. STOPFORD A. BROOKE, M.A. 18mo. 1s.
EARLY ENGLISH LITERATURE. By the same. 2 vols. 8vo. [*Vol. I. In the Press.*

BUTLER.—HUDIBRAS. With Introduction and Notes, by ALFRED MILNES, M.A. Ex. fcap. 8vo. Part I. 3s. 6d. Parts II. and III. 4s. 6d.

CAMPBELL.—SELECTIONS. With Introduction and Notes, by CECIL M. BARROW, M.A., Principal of Victoria College, Palghât. Gl. 8vo. [*In preparation.*

COWPER.—*THE TASK: an Epistle to Joseph Hill, Esq.; TIROCINIUM, or a Review of the Schools; and THE HISTORY OF JOHN GILPIN. Edited, with Notes, by W. BENHAM, B.D. Gl. 8vo. 1s. (*Globe Readings from Standard Authors.*)
THE TASK. With Introduction and Notes, by F. J. ROWE, M.A., and W. T. WEBB, M.A., Professors of English Literature, Presidency College, Calcutta. [*In preparation.*

***DOWDEN.**—SHAKESPERE. By Prof. DOWDEN. 18mo. 1s.

DRYDEN.—SELECT PROSE WORKS. Edited, with Introduction and Notes, by Prof. C. D. YONGE. Fcap. 8vo. 2s. 6d.

***GLOBE READERS.** For Standards I.-VI. Edited by A. F. MURISON. Illustrated. Gl. 8vo.

Primer I. (48 pp.)	3d.	Book III. (232 pp.)	1s. 3d.
Primer II. (48 pp.)	3d.	Book IV. (328 pp.)	1s. 9d.
Book I. (132 pp.)	6d.	Book V. (408 pp.)	2s.
Book II. (136 pp.)	9d.	Book VI. (436 pp.)	2s. 6d.

***THE SHORTER GLOBE READERS.**—Illustrated. Gl. 8vo.

Primer I. (48 pp.)	3d.	Standard III. (178 pp.)	1s.
Primer II. (48 pp.)	3d.	Standard IV. (182 pp.)	1s.
Standard I. (90 pp.)	6d.	Standard V. (216 pp.)	1s. 3d.
Standard II. (124 pp.)	9d.	Standard VI. (228 pp.)	1s. 6d.

***GOLDSMITH.**—THE TRAVELLER, or a Prospect of Society; and the DESERTED VILLAGE. With Notes, Philological and Explanatory, by J. W. HALES, M.A. Cr. 8vo. 6d.

*THE TRAVELLER AND THE DESERTED VILLAGE. With Introduction and Notes, by A. BARRETT, B.A., Professor of English Literature, Elphinstone College, Bombay. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d. The Traveller (separately), 1s., sewed.

THE VICAR OF WAKEFIELD. With Introduction and Notes, by H. LITTLEDALE, B.A., Professor of History and English Literature, Baroda College. Gl. 8vo. [*In preparation.*

*THE VICAR OF WAKEFIELD. With a Memoir of Goldsmith, by Prof. MASSON. Gl. 8vo. 1s. (*Globe Readings from Standard Authors.*)

SELECT ESSAYS. With Introduction and Notes, by Prof. C. D. YONGE. Fcap. 8vo. 2s. 6d.

GOSSE.—A HISTORY OF EIGHTEENTH CENTURY LITERATURE (1660-1780). By EDMUND GOSSE, M.A. Cr. 8vo. 7s. 6d.

GRAY.—POEMS. With Introduction and Notes, by JOHN BRADSHAW, LL.D. Gl. 8vo. [*In the Press.*

*HALES.—LONGER ENGLISH POEMS. With Notes, Philological and Explanatory, and an Introduction on the Teaching of English, by J. W. HALES, M.A., Professor of English Literature at King's College, London. Ex. fcap. 8vo. 4s. 6d.

*HELPS.—ESSAYS WRITTEN IN THE INTERVALS OF BUSINESS. With Introduction and Notes, by F. J. ROWE, M.A., and W. T. WEBB, M.A. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*JOHNSON.—LIVES OF THE POETS. The Six Chief Lives (Milton, Dryden, Swift, Addison, Pope, Gray), with Macaulay's "Life of Johnson." With Preface and Notes by MATTHEW ARNOLD. Cr. 8vo. 4s. 6d.

*LAMB.—TALES FROM SHAKSPEARE. With Preface by the Rev. CANON AINGER, M.A., LL.D. Gl. 8vo. 2s. (*Globe Readings from Standard Authors.*)

*LITERATURE PRIMERS.—Edited by JOHN RICHARD GREEN, LL.D. 18mo. 1s. each.

ENGLISH GRAMMAR. By Rev. R. MORRIS, LL.D.

ENGLISH GRAMMAR EXERCISES. By R. MORRIS, LL.D., and H. C. BOWEN, M.A.

EXERCISES ON MORRIS'S PRIMER OF ENGLISH GRAMMAR. By J. WETHERELL, M.A.

ENGLISH COMPOSITION. By Professor NICHOL.

QUESTIONS AND EXERCISES ON ENGLISH COMPOSITION. By Prof. NICHOL and W. S. M'CORMICK.

ENGLISH LITERATURE. By STOPFORD BROOKE, M.A.

SHAKSPERE. By Professor DOWDEN.

THE CHILDREN'S TREASURY OF LYRICAL POETRY. Selected and arranged with Notes by FRANCIS TURNER PALGRAVE. In Two Parts. 1s. each.

PHILOLOGY. By J. PEILE, Litt.D.

ROMAN LITERATURE. By Prof. A. S. WILKINS, Litt.D.

GREEK LITERATURE. By Prof. JEBB, Litt.D.

HOMER. By the Rt. Hon. W. E. GLADSTONE, M.P.

A HISTORY OF ENGLISH LITERATURE IN FOUR VOLUMES. Cr. 8vo.

EARLY ENGLISH LITERATURE. By STOPFORD BROOKE, M.A. [*In preparation.*

ELIZABETHAN LITERATURE. (1560–1665.) By GEORGE SAINTSBURY. 7s. 6d.

EIGHTEENTH CENTURY LITERATURE. (1660–1780.) By EDMUND GOSSE, M.A. 7s. 6d.

THE MODERN PERIOD. By Prof. DOWDEN. [*In preparation.*

*MACMILLAN'S READING BOOKS.

PRIMER. 18mo. 48 pp. 2d.

BOOK I. for Standard I. 96 pp. 4d.

BOOK II. for Standard II. 144 pp. 5d.

BOOK III. for Standard III. 160 pp. 6d.

BOOK IV. for Standard IV. 176 pp. 8d.

BOOK V. for Standard V. 380 pp. 1s.

BOOK VI. for Standard VI. Cr. 8vo. 430 pp. 2s.

Book VI. is fitted for Higher Classes, and as an Introduction to English Literature

***MACMILLAN'S COPY BOOKS.**—1. Large Post 4to. Price 4d. each. 2. Post Oblong. Price 2d. each.

1. Initiatory Exercises and Short Letters.
2. Words consisting of Short Letters.
3. Long Letters. With Words containing Long Letters—Figures.
4. Words containing Long Letters.

4a. Practising and Revising Copy-Book. For Nos. 1 to 4.

5. Capitals and Short Half-Text. Words beginning with a Capital.
6. Half-Text Words beginning with Capitals—Figures.
7. Small-Hand and Half-Text. With Capitals and Figures.
8. Small-Hand and Half-Text. With Capitals and Figures.

8a. Practising and Revising Copy-Book. For Nos. 5 to 8.

9. Small-Hand Single Headlines—Figures.
10. Small-Hand Single Headlines—Figures.
11. Small-Hand Double Headlines—Figures.
12. Commercial and Arithmetical Examples, &c.

12a. Practising and Revising Copy-Book. For Nos. 8 to 12.

Nos. 3, 4, 5, 6, 7, 8, 9 *may be had with Goodman's Patent Sliding Copies.* Large Post 4to. Price 6d. each.

MARTIN.—*THE POET'S HOUR: Poetry selected and arranged for Children. By Frances Martin. 18mo. 2s. 6d.

*SPRING-TIME WITH THE POETS. By the same. 18mo. 3s. 6d.

***MILTON.**—PARADISE LOST. Books I. and II. With Introduction and Notes, by Michael Macmillan, B.A., Professor of Logic and Moral Philosophy, Elphinstone College, Bombay. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d. Or separately, 1s. 3d.; sewed, 1s. each.

*L'ALLEGRO, IL PENSEROSO, LYCIDAS, ARCADES, SONNETS, &c. With Introduction and Notes, by W. Bell, M.A., Professor of Philosophy and Logic, Government College, Lahore. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*COMUS. By the same. Gl. 8vo. 1s. 3d.; sewed, 1s.

*SAMSON AGONISTES. By H. M. Percival, M.A., Professor of English Literature, Presidency College, Calcutta. Gl. 8vo. 2s.; sewed, 1s. 9d.

*INTRODUCTION TO THE STUDY OF MILTON. By Stopford Brooke, M.A. Fcap. 8vo. 1s. 6d. (*Classical Writers.*)

MORRIS.—Works by the Rev. R. Morris, LL.D.

*PRIMER OF ENGLISH GRAMMAR. 18mo. 1s.

*ELEMENTARY LESSONS IN HISTORICAL ENGLISH GRAMMAR, containing Accidence and Word-Formation. 18mo. 2s. 6d.

*HISTORICAL OUTLINES OF ENGLISH ACCIDENCE, comprising Chapters on the History and Development of the Language, and on Word-Formation. Ex. fcap. 8vo. 6s.

MORRIS and KELLNER.—HISTORICAL OUTLINES OF ENGLISH SYNTAX. By Rev. R. Morris and Dr. L. Kellner. [*In preparation.*

NICHOL.—A SHORT HISTORY OF ENGLISH LITERATURE. By Prof. John Nichol. Gl. 8vo. [*In preparation.*

OLIPHANT.—THE OLD AND MIDDLE ENGLISH. By T. L. Kington Oliphant. New Ed., revised and enlarged, of "The Sources of Standard English." New Ed. Gl. 8vo. [*In the Press.*

THE NEW ENGLISH. By the same. 2 vols. Cr. 8vo. 21s.

***PALGRAVE.**—THE CHILDREN'S TREASURY OF LYRICAL POETRY. Selected and arranged, with Notes, by Francis T. Palgrave. 18mo. 2s. 6d. Also in Two Parts. 1s. each.

PATMORE.—THE CHILDREN'S GARLAND FROM THE BEST POETS. Selected and arranged by Coventry Patmore. Gl. 8vo. 2s. (*Globe Readings from Standard Authors.*)

PLUTARCH.—Being a Selection from the Lives which illustrate Shakespeare. North's Translation. Edited, with Introductions, Notes, Index of Names, and Glossarial Index, by Prof. W. W. Skeat, Litt.D. Cr. 8vo. 6s.

***RANSOME.**—SHORT STUDIES OF SHAKESPEARE'S PLOTS. By Cyril Ransome, Professor of Modern History and Literature, Yorkshire College, Leeds. Cr. 8vo. 3s. 6d.

***RYLAND.**—CHRONOLOGICAL OUTLINES OF ENGLISH LITERATURE. By F. Ryland, M.A. Cr. 8vo. 6s.

SAINTSBURY.—A HISTORY OF ELIZABETHAN LITERATURE. 1560-1665. By George Saintsbury. Cr. 8vo. 7s. 6d.

SCOTT.—*LAY OF THE LAST MINSTREL, and THE LADY OF THE LAKE. Edited, with Introduction and Notes, by Francis Turner Palgrave. Gl. 8vo. 1s. (*Globe Readings from Standard Authors.*)

*THE LAY OF THE LAST MINSTREL. With Introduction and Notes, by G. H. Stuart, M.A., and E. H. Elliot, B.A. Gl. 8vo. Introduction and Canto I. 9d. sewed. Cantos I. to III. 1s. 3d.; sewed, 1s. Cantos IV. to VI. 1s. 3d.; sewed, 1s.

*MARMION, and THE LORD OF THE ISLES. By F. T. Palgrave. Gl. 8vo. 1s. (*Globe Readings from Standard Authors.*)

*MARMION. With Introduction and Notes, by Michael Macmillan, B.A. Gl. 8vo. 3s.; sewed, 2s. 6d.

*THE LADY OF THE LAKE. By G. H. Stuart, M.A. Gl. 8vo. 2s. 6d.; sewed, 2s.

*ROKEBY. With Introduction and Notes, by Michael Macmillan, B.A. Gl. 8vo. 3s.; sewed, 2s. 6d.

SHAKESPEARE.—*A SHAKESPEARIAN GRAMMAR. By Rev. E. A. Abbott, D.D. Gl. 8vo. 6s.

A SHAKESPEARE MANUAL. By F. G. Fleay, M.A. 2d Ed. Ex. fcap. 8vo. 4s. 6d.

*PRIMER OF SHAKESPERE. By Prof. Dowden. 18mo. 1s.

*SHORT STUDIES OF SHAKESPEARE'S PLOTS. By Cyril Ransome, M.A. Cr. 8vo. 3s. 6d.

*THE TEMPEST. With Introduction and Notes, by K. Deighton, late Principal of Agra College. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*MUCH ADO ABOUT NOTHING. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*A MIDSUMMER NIGHT'S DREAM. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*THE MERCHANT OF VENICE. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

AS YOU LIKE IT. By the same. Gl. 8vo.

*TWELFTH NIGHT. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*THE WINTER'S TALE. By the same. Gl. 8vo. 2s.; sewed, 1s. 9d.

*RICHARD II. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*KING JOHN. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*HENRY V. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*RICHARD III. By C. H. Tawney, M.A., Principal and Professor of English Literature, Presidency College, Calcutta. Gl. 8vo. 2s. 6d.; sewed, 2s.

*CORIOLANUS. By K. Deighton. Gl. 8vo. 2s. 6d.; sewed, 2s.

*JULIUS CÆSAR. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*MACBETH. By the same. Gl. 8vo. 1s. 9d.; sewed, 1s. 6d.

*HAMLET. By the same. Gl. 8vo. 2s. 6d.; sewed, 2s.

KING LEAR. By the same. Gl. 8vo.

*OTHELLO. By the same. Gl. 8vo. 2s.; sewed, 1s. 9d.

ANTONY AND CLEOPATRA. By the same. Gl. 8vo. 2s. 6d.; sewed, 2s.

*CYMBELINE. By the same. Gl. 8vo. 2s. 6d.; sewed, 2s.

***SONNENSCHEIN and MEIKLEJOHN.**—THE ENGLISH METHOD OF TEACHING TO READ. By A. Sonnenschein and J. M. D. Meiklejohn, M.A. Fcap. 8vo.

THE NURSERY BOOK, containing all the Two-Letter Words in the Language. 1d. (Also in Large Type on Sheets for School Walls. 5s.)

THE FIRST COURSE, consisting of Short Vowels with Single Consonants. 7d.

THE SECOND COURSE, with Combinations and Bridges, consisting of Short Vowels with Double Consonants. 7d.

THE THIRD AND FOURTH COURSES, consisting of Long Vowels, and all the Double Vowels in the Language. 7d.

*SOUTHEY.—LIFE OF NELSON. With Introduction and Notes, by MICHAEL MACMILLAN, B.A. Gl. 8vo. 3s.; sewed, 2s. 6d.

TAYLOR.—WORDS AND PLACES; or, Etymological Illustrations of History, Ethnology, and Geography. By Rev. ISAAC TAYLOR, Litt.D. With Maps. Gl. 8vo. 6s.

TENNYSON.—THE COLLECTED WORKS OF LORD TENNYSON. An Edition for Schools. In Four Parts. Cr. 8vo. 2s. 6d. each.

TENNYSON FOR THE YOUNG. Edited, with Notes for the Use of Schools, by the Rev. ALFRED AINGER, LL.D., Canon of Bristol. 18mo. 1s. net.

[In preparation.

*SELECTIONS FROM TENNYSON. With Introduction and Notes, by F. J. ROWE, M.A., and W. T. WEBB, M.A. Gl. 8vo. 3s. 6d.

This selection contains:—Recollections of the Arabian Nights, The Lady of Shalott, Œnone, The Lotos Eaters, Ulysses, Tithonus, Morte d'Arthur, Sir Galahad, Dora, Ode on the Death of the Duke of Wellington, and The Revenge.

ENOCH ARDEN. By W. T. WEBB, M.A. Gl. 8vo. [*In the Press.*

*THE COMING OF ARTHUR, AND THE PASSING OF ARTHUR. By F. J. ROWE, M.A. Gl. 8vo. 2s.

THE PRINCESS. By P. M. WALLACE, B.A. [*In the Press.*

THRING.—THE ELEMENTS OF GRAMMAR TAUGHT IN ENGLISH. By EDWARD THRING, M.A. With Questions. 4th Ed. 18mo. 2s.

*VAUGHAN.—WORDS FROM THE POETS. By C. M. VAUGHAN. 18mo. 1s.

WARD.—THE ENGLISH POETS. Selections, with Critical Introductions by various Writers and a General Introduction by MATTHEW ARNOLD. Edited by T. H. WARD, M.A. 4 Vols. Vol. I. CHAUCER TO DONNE.—Vol. II. BEN JONSON TO DRYDEN.—Vol. III. ADDISON TO BLAKE.—Vol. IV. WORDSWORTH TO ROSSETTI. 2d Ed. Cr. 8vo. 7s. 6d. each.

*WETHERELL.—EXERCISES ON MORRIS'S PRIMER OF ENGLISH GRAMMAR. By JOHN WETHERELL, M.A., Headmaster of Towcester Grammar School. 18mo. 1s.

WOODS.—*A FIRST POETRY BOOK. By M. A. WOODS, Head Mistress of the Clifton High School for Girls. Fcap. 8vo. 2s. 6d.

*A SECOND POETRY BOOK. By the same. In Two Parts. 2s. 6d. each.

*A THIRD POETRY BOOK. By the same. 4s. 6d.

HYMNS FOR SCHOOL WORSHIP. By the same. 18mo. 1s. 6d.

WORDSWORTH.—SELECTIONS. With Introduction and Notes, by F. J. ROWE, M.A., and W. T. WEBB, M.A. Gl. 8vo. [*In preparation.*

YONGE.—*A BOOK OF GOLDEN DEEDS. By CHARLOTTE M. YONGE. Gl. 8vo 2s.

*THE ABRIDGED BOOK OF GOLDEN DEEDS. 18mo. 1s.

FRENCH.

BEAUMARCHAIS.—LE BARBIER DE SEVILLE. With Introduction and Notes. By L. P. BLOUET. Fcap. 8vo. 3s. 6d.

*BOWEN.—FIRST LESSONS IN FRENCH. By H. COURTHOPE BOWEN, M.A. Ex. fcap. 8vo. 1s.

BREYMANN.—Works by HERMANN BREYMANN, Ph.D., Professor of Philology in the University of Munich.

A FRENCH GRAMMAR BASED ON PHILOLOGICAL PRINCIPLES. Ex. fcap. 8vo. 4s. 6d.

FIRST FRENCH EXERCISE BOOK. Ex. fcap. 8vo. 4s. 6d.

SECOND FRENCH EXERCISE BOOK. Ex. fcap. 8vo. 2s. 6d.

FASNACHT.—Works by G. E. FASNACHT, late Assistant Master at Westminster.

THE ORGANIC METHOD OF STUDYING LANGUAGES. Ex. fcap. 8vo. I. French. 3s. 6d.

A SYNTHETIC FRENCH GRAMMAR FOR SCHOOLS. Cr. 8vo. 3s. 6d.

GRAMMAR AND GLOSSARY OF THE FRENCH LANGUAGE OF THE SEVENTEENTH CENTURY. Cr. 8vo. [*In preparation.*

MACMILLAN'S PRIMARY SERIES OF FRENCH READING BOOKS.—Edited by G. E. Fasnacht. With Illustrations, Notes, Vocabularies, and Exercises. Gl. 8vo.

FRENCH READINGS FOR CHILDREN. By G. E. Fasnacht. [*May* 1891.

*CORNAZ—NOS ENFANTS ET LEURS AMIS. By Edith Harvey. 1s. 6d.

*DE MAISTRE—LA JEUNE SIBÉRIENNE ET LE LÉPREUX DE LA CITÉ D'AOSTE. By Stephane Barlet, B.Sc. etc. 1s. 6d.

*FLORIAN—FABLES. By Rev. Charles Yeld, M.A., Headmaster of University School, Nottingham. 1s. 6d.

*LA FONTAINE—A SELECTION OF FABLES. By L. M. Moriarty, B.A., Assistant Master at Harrow. 2s. 6d.

LAMARTINE—JEANNE D'ARC. By M. de G. Verrall. [*In the Press.*

*MOLESWORTH—FRENCH LIFE IN LETTERS. By Mrs. Molesworth. 1s. 6d.

*PERRAULT—CONTES DE FÉES. By G. E. Fasnacht. 1s. 6d.

MACMILLAN'S PROGRESSIVE FRENCH COURSE.—By G. E. Fasnacht. Ex. fcap. 8vo.

*First Year, containing Easy Lessons on the Regular Accidence. 1s.

*Second Year, containing an Elementary Grammar with copious Exercises, Notes, and Vocabularies. 2s.

*Third Year, containing a Systematic Syntax, and Lessons in Composition. 2s. 6d.

THE TEACHER'S COMPANION TO MACMILLAN'S PROGRESSIVE FRENCH COURSE. With Copious Notes, Hints for Different Renderings, Synonyms, Philological Remarks, etc. By G. E. Fasnacht. Ex. fcap. 8vo. Each Year 4s. 6d.

***MACMILLAN'S FRENCH COMPOSITION.**—By G. E. Fasnacht. Ex. fcap. 8vo. Part I. Elementary. 2s. 6d. Part II. Advanced. [*In the Press.*

THE TEACHER'S COMPANION TO MACMILLAN'S COURSE OF FRENCH COMPOSITION. By G. E. Fasnacht. Part I. Ex. fcap. 8vo. 4s. 6d.

MACMILLAN'S PROGRESSIVE FRENCH READERS. By G. E. Fasnacht. Ex. fcap. 8vo.

*First Year, containing Tales, Historical Extracts, Letters, Dialogues, Ballads, Nursery Songs, etc., with Two Vocabularies: (1) in the order of subjects; (2) in alphabetical order. With Imitative Exercises. 2s. 6d.

*Second Year, containing Fiction in Prose and Verse, Historical and Descriptive Extracts, Essays, Letters, Dialogues, etc. With Imitative Exercises. 2s. 6d.

MACMILLAN'S FOREIGN SCHOOL CLASSICS. Edited by G. E. Fasnacht. 18mo.

*CORNEILLE—LE CID. By G. E. Fasnacht. 1s.

*DUMAS—LES DEMOISELLES DE ST. CYR. By Victor Oger, Lecturer at University College, Liverpool. 1s. 6d.

LA FONTAINE'S FABLES. Books I.–VI. By L. M. Moriarty, B.A., Assistant Master at Harrow. [*In preparation.*

*MOLIÈRE—L'AVARE. By the same. 1s.

*MOLIÈRE—LE BOURGEOIS GENTILHOMME. By the same. 1s. 6d.

*MOLIÈRE—LES FEMMES SAVANTES. By G. E. Fasnacht. 1s.

*MOLIÈRE—LE MISANTHROPE. By the same. 1s.

*MOLIÈRE—LE MÉDECIN MALGRE LUI. By the same. 1s.

MOLIÈRE—LES PRÉCIEUSES RIDICULES. By the same. [*In the Press.*

*RACINE—BRITANNICUS. By E. Pellissier, M.A. 2s.

*FRENCH READINGS FROM ROMAN HISTORY. Selected from various Authors, by C. Colbeck, M.A., Assistant Master at Harrow. 4s. 6d.

*SAND, GEORGE—LA MARE AU DIABLE. By W. E. Russell, M.A., Assistant Master at Haileybury. 1s.

*SANDEAU, JULES—MADEMOISELLE DE LA SEIGLIÈRE. By H. C. Steel, Assistant Master at Winchester. 1s. 6d.

THIERS'S HISTORY OF THE EGYPTIAN EXPEDITION. By Rev. H. A. Bull, M.A., formerly Assistant Master at Wellington. [*In preparation.*

*VOLTAIRE—CHARLES XII. By G. E. Fasnacht. 3s. 6d.

***MASSON.**—A COMPENDIOUS DICTIONARY OF THE FRENCH LANGUAGE. Adapted from the Dictionaries of Professor A. Elwall. By Gustave Masson. Cr. 8vo. 6s.

MOLIÈRE.—LE MALADE IMAGINAIRE. With Introduction and Notes, by F. Tarver, M.A., Assistant Master at Eton. Fcap. 8vo. 2s. 6d.

***PELLISSIER.**—FRENCH ROOTS AND THEIR FAMILIES. A Synthetic Vocabulary, based upon Derivations. By E. Pellissier, M.A., Assistant Master at Clifton College. Gl. 8vo. 6s.

GERMAN.

BEHAGEL.—THE GERMAN LANGUAGE. By Dr. Otto Behagel. Translated by Emil Trechmann, B.A., Ph.D., Lecturer in Modern Literature in the University of Sydney, N.S.W. Gl. 8vo. [*In the Press.*

HUSS.—A SYSTEM OF ORAL INSTRUCTION IN GERMAN, by means of Progressive Illustrations and Applications of the leading Rules of Grammar. By H. C. O. Huss, Ph.D. Cr. 8vo. 5s.

MACMILLAN'S PROGRESSIVE GERMAN COURSE. By G. E. Fasnacht. Ex. fcap. 8vo.

*First Year. Easy lessons and Rules on the Regular Accidence. 1s. 6d.

*Second Year. Conversational Lessons in Systematic Accidence and Elementary Syntax. With Philological Illustrations and Etymological Vocabulary. 3s. 6d.

Third Year. [*In the Press.*

TEACHER'S COMPANION TO MACMILLAN'S PROGRESSIVE GERMAN COURSE. With copious Notes, Hints for Different Renderings, Synonyms, Philological Remarks, etc. By G. E. Fasnacht. Ex. fcap. 8vo. First Year. 4s. 6d. Second Year. 4s. 6d.

MACMILLAN'S GERMAN COMPOSITION. By G. E. Fasnacht. Ex. fcap. 8vo.

*I. FIRST COURSE. Parallel German-English Extracts and Parallel English-German Syntax. 2s. 6d.

TEACHER'S COMPANION TO MACMILLAN'S GERMAN COMPOSITION. By G. E. Fasnacht. First Course. Gl. 8vo. 4s. 6d.

MACMILLAN'S PROGRESSIVE GERMAN READERS. By G. E. Fasnacht. Ex. fcap. 8vo.

*First Year, containing an Introduction to the German order of Words, with Copious Examples, extracts from German Authors in Prose and Poetry; Notes, and Vocabularies. 2s. 6d.

MACMILLAN'S PRIMARY SERIES OF GERMAN READING BOOKS. Edited by G. E. Fasnacht. With Notes, Vocabularies, and Exercises. Gl. 8vo.

*GRIMM—KINDER UND HAUSMÄRCHEN. By G. E. Fasnacht. 2s. 6d.

*HAUFF—DIE KARAVANE. By Herman Hager, Ph.D., Lecturer in the Owens College, Manchester. 3s.

*SCHMID, CHR. VON—H. VON EICHENFELS. By G. E. Fasnacht. 2s. 6d.

MACMILLAN'S FOREIGN SCHOOL CLASSICS.—Edited by G. E. Fasnacht. 18mo.

FREYTAG (G.)—DOKTOR LUTHER. By F. Storr, M.A., Headmaster of the Modern Side, Merchant Taylors' School. [*In preparation.*

*GOETHE—GÖTZ VON BERLICHINGEN. By H. A. Bull, M.A., Assistant Master at Wellington. 2s.

*GOETHE—FAUST. Part I., followed by an Appendix on Part II. By Jane Lee, Lecturer in German Literature at Newnham College, Cambridge. 4s. 6d.

*HEINE—SELECTIONS FROM THE REISEBILDER AND OTHER PROSE WORKS. By C. Colbeck, M.A., Assistant Master at Harrow. 2s. 6d.

LESSING—MINNA VON BARNHELM. By James Sime, M.A. [*In preparation.*

*SCHILLER—SELECTIONS FROM SCHILLER'S LYRICAL POEMS. With a Memoir of Schiller. By E. J. Turner, B.A., and E. D. A. Morshead, M.A., Assistant Masters at Winchester. 2s. 6d.

*SCHILLER—DIE JUNGFRAU VON ORLEANS. By Joseph Gostwick. 2s. 6d.

*SCHILLER—MARIA STUART. By C. Sheldon, D.Litt., of the Royal Academical Institution, Belfast. 2s. 6d.

*SCHILLER—WILHELM TELL. By G. E. Fasnacht. 2s. 6d.

*SCHILLER—WALLENSTEIN. Part I. DAS LAGER. By H. B. Cotterill, M.A. 2s.

*UHLAND—SELECT BALLADS. Adapted as a First Easy Reading Book for Beginners. With Vocabulary. By G. E. Fasnacht. 1s.

***PYLODET.**—NEW GUIDE TO GERMAN CONVERSATION; containing an Alphabetical List of nearly 800 Familiar Words; followed by Exercises, Vocabulary of Words in frequent use, Familiar Phrases and Dialogues, a Sketch of German Literature, Idiomatic Expressions, etc. By L. Pylodet. 18mo. 2s. 6d.

WHITNEY.—A COMPENDIOUS GERMAN GRAMMAR. By W. D. Whitney, Professor of Sanskrit and Instructor in Modern Languages in Yale College. Cr. 8vo. 4s. 6d.

A GERMAN READER IN PROSE AND VERSE. By the same. With Notes and Vocabulary. Cr. 8vo. 5s.

***WHITNEY and EDGREN.**—A COMPENDIOUS GERMAN AND ENGLISH DICTIONARY, with Notation of Correspondences and Brief Etymologies. By Prof. W. D. Whitney, assisted by A. H. Edgren. Cr. 8vo. 7s. 6d.

THE GERMAN-ENGLISH PART, separately, 5s.

MODERN GREEK.

VINCENT and DICKSON.—HANDBOOK TO MODERN GREEK. By Sir Edgar Vincent, K.C.M.G., and T. G. Dickson, M.A. With Appendix on the relation of Modern and Classical Greek by Prof. Jebb. Cr. 8vo. 6s.

ITALIAN.

DANTE.—THE PURGATORY OF DANTE. With Translation and Notes, by A. J. Butler, M.A. Cr. 8vo. 12s. 6d.

THE PARADISO OF DANTE. With Translation and Notes, by the same. 2d. Ed. Cr. 8vo. 12s. 6d.

READINGS ON THE PURGATORIO OF DANTE. Chiefly based on the Commentary of Benvenuto Da Imola. By the Hon. W. Warren Vernon, M.A. With an Introduction by the Very Rev. the Dean of St. Paul's. 2 vols. Cr. 8vo. 24s.

SPANISH.

CALDERON.—FOUR PLAYS OF CALDERON. With Introduction and Notes. By Norman MacColl, M.A. Cr. 8vo. 14s.

The four plays here given are *El Principe Constante*, *La Vida es Sueno*, *El Alcalde de Zalamea*, and *El Escondido y La Tapada*.

MATHEMATICS.

Arithmetic, Book-keeping, Algebra, Euclid and Pure Geometry, Geometrical Drawing, Mensuration, Trigonometry, Analytical Geometry (Plane and Solid), Problems and Questions in Mathematics, Higher Pure Mathematics, Mechanics (Statics, Dynamics, Hydrostatics, Hydrodynamics: see also Physics), Physics (Sound, Light, Heat, Electricity, Elasticity, Attractions, &c.), Astronomy, Historical.

ARITHMETIC.

*ALDIS.—THE GREAT GIANT ARITHMOS. A most Elementary Arithmetic for Children. By Mary Steadman Aldis. Illustrated. Gl. 8vo. 2s. 6d.

ARMY PRELIMINARY EXAMINATION, SPECIMENS OF PAPERS SET AT THE, 1882-89.—With Answers to the Mathematical Questions. Subjects: Arithmetic, Algebra, Euclid, Geometrical Drawing, Geography, French, English Dictation. Cr. 8vo. 3s. 6d.

*BRADSHAW.—A COURSE OF EASY ARITHMETICAL EXAMPLES FOR BEGINNERS. By J. G. Bradshaw, B.A., Assistant Master at Clifton College. Gl. 8vo. 2s. With Answers, 2s. 6d.

*BROOKSMITH.—ARITHMETIC IN THEORY AND PRACTICE. By J. Brooksmith, M.A. Cr. 8vo. 4s. 6d. KEY. Crown 8vo. 10s. 6d.

*BROOKSMITH.—ARITHMETIC FOR BEGINNERS. By J. and E. J. Brooksmith. Gl. 8vo. 1s. 6d.

CANDLER.—HELP TO ARITHMETIC. Designed for the use of Schools. By H. Candler, Mathematical Master of Uppingham School. 2d Ed. Ex. fcap. 8vo. 2s. 6d.

*DALTON.—RULES AND EXAMPLES IN ARITHMETIC. By the Rev. T. Dalton, M.A., Assistant Master at Eton. New Ed., with Answers. 18mo. 2s. 6d.

*GOYEN.—HIGHER ARITHMETIC AND ELEMENTARY MENSURATION. By P. Goyen, Inspector of Schools, Dunedin, New Zealand. Cr. 8vo. 5s.

*HALL and KNIGHT.—ARITHMETICAL EXERCISES AND EXAMINATION PAPERS. With an Appendix containing Questions in Logarithms and Mensuration. By H. S. Hall, M.A., Master of the Military and Engineering Side, Clifton College, and S. R. Knight, B.A. Gl. 8vo. 2s. 6d.

LOCK.—Works by Rev. J. B. Lock, M.A., Senior Fellow and Bursar of Gonville and Caius College, Cambridge.

*ARITHMETIC FOR SCHOOLS. With Answers and 1000 additional Examples for Exercise. 3d Ed., revised. Gl. 8vo. 4s. 6d. KEY. Cr. 8vo. 10s. 6d.

*ARITHMETIC FOR BEGINNERS. A School Class-Book of Commercial Arithmetic. Gl. 8vo. 2s. 6d. KEY. Cr. 8vo. 8s. 6d.

*A SHILLING BOOK OF ARITHMETIC, FOR ELEMENTARY SCHOOLS. 18mo. 1s. With Answers. 1s. 6d.

*PEDLEY.—EXERCISES IN ARITHMETIC for the Use of Schools. Containing more than 7000 original Examples. By Samuel Pedley. Cr. 8vo. 5s. Also in Two Parts, 2s. 6d. each.

SMITH.—Works by Rev. Barnard Smith, M.A., late Fellow and Senior Bursar of St. Peter's College, Cambridge.

ARITHMETIC AND ALGEBRA, in their Principles and Application; with numerous systematically arranged Examples taken from the Cambridge Examination Papers, with especial reference to the Ordinary Examination for the B.A. Degree. New Ed., carefully revised. Cr. 8vo. 10s. 6d.

*ARITHMETIC FOR SCHOOLS. Cr. 8vo. 4s. 6d. KEY. Cr. 8vo. 8s. 6d.

EXERCISES IN ARITHMETIC. Cr. 8vo. 2s. With Answers, 2s. 6d. Answers separately, 6d.

SCHOOL CLASS-BOOK OF ARITHMETIC. 18mo. 3s. Or separately, in Three Parts, 1s. each. KEYS. Parts I., II., and III., 2s. 6d. each.

SHILLING BOOK OF ARITHMETIC. 18mo. Or separately, Part I., 2d.; Part II., 3d.; Part III., 7d. Answers, 6d. KEY. 18mo. 4s. 6d.

*THE SAME, with Answers. 18mo, cloth. 1s. 6d.

EXAMINATION PAPERS IN ARITHMETIC. 18mo. 1s. 6d. The Same, with Answers. 18mo. 2s. Answers, 6d. KEY. 18mo. 4s. 6d.

THE METRIC SYSTEM OF ARITHMETIC, ITS PRINCIPLES AND APPLICATIONS, with Numerous Examples. 18mo. 3d.

A CHART OF THE METRIC SYSTEM, on a Sheet, size 42 in. by 34 in. on Roller. 3s. 6d. Also a Small Chart on a Card. Price 1d.

EASY LESSONS IN ARITHMETIC, combining Exercises in Reading, Writing, Spelling, and Dictation. Part I. Cr. 8vo. 9d.

EXAMINATION CARDS IN ARITHMETIC. With Answers and Hints. Standards I. and II., in box, 1s. Standards III., IV., and V., in boxes, 1s. each. Standard VI. in Two Parts, in boxes, 1s. each.

A and B papers, of nearly the same difficulty, are given so as to prevent copying, and the colours of the A and B papers differ in each Standard, and from those of every other Standard, so that a master or mistress can see at a glance whether the children have the proper papers.

BOOK-KEEPING.

*THORNTON.—FIRST LESSONS IN BOOK-KEEPING. By J. THORNTON. Cr. 8vo. 2s. 6d. KEY. Oblong 4to. 10s. 6d.

*PRIMER OF BOOK-KEEPING. 18mo. 1s. KEY. Demy 8vo. 2s. 6d.

ALGEBRA.

*DALTON.—RULES AND EXAMPLES IN ALGEBRA. By Rev. T. DALTON, Assistant Master at Eton. Part I. 18mo. 2s. KEY. Cr. 8vo. 7s. 6d. Part II. 18mo. 2s. 6d.

HALL and KNIGHT.—Works by H. S. HALL, M.A., Master of the Military and Engineering Side, Clifton College, and S. R. KNIGHT, B.A.

*ELEMENTARY ALGEBRA FOR SCHOOLS. 6th Ed., revised and corrected. Gl. 8vo, bound in maroon coloured cloth, 3s. 6d.; with Answers, bound in green coloured cloth, 4s. 6d. KEY. 8s. 6d.

*ALGEBRAICAL EXERCISES AND EXAMINATION PAPERS. To accompany ELEMENTARY ALGEBRA. 2d Ed., revised. Gl. 8vo. 2s. 6d.

*HIGHER ALGEBRA. 3d Ed. Cr. 8vo. 7s. 6d. KEY. Cr. 8vo. 10s. 6d.

*JONES and CHEYNE.—ALGEBRAICAL EXERCISES. Progressively Arranged. By Rev. C. A. JONES and C. H. CHEYNE, M.A., late Mathematical Masters at Westminster School. 18mo. 2s. 6d.

KEY. By Rev. W. FAILES, M.A., Mathematical Master at Westminster School. Cr. 8vo. 7s. 6d.

SMITH.—ARITHMETIC AND ALGEBRA, in their Principles and Application; with numerous systematically arranged Examples taken from the Cambridge Examination Papers, with especial reference to the Ordinary Examination for the B.A. Degree. By Rev. BARNARD SMITH, M.A. New Edition, carefully revised. Cr. 8vo. 10s. 6d.

SMITH.—Works by CHARLES SMITH, M.A., Master of Sidney Sussex College, Cambridge.

*ELEMENTARY ALGEBRA. 2d Ed., revised. Gl. 8vo. 4s. 6d.

*A TREATISE ON ALGEBRA. 2d Ed. Cr. 8vo. 7s. 6d. KEY. Cr. 8vo. 10s. 6d.

TODHUNTER.—Works by ISAAC TODHUNTER, F.R.S.

*ALGEBRA FOR BEGINNERS. 18mo. 2s. 6d. KEY. Cr. 8vo. 6s. 6d.

*ALGEBRA FOR COLLEGES AND SCHOOLS. Cr. 8vo. 7s. 6d. KEY. Cr. 8vo. 10s. 6d.

EUCLID AND PURE GEOMETRY.

COCKSHOTT and WALTERS.—A TREATISE ON GEOMETRICAL CONICS. In accordance with the Syllabus of the Association for the Improvement of Geometrical Teaching. By A. COCKSHOTT, M.A., Assistant Master at Eton, and Rev. F. B. WALTERS, M.A., Principal of King William's College, Isle of Man. Cr. 8vo. 5s.

CONSTABLE.—GEOMETRICAL EXERCISES FOR BEGINNERS. By SAMUEL CONSTABLE. Cr. 8vo. 3s. 6d.

CUTHBERTSON.—EUCLIDIAN GEOMETRY. By FRANCIS CUTHBERTSON, M.A., LL.D. Ex. fcap. 8vo. 4s. 6d.

DAY.—PROPERTIES OF CONIC SECTIONS PROVED GEOMETRICALLY. By Rev. H. G. DAY, M.A. Part I. The Ellipse, with an ample collection of Problems. Cr. 8vo. 3s. 6d.

DEAKIN.—RIDER PAPERS ON EUCLID, BOOKS I. AND II. By RUPERT DEAKIN, M.A. 18mo. 1s.

DODGSON.—Works by CHARLES L. DODGSON, M.A., Student and late Mathematical Lecturer, Christ Church, Oxford.

EUCLID, BOOKS I. AND II. 6th Ed., with words substituted for the Algebraical Symbols used in the 1st Ed. Cr. 8vo. 2s.

EUCLID AND HIS MODERN RIVALS. 2d Ed. Cr. 8vo. 6s.

CURIOSA MATHEMATICA. Part I. A New Theory of Parallels. 3d Ed. Cr. 8vo. 2s.

DREW.—GEOMETRICAL TREATISE ON CONIC SECTIONS. By W. H. DREW, M.A. New Ed., enlarged. Cr. 8vo. 5s.

DUPUIS.—ELEMENTARY SYNTHETIC GEOMETRY OF THE POINT, LINE AND CIRCLE IN THE PLANE. By N. F. DUPUIS, M.A., Professor of Pure Mathematics in the University of Queen's College, Kingston, Canada. Gl. 8vo. 4s. 6d.

*HALL and STEVENS.—A TEXT-BOOK OF EUCLID'S ELEMENTS. Including Alternative Proofs, together with additional Theorems and Exercises, classified and arranged. By H. S. HALL, M.A., and F. H. STEVENS, M.A., Masters of the Military and Engineering Side, Clifton College. Gl. 8vo. Book I., 1s.; Books I. and II., 1s. 6d.; Books I.–IV., 3s.; Books III.–IV., 2s.; Books III–VI., 3s.; Books V.–VI. and XI., 2s. 6d.; Books I.–VI. and XI., 4s. 6d.; Book XI., 1s. [KEY. *In preparation.*

HALSTED.—THE ELEMENTS OF GEOMETRY. By G. B. HALSTED, Professor of Pure and Applied Mathematics in the University of Texas. 8vo. 12s. 6d.

HAYWARD.—THE ELEMENTS OF SOLID GEOMETRY. By R. B. HAYWARD, M.A., F.R.S. Gl. 8vo. 3s.

LOCK.—EUCLID FOR BEGINNERS. Being an Introduction to existing Text-Books. By Rev. J. B. LOCK, M.A. [*In the Press.*

MILNE and DAVIS.—GEOMETRICAL CONICS. Part I. The Parabola. By Rev. J. J. MILNE, M.A., and R. F. DAVIS, M.A. Cr. 8vo. 2s.

RICHARDSON.—THE PROGRESSIVE EUCLID. Books I. and II. With Notes, Exercises, and Deductions. Edited by A. T. RICHARDSON, M.A., Senior Mathematical Master at the Isle of Wight College. Gl. 8vo. 2s. 6d.

SYLLABUS OF PLANE GEOMETRY (corresponding to Euclid, Books I.–VI.)—Prepared by the Association for the Improvement of Geometrical Teaching. Cr. 8vo. Sewed, 1s.

SYLLABUS OF MODERN PLANE GEOMETRY.—Prepared by the Association for the Improvement of Geometrical Teaching. Cr. 8vo. Sewed. 1s.

*TODHUNTER.—THE ELEMENTS OF EUCLID. By I. TODHUNTER, F.R.S. 18mo. 3s. 6d. Books I. and II. 1s. KEY. Cr. 8vo. 6s. 6d.

WILSON.—Works by Ven. Archdeacon WILSON, M.A., formerly Headmaster of Clifton College.

ELEMENTARY GEOMETRY. BOOKS I.–V. Containing the Subjects of Euclid's first Six Books. Following the Syllabus of the Geometrical Association. Ex. fcap. 8vo. 4s. 6d.

WILSON.—Works by Ven. Archdeacon WILSON—*continued.*

SOLID GEOMETRY AND CONIC SECTIONS. With Appendices on Transversals and Harmonic Division. Ex. fcap. 8vo. 3s. 6d.

GEOMETRICAL DRAWING.

EAGLES.—CONSTRUCTIVE GEOMETRY OF PLANE CURVES. By T. H. EAGLES, M.A., Instructor in Geometrical Drawing and Lecturer in Architecture at the Royal Indian Engineering College, Cooper's Hill. Cr. 8vo. 12s.

EDGAR and PRITCHARD.—NOTE-BOOK ON PRACTICAL SOLID OR DESCRIPTIVE GEOMETRY. Containing Problems with help for Solutions. By J. H. EDGAR and G. S. PRITCHARD. 4th Ed., revised by A. MEEZE. Gl. 8vo. 4s. 6d.

***KITCHENER.**—A GEOMETRICAL NOTE-BOOK. Containing Easy Problems in Geometrical Drawing preparatory to the Study of Geometry. For the Use of Schools. By F. E. KITCHENER, M.A., Headmaster of the Newcastle-under-Lyme High School. 4to. 2s.

MILLAR.—ELEMENTS OF DESCRIPTIVE GEOMETRY. By J. B. MILLAR, Civil Engineer, Lecturer on Engineering in the Victoria University, Manchester. 2d Ed. Cr. 8vo. 6s.

PLANT.—PRACTICAL PLANE AND DESCRIPTIVE GEOMETRY. By E. C. PLANT. Globe 8vo. [*In preparation.*

MENSURATION.

STEVENS.—ELEMENTARY MENSURATION. With Exercises on the Mensuration of Plane and Solid Figures. By F. H. STEVENS, M.A. Gl. 8vo.

[*In preparation.*

TEBAY.—ELEMENTARY MENSURATION FOR SCHOOLS. By S. TEBAY. Ex. fcap. 8vo. 3s. 6d.

***TODHUNTER.**—MENSURATION FOR BEGINNERS. By ISAAC TODHUNTER, F.R.S. 18mo. 2s. 6d. KEY. By Rev. FR. L. MCCARTHY. Cr. 8vo. 7s. 6d.

TRIGONOMETRY.

BEASLEY.—AN ELEMENTARY TREATISE ON PLANE TRIGONOMETRY. With Examples. By R. D. BEASLEY, M.A. 9th Ed., revised and enlarged. Cr. 8vo. 3s. 6d.

BOTTOMLEY.—FOUR-FIGURE MATHEMATICAL TABLES. Comprising Logarithmic and Trigonometrical Tables, and Tables of Squares, Square Roots, and Reciprocals. By J. T. BOTTOMLEY, M.A., Lecturer in Natural Philosophy in the University of Glasgow. 8vo. 2s. 6d.

HAYWARD.—THE ALGEBRA OF CO-PLANAR VECTORS AND TRIGONOMETRY. By R. B. HAYWARD, M.A., F.R.S., Assistant Master at Harrow.

[*In preparation.*

JOHNSON.—A TREATISE ON TRIGONOMETRY. By W. E. JOHNSON, M.A., late Scholar and Assistant Mathematical Lecturer at King's College, Cambridge. Cr. 8vo. 3s. 6d.

LEVETT and DAVISON.—ELEMENTS OF TRIGONOMETRY. By RAWDON LEVETT and CHARLES DAVISON, Assistant Masters at King Edward's School, Birmingham. [*In the Press.*

LOCK.—Works by Rev. J. B. LOCK, M.A., Senior Fellow and Bursar of Gonville and Caius College, Cambridge.

*TRIGONOMETRY FOR BEGINNERS, as far as the Solution of Triangles. 3d Ed. Gl. 8vo. 2s. 6d. KEY. Cr. 8vo. 6s. 6d.

*ELEMENTARY TRIGONOMETRY. 6th Ed. (in this edition the chapter on logarithms has been carefully revised). Gl. 8vo. 4s. 6d. KEY. Cr. 8vo. 8s. 6d.

HIGHER TRIGONOMETRY. 5th Ed. Gl. 8vo. 4s. 6d. Both Parts complete in One Volume. Gl. 8vo. 7s. 6d.

THE TRIGONOMETRY OF ONE ANGLE. Gl. 8vo. 2s. 6d.

M'CLELLAND and PRESTON.—A TREATISE ON SPHERICAL TRIGONOMETRY. With applications to Spherical Geometry and numerous Examples. By W. J. M'Clelland, M.A., Principal of the Incorporated Society's School, Santry, Dublin, and T. Preston, M.A. Cr. 8vo. 8s. 6d., or: Part I. To the End of Solution of Triangles, 4s. 6d. Part II., 5s.

MATTHEWS.—MANUAL OF LOGARITHMS. By G. F. Matthews, B.A. 8vo. 5s. net.

PALMER.—TEXT-BOOK OF PRACTICAL LOGARITHMS AND TRIGONOMETRY. By J. H. Palmer, Headmaster, R.N., H.M.S. *Cambridge*, Devonport. Gl. 8vo. 4s. 6d.

SNOWBALL.—THE ELEMENTS OF PLANE AND SPHERICAL TRIGONOMETRY. By J. C. Snowball. 14th Ed. Cr. 8vo. 7s. 6d.

TODHUNTER.—Works by Isaac Todhunter, F.R.S.

*TRIGONOMETRY FOR BEGINNERS. 18mo. 2s. 6d. KEY. Cr. 8vo. 8s. 6d.

PLANE TRIGONOMETRY. Cr. 8vo. 5s. KEY. Cr. 8vo. 10s. 6d.
[A New Edition, revised by R. W. Hogg, M.A. *In the Press.*

A TREATISE ON SPHERICAL TRIGONOMETRY. Cr. 8vo. 4s. 6d.

WOLSTENHOLME.—EXAMPLES FOR PRACTICE IN THE USE OF SEVEN-FIGURE LOGARITHMS. By Joseph Wolstenholme, D.Sc., late Professor of Mathematics in the Royal Indian Engineering Coll., Cooper's Hill. 8vo. 5s.

ANALYTICAL GEOMETRY (Plane and Solid).

DYER.—EXERCISES IN ANALYTICAL GEOMETRY. By J. M. Dyer, M.A., Assistant Master at Eton. Illustrated. Cr. 8vo. 4s. 6d.

FERRERS.—AN ELEMENTARY TREATISE ON TRILINEAR CO-ORDINATES, the Method of Reciprocal Polars, and the Theory of Projectors. By the Rev. N. M. Ferrers, D.D., F.R.S., Master of Gonville and Caius College, Cambridge. 4th Ed., revised. Cr. 8vo. 6s. 6d.

FROST.—Works by Percival Frost, D.Sc., F.R.S., Fellow and Mathematical Lecturer at King's College, Cambridge.

AN ELEMENTARY TREATISE ON CURVE TRACING. 8vo. 12s.

SOLID GEOMETRY. 3d Ed. Demy 8vo. 16s.

HINTS FOR THE SOLUTION OF PROBLEMS in the Third Edition of SOLID GEOMETRY. 8vo. 8s. 6d.

JOHNSON.—CURVE TRACING IN CARTESIAN CO-ORDINATES. By W. Woolsey Johnson, Professor of Mathematics at the U.S. Naval Academy, Annapolis, Maryland. Cr. 8vo. 4s. 6d.

M'CLELLAND.—THE GEOMETRY OF THE CIRCLE. By W J. M'Clelland, M.A. Cr. 8vo. [*In the Press.*

PUCKLE.—AN ELEMENTARY TREATISE ON CONIC SECTIONS AND ALGEBRAIC GEOMETRY. With Numerous Examples and Hints for their Solution. By G. H. Puckle, M.A. 5th Ed., revised and enlarged. Cr. 8vo. 7s. 6d.

SMITH.—Works by Charles Smith, M.A., Master of Sidney Sussex College, Cambridge.

CONIC SECTIONS. 7th Ed. Cr. 8vo. 7s. 6d.

SOLUTIONS TO CONIC SECTIONS. Cr. 8vo. 10s. 6d.

AN ELEMENTARY TREATISE ON SOLID GEOMETRY. 2d Ed. Cr. 8vo. 9s. 6d.

TODHUNTER.—Works by Isaac Todhunter, F.R.S.

PLANE CO-ORDINATE GEOMETRY, as applied to the Straight Line and the Conic Sections. Cr. 8vo. 7s. 6d.

KEY. By C. W. Bourne, M.A., Headmaster of King's College School. Cr. 8vo. 10s. 6d.

EXAMPLES OF ANALYTICAL GEOMETRY OF THREE DIMENSIONS. New Ed., revised. Cr. 8vo. 4s.

PROBLEMS AND QUESTIONS IN MATHEMATICS.

ARMY PRELIMINARY EXAMINATION, 1882-1889, Specimens of Papers set at the. With Answers to the Mathematical Questions. Subjects: Arithmetic, Algebra, Euclid, Geometrical Drawing, Geography, French, English Dictation. Cr. 8vo. 3s. 6d.

CAMBRIDGE SENATE-HOUSE PROBLEMS AND RIDERS, WITH SOLUTIONS:—

1875—PROBLEMS AND RIDERS. By A. G. GREENHILL, F.R.S. Cr. 8vo. 8s. 6d.

1878—SOLUTIONS OF SENATE-HOUSE PROBLEMS. By the Mathematical Moderators and Examiners. Edited by J. W. L. GLAISHER, F.R.S., Fellow of Trinity College, Cambridge. 12s.

CHRISTIE.—A COLLECTION OF ELEMENTARY TEST-QUESTIONS IN PURE AND MIXED MATHEMATICS; with Answers and Appendices on Synthetic Division, and on the Solution of Numerical Equations by Horner's Method. By JAMES R. CHRISTIE, F.R.S. Cr. 8vo. 8s. 6d.

CLIFFORD.—MATHEMATICAL PAPERS. By W. K. CLIFFORD. Edited by R. TUCKER. With an Introduction by H. J. STEPHEN SMITH, M.A. 8vo. 30s.

MILNE.—Works by Rev. JOHN J. MILNE, Private Tutor.

WEEKLY PROBLEM PAPERS. With Notes intended for the use of Students preparing for Mathematical Scholarships, and for Junior Members of the Universities who are reading for Mathematical Honours. Pott 8vo. 4s. 6d.

SOLUTIONS TO WEEKLY PROBLEM PAPERS. Cr. 8vo. 10s. 6d.

COMPANION TO WEEKLY PROBLEM PAPERS. Cr. 8vo. 10s. 6d.

SANDHURST MATHEMATICAL PAPERS, for admission into the Royal Military College, 1881-1889. Edited by E. J. BROOKSMITH, B.A., Instructor in Mathematics at the Royal Military Academy, Woolwich. Cr. 8vo. 3s. 6d.

WOOLWICH MATHEMATICAL PAPERS, for Admission into the Royal Military Academy, Woolwich, 1880–1888 inclusive. Edited by E. J. BROOKSMITH, B.A. Cr. 8vo. 6s.

WOLSTENHOLME.—Works by JOSEPH WOLSTENHOLME, D.Sc., late Professor of Mathematics in the Royal Engineering Coll., Cooper's Hill.

MATHEMATICAL PROBLEMS, on Subjects included in the First and Second Divisions of the Schedule of Subjects for the Cambridge Mathematical Tripos Examination. New Ed., greatly enlarged. 8vo. 18s.

EXAMPLES FOR PRACTICE IN THE USE OF SEVEN-FIGURE LOGARITHMS. 8vo. 5s.

HIGHER PURE MATHEMATICS.

AIRY.—Works by Sir G. B. AIRY, K.C.B., formerly Astronomer-Royal.

ELEMENTARY TREATISE ON PARTIAL DIFFERENTIAL EQUATIONS. With Diagrams. 2d Ed. Cr. 8vo. 5s. 6d.

ON THE ALGEBRAICAL AND NUMERICAL THEORY OF ERRORS OF OBSERVATIONS AND THE COMBINATION OF OBSERVATIONS. 2d Ed., revised. Cr. 8vo. 6s. 6d.

BOOLE.—THE CALCULUS OF FINITE DIFFERENCES. By G. BOOLE. 3d Ed., revised by J. F. MOULTON, Q.C. Cr. 8vo. 10s. 6d.

CARLL.—A TREATISE ON THE CALCULUS OF VARIATIONS. By LEWIS B. CARLL. Arranged with the purpose of Introducing, as well as Illustrating, its Principles to the Reader by means of Problems, and Designed to present in all Important Particulars a Complete View of the Present State of the Science. 8vo. 21s.

EDWARDS.—THE DIFFERENTIAL CALCULUS. By JOSEPH EDWARDS, M.A. With Applications and numerous Examples. Cr. 8vo. 10s. 6d.

FERRERS.—AN ELEMENTARY TREATISE ON SPHERICAL HARMONICS, AND SUBJECTS CONNECTED WITH THEM. By Rev. N. M. FERRERS, D.D., F.R.S., Master of Gonville and Caius College, Cambridge. Cr. 8vo. 7s. 6d.

FORSYTH.—A TREATISE ON DIFFERENTIAL EQUATIONS. By Andrew Russell Forsyth, F.R.S., Fellow and Assistant Tutor of Trinity College, Cambridge. 2d Ed. 8vo. 14s.

FROST.—AN ELEMENTARY TREATISE ON CURVE TRACING. By Percival Frost, M.A., D.Sc. 8vo. 12s.

GRAHAM.—GEOMETRY OF POSITION. By R. H. Graham. Cr. 8vo. 7s. 6d.

GREENHILL.—DIFFERENTIAL AND INTEGRAL CALCULUS. By A. G. Greenhill, Professor of Mathematics to the Senior Class of Artillery Officers, Woolwich. New Ed. Cr. 8vo. 10s. 6d.

APPLICATIONS OF ELLIPTIC FUNCTIONS. By the same. [*In the Press.*

JOHNSON.—Works by William Woolsey Johnson, Professor of Mathematics at the U.S. Naval Academy, Annapolis, Maryland.

INTEGRAL CALCULUS, an Elementary Treatise on the. Founded on the Method of Rates or Fluxions. 8vo. 9s.

CURVE TRACING IN CARTESIAN CO-ORDINATES. Cr. 8vo. 4s. 6d.

A TREATISE ON ORDINARY AND DIFFERENTIAL EQUATIONS. Ex. cr. 8vo. 15s.

KELLAND and TAIT.—INTRODUCTION TO QUATERNIONS, with numerous examples. By P. Kelland and P. G. Tait, Professors in the Department of Mathematics in the University of Edinburgh. 2d Ed. Cr. 8vo. 7s. 6d.

KEMPE.—HOW TO DRAW A STRAIGHT LINE: a Lecture on Linkages. By A. B. Kempe. Illustrated. Cr. 8vo. 1s. 6d.

KNOX.—DIFFERENTIAL CALCULUS FOR BEGINNERS. By Alexander Knox. Fcap. 8vo. 3s. 6d.

MUIR.—THE THEORY OF DETERMINANTS IN THE HISTORICAL ORDER OF ITS DEVELOPMENT. Part I. Determinants in General. Leibnitz (1693) to Cayley (1841). By Thos. Muir, Mathematical Master in the High School of Glasgow. 8vo. 10s. 6d.

RICE and JOHNSON.—DIFFERENTIAL CALCULUS, an Elementary Treatise on the. Founded on the Method of Rates or Fluxions. By J. M. Rice, Professor of Mathematics in the United States Navy, and W. W. Johnson, Professor of Mathematics at the United States Naval Academy. 3d Ed., revised and corrected. 8vo. 18s. Abridged Ed. 9s.

TODHUNTER.—Works by Isaac Todhunter, F.R.S.

AN ELEMENTARY TREATISE ON THE THEORY OF EQUATIONS. Cr. 8vo. 7s. 6d.

A TREATISE ON THE DIFFERENTIAL CALCULUS. Cr. 8vo. 10s. 6d. KEY. Cr. 8vo. 10s. 6d.

A TREATISE ON THE INTEGRAL CALCULUS AND ITS APPLICATIONS. Cr. 8vo. 10s. 6d. KEY. Cr. 8vo. 10s. 6d.

A HISTORY OF THE MATHEMATICAL THEORY OF PROBABILITY, from the time of Pascal to that of Laplace. 8vo. 18s.

AN ELEMENTARY TREATISE ON LAPLACE'S, LAME'S, AND BESSEL'S FUNCTIONS. Cr. 8vo. 10s. 6d.

MECHANICS: Statics, Dynamics, Hydrostatics, Hydrodynamics. (See also Physics.)

ALEXANDER and THOMSON.—ELEMENTARY APPLIED MECHANICS. By Prof. T. Alexander and A. W. Thomson. Part II. Transverse Stress. Cr. 8vo. 10s. 6d.

BALL.—EXPERIMENTAL MECHANICS. A Course of Lectures delivered at the Royal College of Science for Ireland. By Sir R. S. Ball, F.R.S. 2d Ed. Illustrated. Cr. 8vo. 6s.

CLIFFORD.—THE ELEMENTS OF DYNAMIC. An Introduction to the Study of Motion and Rest in Solid and Fluid Bodies. By W. K. Clifford. Part I.—Kinematic. Cr. 8vo. Books I.-III. 7s. 6d.; Book IV. and Appendix, 6s.

COTTERILL.—APPLIED MECHANICS: An Elementary General Introduction to the Theory of Structures and Machines. By J. H. COTTERILL, F.R.S., Professor of Applied Mechanics in the Royal Naval College, Greenwich. 8vo. 18s.

COTTERILL and SLADE.—LESSONS IN APPLIED MECHANICS. By Prof. J. H. COTTERILL and J. H. SLADE. Fcap. 8vo. 5s. 6d.

DYNAMICS, SYLLABUS OF ELEMENTARY. Part I. Linear Dynamics. With an Appendix on the Meanings of the Symbols in Physical Equations. Prepared by the Association for the Improvement of Geometrical Teaching. 4to. 1s.

GANGUILLET and KUTTER.—A GENERAL FORMULA FOR THE UNIFORM FLOW OF WATER IN RIVERS AND OTHER CHANNELS. By E. GANGUILLET and W. R. KUTTER, Engineers in Berne, Switzerland. Translated from the German, with numerous Additions, including Tables and Diagrams, and the Elements of over 1200 Gaugings of Rivers, Small Channels, and Pipes in English Measure, by RUDOLPH HERING, Assoc. Am. Soc. C.E., M. Inst. C.E., and JOHN C. TRAUTWINE Jun., Assoc. Am. Soc. C.E., Assoc. Inst. C.E. 8vo. 17s.

GRAHAM.—GEOMETRY OF POSITION. By R. H. GRAHAM. Cr. 8vo. 7s. 6d.

GREAVES.—Works by JOHN GREAVES, M.A., Fellow and Mathematical Lecturer at Christ's College, Cambridge.

*STATICS FOR BEGINNERS. Gl. 8vo. 8s. 6d.

A TREATISE ON ELEMENTARY STATICS. 2d Ed. Cr. 8vo. 6s. 6d.

GREENHILL.—HYDROSTATICS. By A. G. GREENHILL, Professor of Mathematics to the Senior Class of Artillery Officers, Woolwich. Cr. 8vo. [*In preparation.*

***HICKS.**—ELEMENTARY DYNAMICS OF PARTICLES AND SOLIDS. By W. M. HICKS, Principal and Professor of Mathematics and Physics, Firth College, Sheffield. Cr. 8vo. 6s. 6d.

JELLETT.—A TREATISE ON THE THEORY OF FRICTION. By JOHN H. JELLETT, B.D., late Provost of Trinity College, Dublin. 8vo. 8s. 6d.

KENNEDY.—THE MECHANICS OF MACHINERY. By A. B. W. KENNEDY, F.R.S. Illustrated. Cr. 8vo. 12s. 6d.

LOCK.—Works by Rev. J. B. LOCK, M.A.

*ELEMENTARY STATICS. 2d Ed. Gl. 8vo. 4s. 6d.

*ELEMENTARY DYNAMICS. 3d Ed. Gl. 8vo. 4s. 6d.

MECHANICS FOR BEGINNERS. Gl. 8vo. [*In the Press.*

MACGREGOR.—KINEMATICS AND DYNAMICS. An Elementary Treatise. By J. G. MACGREGOR, D.Sc., Munro Professor of Physics in Dalhousie College, Halifax, Nova Scotia. Illustrated. Cr. 8vo. 10s. 6d.

PARKINSON.—AN ELEMENTARY TREATISE ON MECHANICS. By S. PARKINSON, D.D., F.R.S., late Tutor and Prælector of St. John's College, Cambridge. 6th Ed., revised. Cr. 8vo. 9s. 6d.

PIRIE.—LESSONS ON RIGID DYNAMICS. By Rev. G. PIRIE, M.A., Professor of Mathematics in the University of Aberdeen. Cr. 8vo. 6s.

ROUTH.—Works by EDWARD JOHN ROUTH, D.Sc., LL.D., F.R.S., Hon. Fellow of St. Peter's College, Cambridge.

A TREATISE ON THE DYNAMICS OF THE SYSTEM OF RIGID BODIES. With numerous Examples. Two Vols. 8vo. Vol. I.—Elementary Parts. 5th Ed. 14s. Vol. II.—The Advanced Parts. 4th Ed. 14s.

STABILITY OF A GIVEN STATE OF MOTION, PARTICULARLY STEADY MOTION. Adams Prize Essay for 1877. 8vo. 8s. 6d.

***SANDERSON.**—HYDROSTATICS FOR BEGINNERS. By F. W. SANDERSON, M.A., Assistant Master at Dulwich College. Gl. 8vo. 4s. 6d.

TAIT and STEELE.—A TREATISE ON DYNAMICS OF A PARTICLE. By Professor TAIT, M.A., and W. J. STEELE, B.A. 6th Ed., revised. Cr. 8vo. 12s.

TODHUNTER.—Works by ISAAC TODHUNTER, F.R.S.

*MECHANICS FOR BEGINNERS. 18mo. 4s. 6d. KEY. Cr. 8vo. 6s. 6d.

A TREATISE ON ANALYTICAL STATICS. 5th Ed. Edited by Prof. J. D. EVERETT, F.R.S. Cr. 8vo. 10s. 6d.

PHYSICS: Sound, Light, Heat, Electricity, Elasticity, Attractions, etc. (See also Mechanics.)

AIRY.—Works by Sir G. B. AIRY, K.C.B., formerly Astronomer-Royal.

ON SOUND AND ATMOSPHERIC VIBRATIONS. With the Mathematical Elements of Music. 2d Ed., revised and enlarged. Cr. 8vo. 9s.

GRAVITATION: An Elementary Explanation of the Principal Perturbations in the Solar System. 2d Ed. Cr. 8vo. 7s. 6d.

CLAUSIUS.—MECHANICAL THEORY OF HEAT. By R. CLAUSIUS. Translated by W. R. BROWNE, M.A. Cr. 8vo. 10s. 6d.

CUMMING.—AN INTRODUCTION TO THE THEORY OF ELECTRICITY. By LINNÆUS CUMMING, M.A., Assistant Master at Rugby. Illustrated. Cr. 8vo. 8s. 6d.

DANIELL.—A TEXT-BOOK OF THE PRINCIPLES OF PHYSICS. By ALFRED DANIELL, D.Sc. Illustrated. 2d Ed., revised and enlarged. 8vo. 21s.

DAY.—ELECTRIC LIGHT ARITHMETIC. By R. E. DAY, Evening Lecturer in Experimental Physics at King's College, London. Pott 8vo. 2s.

EVERETT.—UNITS AND PHYSICAL CONSTANTS. By J. D. EVERETT, F.R.S., Professor of Natural Philosophy, Queen's College, Belfast. 2d Ed. Ex. fcap. 8vo. 5s.

FERRERS.—AN ELEMENTARY TREATISE ON SPHERICAL HARMONICS, and Subjects connected with them. By Rev. N. M. FERRERS, D.D., F.R.S., Master of Gonville and Caius College, Cambridge. Cr. 8vo. 7s. 6d.

FESSENDEN.—PHYSICS FOR PUBLIC SCHOOLS. By C. FESSENDEN. Illustrated. Fcap. 8vo. [*In the Press.*

GRAY.—THE THEORY AND PRACTICE OF ABSOLUTE MEASUREMENTS IN ELECTRICITY AND MAGNETISM. By A. GRAY, F.R.S.E., Professor of Physics in the University College of North Wales. Two Vols. Cr. 8vo. Vol. I. 12s. 6d. [Vol. II. *In the Press.*

ABSOLUTE MEASUREMENTS IN ELECTRICITY AND MAGNETISM. 2d Ed., revised and greatly enlarged. Fcap. 8vo. 5s. 6d.

IBBETSON.—THE MATHEMATICAL THEORY OF PERFECTLY ELASTIC SOLIDS, with a Short Account of Viscous Fluids. By W. J. IBBETSON, late Senior Scholar of Clare College, Cambridge. 8vo. 21s.

***JONES.**—EXAMPLES IN PHYSICS. Containing over 1000 Problems with Answers and numerous solved Examples. Suitable for candidates preparing for the Intermediate, Science, Preliminary, Scientific, and other Examinations of the University of London. By D. E. JONES, B.Sc., Professor of Physics in the University College of Wales, Aberystwyth. Fcap. 8vo. 3s. 6d.

***ELEMENTARY LESSONS IN HEAT, LIGHT, AND SOUND.** Gl. 8vo. 2s. 6d.

LOCKYER.—CONTRIBUTIONS TO SOLAR PHYSICS. By J. NORMAN LOCKYER, F.R.S. With Illustrations. Royal 8vo. 31s. 6d.

LODGE.—MODERN VIEWS OF ELECTRICITY. By OLIVER J. LODGE, F.R.S., Professor of Experimental Physics in University College, Liverpool. Illustrated. Cr. 8vo. 6s. 6d.

LOEWY.—*QUESTIONS AND EXAMPLES ON EXPERIMENTAL PHYSICS: Sound, Light, Heat, Electricity, and Magnetism. By B. LOEWY, Examiner in Experimental Physics to the College of Preceptors. Fcap. 8vo. 2s.

*A GRADUATED COURSE OF NATURAL SCIENCE FOR ELEMENTARY AND TECHNICAL SCHOOLS AND COLLEGES. By the same. In Three Parts. Part I. FIRST YEAR'S COURSE. Gl. 8vo. 2s.

LUPTON.—NUMERICAL TABLES AND CONSTANTS IN ELEMENTARY SCIENCE. By S. LUPTON, M.A., late Assistant Master at Harrow. Ex. fcap. 8vo. 2s. 6d.

MACFARLANE.—PHYSICAL ARITHMETIC. By A. MACFARLANE, D.Sc., late Examiner in Mathematics at the University of Edinburgh. Cr. 8vo. 7s. 6d.

***MAYER.**—SOUND: A Series of Simple, Entertaining, and Inexpensive Experiments in the Phenomena of Sound. By A. M. Mayer, Professor of Physics in the Stevens Institute of Technology. Illustrated. Cr. 8vo. 3s. 6d.

***MAYER and BARNARD.**—LIGHT: A Series of Simple, Entertaining, and Inexpensive Experiments in the Phenomena of Light. By A. M. Mayer and C. Barnard. Illustrated. Cr. 8vo. 2s. 6d.

MOLLOY.—GLEANINGS IN SCIENCE: Popular Lectures on Scientific Subjects. By the Rev. Gerald Molloy, D.Sc., Rector of the Catholic University of Ireland. 8vo. 7s. 6d.

NEWTON.—PRINCIPIA. Edited by Prof. Sir W. Thomson, P.R.S., and Prof. Blackburne. 4to. 31s. 6d.

THE FIRST THREE SECTIONS OF NEWTON'S PRINCIPIA. With Notes and Illustrations. Also a Collection of Problems, principally intended as Examples of Newton's Methods. By P. Frost, M.A., D.Sc. 3d Ed. 8vo. 12s.

PARKINSON.—A TREATISE ON OPTICS. By S. Parkinson, D.D., F.R.S., late Tutor and Prælector of St. John's College, Cambridge. 4th Ed., revised and enlarged. Cr. 8vo. 10s. 6d.

PEABODY.—THERMODYNAMICS OF THE STEAM-ENGINE AND OTHER HEAT-ENGINES. By Cecil H. Peabody, Associate Professor of Steam Engineering, Massachusetts Institute of Technology. 8vo. 21s.

PERRY.—STEAM: An Elementary Treatise. By John Perry, Professor of Mechanical Engineering and Applied Mechanics at the Technical College, Finsbury. 18mo. 4s. 6d.

PICKERING.—ELEMENTS OF PHYSICAL MANIPULATION. By Prof. Edward C. Pickering. Medium 8vo. Part I., 12s. 6d. Part II., 14s.

PRESTON.—THE THEORY OF LIGHT. By Thomas Preston, M.A. Illustrated. 8vo. 12s. 6d.

THE THEORY OF HEAT. By the same Author. 8vo. [*In preparation.*

RAYLEIGH.—THE THEORY OF SOUND. By Lord Rayleigh, F.R.S. 8vo. Vol. I., 12s. 6d. Vol. II., 12s. 6d. [Vol. III. *In the Press.*

SHANN.—AN ELEMENTARY TREATISE ON HEAT, IN RELATION TO STEAM AND THE STEAM-ENGINE. By G. Shann, M.A. Illustrated. Cr. 8vo. 4s. 6d.

SPOTTISWOODE.—POLARISATION OF LIGHT. By the late W. Spottiswoode, F.R.S. Illustrated. Cr. 8vo. 3s. 6d.

STEWART.—Works by Balfour Stewart, F.R.S., late Langworthy Professor of Physics in the Owens College, Victoria University, Manchester.

*PRIMER OF PHYSICS. Illustrated. With Questions. 18mo. 1s.

*LESSONS IN ELEMENTARY PHYSICS. Illustrated. Fcap. 8vo. 4s. 6d.

*QUESTIONS. By Prof. T. H. Core. Fcap. 8vo. 2s.

STEWART and GEE.—LESSONS IN ELEMENTARY PRACTICAL PHYSICS. By Balfour Stewart, F.R.S., and W. W. Haldane Gee, B.Sc. Cr. 8vo. Vol. I. General Physical Processes. 6s. Vol. II. Electricity and Magnetism. 7s. 6d. [Vol. III. Optics, Heat, and Sound. *In the Press.*

*PRACTICAL PHYSICS FOR SCHOOLS AND THE JUNIOR STUDENTS OF COLLEGES. Gl. 8vo. Vol. I. Electricity and Magnetism. 2s. 6d. [Vol. II. Optics, Heat, and Sound. *In the Press.*

STOKES.—ON LIGHT. Burnett Lectures, delivered in Aberdeen in 1883-4-5. By Sir G. G. Stokes, F.R.S., Lucasian Professor of Mathematics in the University of Cambridge. First Course: On the Nature of Light. Second Course: On Light as a Means of Investigation. Third Course: On the Beneficial Effects of Light. Cr. 8vo. 7s. 6d.

*** The 2d and 3d Courses may be had separately. Cr. 8vo. 2s. 6d. each.

STONE.—AN ELEMENTARY TREATISE ON SOUND. By W. H. Stone. Illustrated. Fcap. 8vo. 3s. 6d.

TAIT.—HEAT. By P. G. Tait, Professor of Natural Philosophy in the University of Edinburgh. Cr. 8vo. 6s.

LECTURES ON SOME RECENT ADVANCES IN PHYSICAL SCIENCE. By the same. 3d Edition. Crown 8vo. 9s.

TAYLOR.—SOUND AND MUSIC. An Elementary Treatise on the Physical Constitution of Musical Sounds and Harmony, including the Chief Acoustical Discoveries of Professor Helmholtz. By SEDLEY TAYLOR, M.A. Illustrated. 2d Ed. Ex. cr. 8vo. 8s. 6d.

***THOMPSON.**—ELEMENTARY LESSONS IN ELECTRICITY AND MAGNETISM. By SILVANUS P. THOMPSON, Principal and Professor of Physics in the Technical College, Finsbury. Illustrated. New Ed., revised. Fcap. 8vo. 4s. 6d.

THOMSON.—Works by J. J. THOMSON, Professor of Experimental Physics in the University of Cambridge.

A TREATISE ON THE MOTION OF VORTEX RINGS. Adams Prize Essay, 1882. 8vo. 6s.

APPLICATIONS OF DYNAMICS TO PHYSICS AND CHEMISTRY. Cr. 8vo. 7s. 6d.

THOMSON.—Works by Sir W. THOMSON, P.R.S., Professor of Natural Philosophy in the University of Glasgow.

ELECTROSTATICS AND MAGNETISM, REPRINTS OF PAPERS ON. 2d Ed. 8vo. 18s.

POPULAR LECTURES AND ADDRESSES. 3 Vols. Illustrated. Cr. 8vo. Vol. I. CONSTITUTION OF MATTER. 6s. Vol. III. NAVIGATION. [*In the Press.*

TODHUNTER.—Works by ISAAC TODHUNTER, F.R.S.

AN ELEMENTARY TREATISE ON LAPLACE'S, LAME'S, AND BESSEL'S FUNCTIONS. Crown 8vo. 10s. 6d.

A HISTORY OF THE MATHEMATICAL THEORIES OF ATTRACTION, AND THE FIGURE OF THE EARTH, from the time of Newton to that of Laplace. 2 vols. 8vo. 24s.

TURNER.—A COLLECTION OF EXAMPLES ON HEAT AND ELECTRICITY. By H. H. TURNER, Fellow of Trinity College, Cambridge. Cr. 8vo. 2s. 6d.

WRIGHT.—LIGHT: A Course of Experimental Optics, chiefly with the Lantern. By LEWIS WRIGHT. Illustrated. Cr. 8vo. 7s. 6d.

ASTRONOMY.

AIRY.—Works by Sir G. B. AIRY, K.C.B., formerly Astronomer-Royal.

*POPULAR ASTRONOMY. 18mo. 4s. 6d.

GRAVITATION: An Elementary Explanation of the Principal Perturbations in the Solar System. 2d Ed. Cr. 8vo. 7s. 6d.

CHEYNE.—AN ELEMENTARY TREATISE ON THE PLANETARY THEORY. By C. H. H. CHEYNE. With Problems. 3d Ed. Edited by Rev. A. FREEMAN, M.A., F.R.A.S. Cr. 8vo. 7s. 6d.

CLARK and SADLER.—THE STAR GUIDE. By L. CLARK and H. SADLER. Roy. 8vo. 5s.

CROSSLEY, GLEDHILL, and WILSON.—A HANDBOOK OF DOUBLE STARS. By E. CROSSLEY, J. GLEDHILL, and J. M. WILSON. 8vo. 21s.

CORRECTIONS TO THE HANDBOOK OF DOUBLE STARS. 8vo. 1s.

FORBES.—TRANSIT OF VENUS. By G. FORBES, Professor of Natural Philosophy in the Andersonian University, Glasgow. Illustrated. Cr. 8vo. 3s. 6d.

GODFRAY.—Works by HUGH GODFRAY, M.A., Mathematical Lecturer at Pembroke College, Cambridge.

A TREATISE ON ASTRONOMY. 4th Ed. 8vo. 12s. 6d.

AN ELEMENTARY TREATISE ON THE LUNAR THEORY, with a brief Sketch of the Problem up to the time of Newton. 2d Ed., revised. Cr. 8vo. 5s. 6d.

LOCKYER.—Works by J. NORMAN LOCKYER, F.R.S.

*PRIMER OF ASTRONOMY. Illustrated. 18mo. 1s.

*ELEMENTARY LESSONS IN ASTRONOMY. With Spectra of the Sun, Stars, and Nebulæ, and numerous Illustrations. 36th Thousand. Revised throughout. Fcap. 8vo. 5s. 6d.

*QUESTIONS ON LOCKYER'S ELEMENTARY LESSONS IN ASTRONOMY. By J. FORBES ROBERTSON. 18mo. 1s. 6d.

THE CHEMISTRY OF THE SUN. Illustrated. 8vo. 14s.

THE METEORITIC HYPOTHESIS OF THE ORIGIN OF COSMICAL SYSTEMS. Illustrated. 8vo. 17s. net.

THE EVOLUTION OF THE HEAVENS AND THE EARTH. Cr. 8vo. Illustrated. [*In the Press.*

LOCKYER and SEABROKE.—STAR-GAZING PAST AND PRESENT. By J. NORMAN LOCKYER, F.R.S. Expanded from Shorthand Notes with the assistance of G. M. SEABROKE, F.R.A.S. Royal 8vo. 21s.

NEWCOMB.—POPULAR ASTRONOMY. By S. NEWCOMB, LL.D., Professor U.S. Naval Observatory. Illustrated. 2d Ed., revised. 8vo. 18s.

HISTORICAL.

BALL.—A SHORT ACCOUNT OF THE HISTORY OF MATHEMATICS. By W. W. R. BALL, M.A. Cr. 8vo. 10s. 6d.

TODHUNTER.—Works by ISAAC TODHUNTER, F.R.S.

A HISTORY OF THE MATHEMATICAL THEORY OF PROBABILITY, from the time of Pascal to that of Laplace. 8vo. 18s.

A HISTORY OF THE MATHEMATICAL THEORIES OF ATTRACTION, AND THE FIGURE OF THE EARTH, from the time of Newton to that of Laplace. 2 vols. 8vo. 24s.

NATURAL SCIENCES.

Chemistry; Physical Geography, Geology, and Mineralogy; Biology; Medicine.

(For MECHANICS, PHYSICS, and ASTRONOMY, see *MATHEMATICS.*)

CHEMISTRY.

ARMSTRONG.—A MANUAL OF INORGANIC CHEMISTRY. By HENRY ARMSTRONG, F.R.S., Professor of Chemistry in the City and Guilds of London Technical Institute. Cr. 8vo. [*In preparation.*

***COHEN.**—THE OWENS COLLEGE COURSE OF PRACTICAL ORGANIC CHEMISTRY. By JULIUS B. COHEN, Ph.D., Assistant Lecturer on Chemistry in the Owens College, Manchester. With a Preface by Sir HENRY ROSCOE, F.R.S., and C. SCHORLEMMER, F.R.S. Fcap. 8vo. 2s. 6d.

COOKE.—ELEMENTS OF CHEMICAL PHYSICS. By JOSIAH P. COOKE, Jun., Erving Professor of Chemistry and Mineralogy in Harvard University. 4th Ed. 8vo. 21s.

FLEISCHER.—A SYSTEM OF VOLUMETRIC ANALYSIS. By EMIL FLEISCHER. Translated, with Notes and Additions, by M. M. P. MUIR, F.R.S.E. Illustrated. Cr. 8vo. 7s. 6d.

FRANKLAND.—A HANDBOOK OF AGRICULTURAL CHEMICAL ANALYSIS. By P. F. FRANKLAND, F.R.S., Professor of Chemistry in University College, Dundee. Cr. 8vo. 7s. 6d.

HARTLEY.—A COURSE OF QUANTITATIVE ANALYSIS FOR STUDENTS. By W. NOEL HARTLEY, F.R.S., Professor of Chemistry and of Applied Chemistry, Science and Art Department, Royal College of Science, Dublin. Gl. 8vo. 5s.

HIORNS.—PRACTICAL METALLURGY AND ASSAYING. A Text-Book for the use of Teachers, Students, and Assayers. By ARTHUR H. HIORNS, Principal of the School of Metallurgy, Birmingham and Midland Institute. Illustrated. Gl. 8vo. 6s.

A TEXT-BOOK OF ELEMENTARY METALLURGY FOR THE USE OF STUDENTS. To which is added an Appendix of Examination Questions, embracing the whole of the Questions set in the three stages of the subject by the Science and Art Department for the past twenty years. By the same. Gl. 8vo. 4s.

IRON AND STEEL MANUFACTURE. A Text-Book for Beginners. By the same. Illustrated. Gl. 8vo. 3s. 6d.

MIXED METALS OR METALLIC ALLOYS. By the same. Gl. 8vo. 6s.

JONES.—*THE OWENS COLLEGE JUNIOR COURSE OF PRACTICAL CHEMISTRY. By Francis Jones, F.R.S.E., Chemical Master at the Grammar School, Manchester. With Preface by Sir Henry Roscoe, F.R.S. Illustrated. Fcap. 8vo. 2s. 6d.

*QUESTIONS ON CHEMISTRY. A Series of Problems and Exercises in Inorganic and Organic Chemistry. By the same. Fcap. 8vo. 3s.

LANDAUER.—BLOWPIPE ANALYSIS. By J. Landauer. Authorised English Edition by J. Taylor and W. E. Kay, of Owens College, Manchester.

[*New Edition in preparation.*

LOCKYER.—THE CHEMISTRY OF THE SUN. By J. Norman Lockyer, F.R.S. Illustrated. 8vo. 14s.

LUPTON.—CHEMICAL ARITHMETIC. With 1200 Problems. By S. Lupton, M.A. 2d Ed., revised and abridged. Fcap. 8vo. 4s. 6d.

MANSFIELD.—A THEORY OF SALTS. By C. B. Mansfield. Crown 8vo. 14s.

MELDOLA.—THE CHEMISTRY OF PHOTOGRAPHY. By Raphael Meldola, F.R.S., Professor of Chemistry in the Technical College, Finsbury. Cr. 8vo. 6s.

MEYER. HISTORY OF CHEMISTRY. By Ernst von Meyer. Translated by George McGowan, Ph.D. 8vo. 14s. net.

MIXTER.—AN ELEMENTARY TEXT-BOOK OF CHEMISTRY. By William G. Mixter, Professor of Chemistry in the Sheffield Scientific School of Yale College. 2d and revised Ed. Cr. 8vo. 7s. 6d.

MUIR.—PRACTICAL CHEMISTRY FOR MEDICAL STUDENTS. Specially arranged for the first M.B. Course. By M. M. P. Muir, F.R.S.E., Fellow and Prælector in Chemistry at Gonville and Caius College, Cambridge. Fcap. 8vo. 1s. 6d.

MUIR and WILSON.—THE ELEMENTS OF THERMAL CHEMISTRY. By M. M. P. Muir, F.R.S.E.; assisted by D. M. Wilson. 8vo. 12s. 6d.

OSTWALD.—OUTLINES OF GENERAL CHEMISTRY (PHYSICAL AND THEORETICAL). By Prof. W. Ostwald. Translated by James Walker, D.Sc., Ph.D. 8vo. 10s. net.

RAMSAY.—EXPERIMENTAL PROOFS OF CHEMICAL THEORY FOR BEGINNERS. By William Ramsay, F.R.S., Professor of Chemistry in University College, London. Pott 8vo. 2s. 6d.

REMSEN.—Works by Ira Remsen, Professor of Chemistry in the Johns Hopkins University, U.S.A.

COMPOUNDS OF CARBON: or, Organic Chemistry, an Introduction to the Study of. Cr. 8vo. 6s. 6d.

AN INTRODUCTION TO THE STUDY OF CHEMISTRY (INORGANIC CHEMISTRY). Cr. 8vo. 6s. 6d.

*THE ELEMENTS OF CHEMISTRY. A Text-Book for Beginners. Fcap. 8vo. 2s. 6d.

A TEXT-BOOK OF INORGANIC CHEMISTRY. 8vo. 16s.

ROSCOE.—Works by Sir Henry E. Roscoe, F.R.S., formerly Professor of Chemistry in the Owens College, Victoria University, Manchester.

*PRIMER OF CHEMISTRY. Illustrated. With Questions. 18mo. 1s.

*LESSONS IN ELEMENTARY CHEMISTRY, INORGANIC AND ORGANIC. With Illustrations and Chromolitho of the Solar Spectrum, and of the Alkalies and Alkaline Earths. Fcap. 8vo. 4s. 6d.

ROSCOE and SCHORLEMMER.—INORGANIC AND ORGANIC CHEMISTRY. A Complete Treatise on Inorganic and Organic Chemistry. By Sir Henry E. Roscoe, F.R.S., and Prof. C. Schorlemmer, F.R.S. Illustrated. 8vo.

Vols. I. and II. INORGANIC CHEMISTRY. Vol. I.—The Non-Metallic Elements. 2d Ed. 21s. Vol. II. Part I.—Metals. 18s. Part II.—Metals. 18s.

Vol. III.—ORGANIC CHEMISTRY. THE CHEMISTRY OF THE HYDROCARBONS and their Derivatives. Five Parts. Parts I., II., and IV. 21s. Parts III. and V. 18s. each.

ROSCOE and SCHUSTER.—SPECTRUM ANALYSIS. Lectures delivered in 1868. By Sir HENRY ROSCOE, F.R.S. 4th Ed., revised and considerably enlarged by the Author and by A. SCHUSTER, F.R.S., Ph.D., Professor of Applied Mathematics in the Owens College, Victoria University. With Appendices, Illustrations, and Plates. 8vo. 21s.

***THORPE.**—A SERIES OF CHEMICAL PROBLEMS. With Key. For use in Colleges and Schools. By T. E. THORPE, B.Sc. (Vic.), Ph.D., F.R.S. Revised and Enlarged by W. TATE, Assoc.N.S.S. With Preface by Sir H. E. ROSCOE, F.R.S. New Ed. Fcap. 8vo. 2s.

THORPE and RÜCKER.—A TREATISE ON CHEMICAL PHYSICS. By Prof. T. E. THORPE, F.R.S., and Prof. A. W. RÜCKER, F.R.S. Illustrated. 8vo. *[In preparation.*

WILLIAMS.—ELEMENTS OF CRYSTALLOGRAPHY FOR STUDENTS OF CHEMISTRY, PHYSICS, AND MINERALOGY. By G. H. WILLIAMS, Ph.D. Cr. 8vo.

WURTZ.—A HISTORY OF CHEMICAL THEORY. By AD. WURTZ. Translated by HENRY WATTS, F.R.S. Crown 8vo. 6s.

PHYSICAL GEOGRAPHY, GEOLOGY, AND MINERALOGY.

BLANFORD.—THE RUDIMENTS OF PHYSICAL GEOGRAPHY FOR THE USE OF INDIAN SCHOOLS; with a Glossary of Technical Terms employed. By H. F. BLANFORD, F.G.S. Illustrated. Cr. 8vo. 2s. 6d.

FERREL.—A POPULAR TREATISE ON THE WINDS. Comprising the General Motions of the Atmosphere, Monsoons, Cyclones, Tornadoes, Waterspouts, Hailstorms, etc. By WILLIAM FERREL, M.A., Member of the American National Academy of Sciences. 8vo. 18s.

FISHER.—PHYSICS OF THE EARTH'S CRUST. By the Rev. OSMOND FISHER, M.A., F.G.S., Hon. Fellow of King's College, London. 2d Ed., altered and enlarged. 8vo. 12s.

GEIKIE.—Works by ARCHIBALD GEIKIE, LL.D., F.R.S., Director-General of the Geological Survey of Great Britain and Ireland.

*PRIMER OF PHYSICAL GEOGRAPHY. Illustrated. With Questions. 18mo. 1s.

*ELEMENTARY LESSONS IN PHYSICAL GEOGRAPHY. Illustrated. Fcap. 8vo. 4s. 6d. *QUESTIONS ON THE SAME. 1s. 6d.

*PRIMER OF GEOLOGY. Illustrated. 18mo. 1s.

*CLASS-BOOK OF GEOLOGY. Illustrated. New and Cheaper Ed. Cr. 8vo. 4s. 6d.

TEXT-BOOK OF GEOLOGY. Illustrated. 2d Ed., 7th Thousand, revised and enlarged. 8vo. 28s.

OUTLINES OF FIELD GEOLOGY. Illustrated. Ex. fcap. 8vo. 3s. 6d.

THE SCENERY AND GEOLOGY OF SCOTLAND, VIEWED IN CONNEXION WITH ITS PHYSICAL GEOLOGY. Illustrated. Cr. 8vo. 12s. 6d.

HUXLEY.—PHYSIOGRAPHY. An Introduction to the Study of Nature. By T. H. HUXLEY, F.R.S. Illustrated. New and Cheaper Edition. Cr. 8vo. 6s.

LOCKYER.—OUTLINES OF PHYSIOGRAPHY—THE MOVEMENTS OF THE EARTH. By J. NORMAN LOCKYER, F.R.S., Examiner in Physiography for the Science and Art Department. Illustrated. Cr. 8vo. Sewed, 1s. 6d.

PHILLIPS. A TREATISE ON ORE DEPOSITS. By J. ARTHUR PHILLIPS, F.R.S. Illustrated. 8vo. 25s.

ROSENBUSCH and IDDINGS.—MICROSCOPICAL PHYSIOGRAPHY OF THE ROCK-MAKING MINERALS: AN AID TO THE MICROSCOPICAL STUDY OF ROCKS. By H. ROSENBUSCH. Translated and Abridged by J. P. IDDINGS. Illustrated. 8vo. 24s.

BIOLOGY.

ALLEN.—ON THE COLOURS OF FLOWERS, as Illustrated in the British Flora. By GRANT ALLEN. Illustrated. Cr. 8vo. 3s. 6d.

BALFOUR.—A TREATISE ON COMPARATIVE EMBRYOLOGY. By F. M. BALFOUR, F.R.S., Fellow and Lecturer of Trinity College, Cambridge. Illustrated. 2d Ed., reprinted without alteration from the 1st Ed. 2 vols. 8vo. Vol. I. 18s. Vol. II. 21s.

BALFOUR and WARD.—A GENERAL TEXT-BOOK OF BOTANY. By ISAAC BAYLEY BALFOUR, F.R.S., Professor of Botany in the University of Edinburgh, and H. MARSHALL WARD, F.R.S., Professor of Botany in the Royal Indian Engineering College, Cooper's Hill. 8vo. [*In preparation.*

***BETTANY.**—FIRST LESSONS IN PRACTICAL BOTANY. By G. T. BETTANY. 18mo. 1s.

***BOWER.**—A COURSE OF PRACTICAL INSTRUCTION IN BOTANY. By F. O. BOWER, D.Sc., Regius Professor of Botany in the University of Glasgow. Cr. 8vo. 10s. 6d.

BUCKTON.—MONOGRAPH OF THE BRITISH CICADÆ, OR TETTIGIDÆ. By G. B. BUCKTON. In 8 parts, Quarterly. Part I. January, 1890. 8vo. Parts I.-V. ready. 8s. each, net. Vol I. 33s. 6d. net.

CHURCH and SCOTT.—MANUAL OF VEGETABLE PHYSIOLOGY. By Professor A. H. CHURCH, and D. H. SCOTT, D.Sc., Lecturer in the Normal School of Science. Illustrated. Cr. 8vo. [*In preparation.*

COPE.—THE ORIGIN OF THE FITTEST. Essays on Evolution. By E. D. COPE, M.A., Ph.D. 8vo. 12s. 6d.

COUES.—HANDBOOK OF FIELD AND GENERAL ORNITHOLOGY. By Prof. ELLIOTT COUES, M.A. Illustrated. 8vo. 10s. net.

DARWIN.—MEMORIAL NOTICES OF CHARLES DARWIN, F.R.S., etc. By T. H. HUXLEY, F.R.S., G. J. ROMANES, F.R.S., ARCHIBALD GEIKIE, F.R.S., and W. THISELTON DYER, F.R.S. Reprinted from *Nature*. With a Portrait. Cr. 8vo. 2s. 6d.

EIMER.—ORGANIC EVOLUTION AS THE RESULT OF THE INHERITANCE OF ACQUIRED CHARACTERS ACCORDING TO THE LAWS OF ORGANIC GROWTH. By Dr. G. H. THEODOR EIMER. Translated by J. T. CUNNINGHAM, F.R.S.E., late Fellow of University College, Oxford. 8vo. 12s. 6d.

FEARNLEY.—A MANUAL OF ELEMENTARY PRACTICAL HISTOLOGY. By WILLIAM FEARNLEY. Illustrated. Cr. 8vo. 7s. 6d.

FLOWER and GADOW.—AN INTRODUCTION TO THE OSTEOLOGY OF THE MAMMALIA. By W. H. FLOWER, F.R.S., Director of the Natural History Departments of the British Museum. Illustrated. 3d Ed. Revised with the assistance of HANS GADOW, Ph.D., Lecturer on the Advanced Morphology of Vertebrates in the University of Cambridge. Cr. 8vo. 10s. 6d.

FOSTER.—Works by MICHAEL FOSTER, M.D., Professor of Physiology in the University of Cambridge.

*PRIMER OF PHYSIOLOGY. Illustrated. 18mo. 1s.

A TEXT-BOOK OF PHYSIOLOGY. Illustrated. 5th Ed., largely revised. 8vo. Part I., comprising Book I. Blood—The Tissues of Movement, The Vascular Mechanism. 10s. 6d. Part II., comprising Book II. The Tissues of Chemical Action, with their Respective Mechanisms—Nutrition. 10s. 6d. Part III. The Central Nervous System. 7s. 6d.

OSTER and BALFOUR.—THE ELEMENTS OF EMBRYOLOGY. By Prof. MICHAEL FOSTER, M.D., and the late F. M. BALFOUR, F.R.S., Professor of Animal Morphology in the University of Cambridge. 2d Ed., revised. Edited by A. SEDGWICK, M.A., Fellow and Assistant Lecturer of Trinity College, Cambridge, and W. HEAPE, M.A., late Demonstrator in the Morphological Laboratory of the University of Cambridge. Illustrated. Cr. 8vo. 10s. 6d.

FOSTER and LANGLEY.—A COURSE OF ELEMENTARY PRACTICAL PHYSIOLOGY AND HISTOLOGY. By Prof. MICHAEL FOSTER, M.D., and J. N. LANGLEY, F.R.S., Fellow of Trinity College, Cambridge. 6th Ed. Cr. 8vo. 7s. 6d.

GAMGEE.—A TEXT-BOOK OF THE PHYSIOLOGICAL CHEMISTRY OF THE ANIMAL BODY. Including an Account of the Chemical Changes occurring in Disease. By A. GAMGEE, M.D., F.R.S. Illustrated. 8vo. Vol. I. 18s.

GOODALE.—PHYSIOLOGICAL BOTANY. I. Outlines of the Histology of Phænogamous Plants. II. Vegetable Physiology. By GEORGE LINCOLN GOODALE, M.A., M.D., Professor of Botany in Harvard University. 8vo. 10s. 6d.

GRAY.—STRUCTURAL BOTANY, OR ORGANOGRAPHY ON THE BASIS OF MORPHOLOGY. To which are added the Principles of Taxonomy and Phytography, and a Glossary of Botanical Terms. By Prof. ASA GRAY, LL.D. 8vo. 10s. 6d.

THE SCIENTIFIC PAPERS OF ASA GRAY. Selected by C. SPRAGUE SARGENT. 2 vols. Vol. I. Reviews of Works on Botany and Related Subjects, 1834-1887. Vol. II. Essays, Biographical Sketches, 1841-1886. 8vo. 21s.

HAMILTON.—A SYSTEMATIC AND PRACTICAL TEXT-BOOK OF PATHOLOGY. By D. J. HAMILTON, F.R.S.E., Professor of Pathological Anatomy in the University of Aberdeen. Illustrated. 8vo. Vol. I. 25s.

HARTIG.—TEXT-BOOK OF THE DISEASES OF TREES. By Dr. ROBERT HARTIG. Translated by WM. SOMERVILLE, B.Sc., D.Œ., Lecturer on Forestry in the University of Edinburgh. Edited, with Introduction, by Prof. H. MARSHALL WARD. 8vo. [*In preparation.*

HOOKER.—Works by Sir JOSEPH HOOKER, F.R.S., &c.

*PRIMER OF BOTANY. Illustrated. 18mo. 1s.

THE STUDENT'S FLORA OF THE BRITISH ISLANDS. 3d Ed., revised. Gl. 8vo. 10s. 6d.

HOWES.—AN ATLAS OF PRACTICAL ELEMENTARY BIOLOGY. By G. B. HOWES, Assistant Professor of Zoology, Normal School of Science and Royal School of Mines. With a Preface by Prof. T. H. HUXLEY, F.R.S. 4to. 14s.

HUXLEY.—Works by Prof. T. H. HUXLEY, F.R.S.

*INTRODUCTORY PRIMER OF SCIENCE. 18mo. 1s.

*LESSONS IN ELEMENTARY PHYSIOLOGY. Illustrated. Fcap. 8vo. 4s. 6d.

*QUESTIONS ON HUXLEY'S PHYSIOLOGY. By T. ALCOCK, M.D. 18mo. 1s. 6d.

HUXLEY and MARTIN.—A COURSE OF PRACTICAL INSTRUCTION IN ELEMENTARY BIOLOGY. By Prof. T. H. HUXLEY, F.R.S., assisted by H. N. MARTIN, F.R.S., Professor of Biology in the Johns Hopkins University, U.S.A. New Ed., revised and extended by G. B. HOWES and D. H. SCOTT, Ph.D., Assistant Professors, Normal School of Science and Royal School of Mines. With a Preface by T. H. HUXLEY, F.R.S. Cr. 8vo. 10s. 6d.

KLEIN.—Works by E. KLEIN, F.R.S., Lecturer on General Anatomy and Physiology in the Medical School of St. Bartholomew's Hospital, Professor of Bacteriology at the College of State Medicine, London.

MICRO-ORGANISMS AND DISEASE. An Introduction into the Study of Specific Micro-Organisms. Illustrated. 3d Ed., revised. Cr. 8vo. 6s.

THE BACTERIA IN ASIATIC CHOLERA. Cr. 8vo. 5s.

LANG.—TEXT-BOOK OF COMPARATIVE ANATOMY. By Dr. ARNOLD LANG, Professor of Zoology in the University of Zurich. Translated by H. M. BERNARD, M.A., and M. BERNARD. Introduction by Prof. E. HAECKEL. 2 vols. Illustrated. 8vo. [*In the Press.*

LANKESTER.—Works by E. RAY LANKESTER, F.R.S., Linacre Professor of Human and Comparative Anatomy in the University of Oxford.

A TEXT-BOOK OF ZOOLOGY. 8vo. [*In preparation.*

THE ADVANCEMENT OF SCIENCE. Occasional Essays and Addresses. 8vo. 10s. 6d.

LUBBOCK.—Works by the Right Hon. Sir JOHN LUBBOCK, F.R.S., D.C.L.

THE ORIGIN AND METAMORPHOSES OF INSECTS. Illustrated. Cr. 8vo. 3s. 6d.

ON BRITISH WILD FLOWERS CONSIDERED IN RELATION TO INSECTS. Illustrated. Cr. 8vo. 4s. 6d.

FLOWERS, FRUITS, AND LEAVES. Illustrated. 2d Ed. Cr. 8vo. 4s. 6d.

SCIENTIFIC LECTURES. 2d Ed. 8vo. 8s. 6d.

FIFTY YEARS OF SCIENCE. Being the Address delivered at York to the British Association, August 1881. 5th Ed. Cr. 8vo. 2s. 6d.

MARTIN and MOALE.—ON THE DISSECTION OF VERTEBRATE ANIMALS. By Prof. H. N. MARTIN and W. A. MOALE. Cr. 8vo. [*In preparation.*

MIVART.—LESSONS IN ELEMENTARY ANATOMY. By ST. GEORGE MIVART, F.R.S., Lecturer on Comparative Anatomy at St. Mary's Hospital. Illustrated. Fcap. 8vo. 6s. 6d.

MÜLLER.—THE FERTILISATION OF FLOWERS. By HERMANN MÜLLER. Translated and Edited by D'ARCY W. THOMPSON, B.A., Professor of Biology in University College, Dundee. With a Preface by C. DARWIN, F.R.S. Illustrated. 8vo. 21s.

OLIVER.—Works by DANIEL OLIVER, F.R.S., late Professor of Botany in University College, London.

*LESSONS IN ELEMENTARY BOTANY. Illustrated. Fcap. 8vo. 4s. 6d.

FIRST BOOK OF INDIAN BOTANY. Illustrated. Ex. fcap. 8vo. 6s. 6d.

PARKER.—Works by T. JEFFERY PARKER, F.R.S., Professor of Biology in the University of Otago, New Zealand.

A COURSE OF INSTRUCTION IN ZOOTOMY (VERTEBRATA). Illustrated. Cr. 8vo. 8s. 6d.

LESSONS IN ELEMENTARY BIOLOGY. Illustrated. Cr. 8vo. [*In the Press.*

PARKER and BETTANY.—THE MORPHOLOGY OF THE SKULL. By Prof. W. K. PARKER, F.R.S., and G. T. BETTANY. Illustrated. Cr. 8vo. 10s. 6d.

ROMANES.—THE SCIENTIFIC EVIDENCES OF ORGANIC EVOLUTION. By GEORGE J. ROMANES, F.R.S., Zoological Secretary of the Linnean Society. Cr. 8vo. 2s. 6d.

SEDGWICK.—A SUPPLEMENT TO F. M. BALFOUR'S TREATISE ON EMBRYOLOGY. By ADAM SEDGWICK, F.R.S., Fellow and Lecturer of Trinity College, Cambridge. Illustrated. 8vo. [*In preparation.*

SHUFELDT.—THE MYOLOGY OF THE RAVEN (*Corvus corax sinuatus*). A Guide to the Study of the Muscular System in Birds. By R. W. SHUFELDT. Illustrated. 8vo. 13s. net.

SMITH.—DISEASES OF FIELD AND GARDEN CROPS, CHIEFLY SUCH AS ARE CAUSED BY FUNGI. By W. G. SMITH, F.L.S. Illustrated. Fcap. 8vo. 4s. 6d.

STEWART and CORRY.—A FLORA OF THE NORTH-EAST OF IRELAND. Including the Phanerogamia, the Cryptogamia Vascularia, and the Muscineæ. By S. A. STEWART, Curator of the Collections in the Belfast Museum, and the late T. H. CORRY, M.A., Lecturer on Botany in the University Medical and Science Schools, Cambridge. Cr. 8vo. 5s. 6d.

WALLACE.—DARWINISM: An Exposition of the Theory of Natural Selection, with some of its Applications. By ALFRED RUSSEL WALLACE, LL.D., F.R.S. 3d Ed. Cr. 8vo. 9s.

WARD.—TIMBER AND SOME OF ITS DISEASES. By H. MARSHALL WARD, F.R.S., Professor of Botany in the Royal Indian Engineering College, Cooper's Hill. Illustrated. Cr. 8vo. 6s.

WIEDERSHEIM.—ELEMENTS OF THE COMPARATIVE ANATOMY OF VERTEBRATES. By Prof. R. WIEDERSHEIM. Adapted by W. NEWTON PARKER, Professor of Biology in the University College of South Wales and Monmouthshire. With Additions. Illustrated. 8vo. 12s. 6d.

MEDICINE.

BLYTH.—A MANUAL OF PUBLIC HEALTH. By A. WYNTER BLYTH, M.R.C.S. 8vo. 17s. net.

BRUNTON.—Works by T. Lauder Brunton, M.D., F.R.S., Examiner in Materia Medica in the University of London, in the Victoria University, and in the Royal College of Physicians, London.

A TEXT-BOOK OF PHARMACOLOGY, THERAPEUTICS, AND MATERIA MEDICA. Adapted to the United States Pharmacopœia by F. H. Williams, M.D., Boston, Mass. 3d Ed. Adapted to the New British Pharmacopœia, 1885. 8vo. 21s.

TABLES OF MATERIA MEDICA: A Companion to the Materia Medica Museum. Illustrated. Cheaper Issue. 8vo. 5s.

ON THE CONNECTION BETWEEN CHEMICAL CONSTITUTION AND PHYSIOLOGICAL ACTION, BEING AN INTRODUCTION TO MODERN THERAPEUTICS. Croonian Lectures. 8vo. [*In the Press.*

GRIFFITHS.—LESSONS ON PRESCRIPTIONS AND THE ART OF PRESCRIBING. By W. Handsel Griffiths. Adapted to the Pharmacopœia, 1885. 18mo. 3s. 6d.

HAMILTON.—A TEXT-BOOK OF PATHOLOGY, SYSTEMATIC AND PRACTICAL. By D. J. Hamilton, F.R.S.E., Professor of Pathological Anatomy, University of Aberdeen. Illustrated. Vol. I. 8vo. 25s.

KLEIN.—Works by E. Klein, F.R.S., Lecturer on General Anatomy and Physiology in the Medical School of St. Bartholomew's Hospital, London.

MICRO-ORGANISMS AND DISEASE. An Introduction into the Study of Specific Micro-Organisms. Illustrated. 3d Ed., revised. Cr. 8vo. 6s.

THE BACTERIA IN ASIATIC CHOLERA. Cr. 8vo. 5s.

WHITE.—A TEXT-BOOK OF GENERAL THERAPEUTICS. By W. Hale White, M.D., Senior Assistant Physician to and Lecturer in Materia Medica at Guy's Hospital. Illustrated. Cr. 8vo. 8s. 6d.

*** Contents: General Medical Climatology—Classification of Climates—Treatment of Disease by Climate—Treatment by Compressed Air—Oertel's Method of Treatment for Chronic Cardiac Disease—Diet—Diet in Different Diseases—On the Drinking of Water—Baths and other external applications of Water—Cold Water as an Antipyretic—Artificial Baths—On Lavage, or washing out the Stomach—Massage—The Weir Mitchell Method—Venesection—Electricity, Physical and Physiological Facts of Importance in Medicine—Electrodiagnosis—Electrotherapeutics—Hypnotism, its Mode of Production, Phenomena, and Subdivisions—The Therapeutic Applications of Hypnotism—Metallo-Therapy—Treatment of Diseases of the Spinal Cord by Suspension—General Treatment of Acute Diseases, Convalescence and Insomnia—Index.

ZIEGLER—MACALISTER.—TEXT-BOOK OF PATHOLOGICAL ANATOMY AND PATHOGENESIS. By Prof. E. Ziegler. Translated and Edited by Donald Macalister, M.A., M.D., Fellow and Medical Lecturer of St. John's College, Cambridge. Illustrated. 8vo.

Part I.—GENERAL PATHOLOGICAL ANATOMY. 2d Ed. 12s. 6d.

Part II.—SPECIAL PATHOLOGICAL ANATOMY. Sections I.–VIII. 2d Ed. 12s. 6d. Sections IX.–XII. 12s. 6d.

HUMAN SCIENCES.

Mental and Moral Philosophy; Political Economy; Law and Politics; Anthropology; Education.

MENTAL AND MORAL PHILOSOPHY.

BALDWIN.—HANDBOOK OF PSYCHOLOGY: SENSES AND INTELLECT. By Prof. J. M. Baldwin, M.A., LL.D. 2d Ed., revised. 8vo. 12s. 6d.

BOOLE.—THE MATHEMATICAL ANALYSIS OF LOGIC. Being an Essay towards a Calculus of Deductive Reasoning. By George Boole. 8vo. 5s.

CALDERWOOD.—HANDBOOK OF MORAL PHILOSOPHY. By Rev. Henry Calderwood, LL.D., Professor of Moral Philosophy in the University of Edinburgh. 14th Ed., largely rewritten. Cr. 8vo. 6s.

CLIFFORD.—SEEING AND THINKING. By the late Prof. W. K. Clifford, F.R.S. With Diagrams. Cr. 8vo. 3s. 6d.

HÖFFDING.—OUTLINES OF PSYCHOLOGY. By Prof. H. Höffding. Translated by M. E. Lowndes. Cr. 8vo. 6s.

JAMES.—THE PRINCIPLES OF PSYCHOLOGY. By Wm. James, Professor of Psychology in Harvard University. 2 vols. 8vo. 25s. net.

JARDINE.—THE ELEMENTS OF THE PSYCHOLOGY OF COGNITION. By Rev. Robert Jardine, D.Sc. 3d Ed., revised. Cr. 8vo. 6s. 6d.

JEVONS.—Works by W. Stanley Jevons, F.R.S.

*PRIMER OF LOGIC. 18mo. 1s.

*ELEMENTARY LESSONS IN LOGIC, Deductive and Inductive, with Copious Questions and Examples, and a Vocabulary of Logical Terms. Fcap. 8vo. 3s. 6d.

THE PRINCIPLES OF SCIENCE. A Treatise on Logic and Scientific Method. New and revised Ed. Cr. 8vo. 12s. 6d.

STUDIES IN DEDUCTIVE LOGIC. 2d Ed. Cr. 8vo. 6s.

PURE LOGIC: AND OTHER MINOR WORKS. Edited by R. Adamson, M.A., LL.D., Professor of Logic at Owens College, Manchester, and Harriet A. Jevons. With a Preface by Prof. Adamson. 8vo. 10s. 6d.

KANT—MAX MÜLLER.—CRITIQUE OF PURE REASON. By Immanuel Kant. 2 vols. 8vo. 16s. each. Vol. I. HISTORICAL INTRODUCTION, by Ludwig Noiré; Vol. II. CRITIQUE OF PURE REASON, translated by F. Max Müller.

KANT—MAHAFFY and BERNARD.—KANT'S CRITICAL PHILOSOPHY FOR ENGLISH READERS. By J. P. Mahaffy, D.D., Professor of Ancient History in the University of Dublin, and John H. Bernard, B.D., Fellow of Trinity College, Dublin. A new and complete Edition in 2 vols. Cr. 8vo.

Vol. I. The Kritik of Pure Reason explained and defended. 7s. 6d.

Vol. II. The Prolegomena. Translated with Notes and Appendices. 6s.

KEYNES.—FORMAL LOGIC, Studies and Exercises in. Including a Generalisation of Logical Processes in their application to Complex Inferences. By John Neville Keynes, M.A. 2d Ed., revised and enlarged. Cr. 8vo. 10s. 6d.

McCOSH.—Works by James McCosh, D.D., President of Princeton College.

PSYCHOLOGY. Cr. 8vo.

I. THE COGNITIVE POWERS. 6s. 6d.

II. THE MOTIVE POWERS. 6s. 6d.

FIRST AND FUNDAMENTAL TRUTHS: being a Treatise on Metaphysics. Ex. cr. 8vo. 9s.

THE PREVAILING TYPES OF PHILOSOPHY. CAN THEY LOGICALLY REACH REALITY? 8vo. 3s. 6d.

MAURICE.—MORAL AND METAPHYSICAL PHILOSOPHY. By F. D. Maurice, M.A., late Professor of Moral Philosophy in the University of Cambridge. Vol. I.—Ancient Philosophy and the First to the Thirteenth Centuries. Vol. II.—Fourteenth Century and the French Revolution, with a glimpse into the Nineteenth Century. 4th Ed. 2 vols. 8vo. 16s.

***RAY.**—A TEXT-BOOK OF DEDUCTIVE LOGIC FOR THE USE OF STUDENTS. By P. K. Ray, D.Sc., Professor of Logic and Philosophy, Presidency College, Calcutta. 4th Ed. Globe 8vo. 4s. 6d.

SIDGWICK.—Works by Henry Sidgwick, LL.D., D.C.L., Knightbridge Professor of Moral Philosophy in the University of Cambridge.

THE METHODS OF ETHICS. 4th Ed. 8vo. 14s. A Supplement to the 2d Ed., containing all the important Additions and Alterations in the 3d Ed. 8vo. 6s.

OUTLINES OF THE HISTORY OF ETHICS, for English Readers. 2d Ed., revised. Cr. 8vo. 3s. 6d.

VENN.—Works by John Venn, F.R.S., Examiner in Moral Philosophy in the University of London.

THE LOGIC OF CHANCE. An Essay on the Foundations and Province of the Theory of Probability, with special Reference to its Logical Bearings and its Application to Moral and Social Science. 3d Ed., rewritten and greatly enlarged. Cr. 8vo. 10s. 6d.

VENN.—Works by JOHN VENN, F.R.S.—*continued.*

SYMBOLIC LOGIC. Cr. 8vo. 10s. 6d.

THE PRINCIPLES OF EMPIRICAL OR INDUCTIVE LOGIC. 8vo. 18s.

POLITICAL ECONOMY.

BÖHM-BAWERK.—CAPITAL AND INTEREST. Translated by WILLIAM SMART, M.A. 8vo. 14s.

THE POSITIVE THEORY OF CAPITAL. By the same Author and Translator. 8vo. [*In the Press.*

CAIRNES.—THE CHARACTER AND LOGICAL METHOD OF POLITICAL ECONOMY. By J. E. CAIRNES. Cr. 8vo. 6s.

SOME LEADING PRINCIPLES OF POLITICAL ECONOMY NEWLY EXPOUNDED. By the same. 8vo. 14s.

COSSA.—GUIDE TO THE STUDY OF POLITICAL ECONOMY. By Dr. L. COSSA. Translated. With a Preface by W. S. JEVONS, F.R.S. Cr. 8vo. 4s. 6d.

***FAWCETT.**—POLITICAL ECONOMY FOR BEGINNERS, WITH QUESTIONS. By Mrs. HENRY FAWCETT. 7th Ed. 18mo. 2s. 6d.

FAWCETT.—A MANUAL OF POLITICAL ECONOMY. By the Right Hon. HENRY FAWCETT, F.R.S. 7th Ed., revised. With a Chapter on "State Socialism and the Nationalisation of the Land," and an Index. Cr. 8vo. 12s. 6d.

AN EXPLANATORY DIGEST of the above. By C. A. WATERS, B.A. Cr. 8vo. 2s. 6d.

GILMAN.—PROFIT-SHARING BETWEEN EMPLOYER AND EMPLOYEE. A Study in the Evolution of the Wages System. By N. P. GILMAN. Cr. 8vo. 7s. 6d.

GUNTON.—WEALTH AND PROGRESS: A Critical Examination of the Wages Question and its Economic Relation to Social Reform. By GEORGE GUNTON. Cr. 8vo. 6s.

HOWELL.—THE CONFLICTS OF CAPITAL AND LABOUR HISTORICALLY AND ECONOMICALLY CONSIDERED. Being a History and Review of the Trade Unions of Great Britain, showing their Origin, Progress, Constitution, and Objects, in their varied Political, Social, Economical, and Industrial Aspects. By GEORGE HOWELL, M.P. 2d Ed., revised. Cr. 8vo. 7s. 6d.

JEVONS.—Works by W. STANLEY JEVONS, F.R.S.

*PRIMER OF POLITICAL ECONOMY. 18mo. 1s.

THE THEORY OF POLITICAL ECONOMY. 3d Ed., revised. 8vo. 10s. 6d.

KEYNES.—THE SCOPE AND METHOD OF POLITICAL ECONOMY. By J. N. KEYNES, M.A. 7s. net.

MARSHALL.—THE ECONOMICS OF INDUSTRY. By A. MARSHALL, M.A., Professor of Political Economy in the University of Cambridge, and MARY P. MARSHALL. Ex. fcap. 8vo. 2s. 6d.

MARSHALL.—PRINCIPLES OF ECONOMICS. By ALFRED MARSHALL, M.A. 2 vols. 8vo. Vol. I. 12s. 6d. net.

PALGRAVE.—A DICTIONARY OF POLITICAL ECONOMY. By various Writers. Edited by R. H. INGLIS PALGRAVE, F.R.S. 3s. 6d. each, net. [No. I. *May* 1891.

PANTALEONI.—MANUAL OF POLITICAL ECONOMY. By Prof. M. PANTALEONI. Translated by T. BOSTON BRUCE. [*In preparation.*

SIDGWICK.—THE PRINCIPLES OF POLITICAL ECONOMY. By HENRY SIDGWICK, LL.D., D.C.L., Knightbridge Professor of Moral Philosophy in the University of Cambridge. 2d Ed., revised. 8vo. 16s.

WALKER.—Works by FRANCIS A. WALKER, M.A.

FIRST LESSONS IN POLITICAL ECONOMY. Cr. 8vo. 5s.

A BRIEF TEXT-BOOK OF POLITICAL ECONOMY. Cr. 8vo. 6s. 6d.

POLITICAL ECONOMY. 2d Ed., revised and enlarged. 8vo. 12s. 6d.

THE WAGES QUESTION. 8vo. 14s.

***WICKSTEED.**—ALPHABET OF ECONOMIC SCIENCE. By PHILIP H. WICKSTEED, M.A. Part I. Elements of the Theory of Value or Worth. Gl. 8vo. 2s. 6d.

LAW AND POLITICS.

ADAMS and CUNNINGHAM.—THE SWISS CONFEDERATION. By Sir F. O. Adams and C. Cunningham. 8vo. 14s.

ANGLO-SAXON LAW, ESSAYS ON.—Contents: Anglo-Saxon Law Courts, Land and Family Law, and Legal Procedure. 8vo. 18s.

BALL.—THE STUDENT'S GUIDE TO THE BAR. By Walter W. R. Ball, M.A., Fellow and Assistant Tutor of Trinity College, Cambridge. 4th Ed., revised. Cr. 8vo. 2s. 6d.

BIGELOW.—HISTORY OF PROCEDURE IN ENGLAND FROM THE NORMAN CONQUEST. The Norman Period, 1066-1204. By Melville M. Bigelow, Ph.D., Harvard University. 8vo. 16s.

BOUTMY.—STUDIES IN CONSTITUTIONAL LAW. By Emile Boutmy. Translated by Mrs. Dicey, with Preface by Prof. A. V. Dicey. Crown 8vo. [*In the Press.*

THE ENGLISH CONSTITUTION. Translated by Mrs. Eaden, with Introduction by Sir F. Pollock, Bart. Cr. 8vo. 6s.

BRYCE.—THE AMERICAN COMMONWEALTH. By James Bryce, M.P., D.C.L., Regius Professor of Civil Law in the University of Oxford. Two Volumes. Ex. cr. 8vo. 25s. Part I. The National Government. Part II. The State Governments. Part III. The Party System. Part IV. Public Opinion. Part V. Illustrations and Reflections. Part VI. Social Institutions.

***BUCKLAND.**—OUR NATIONAL INSTITUTIONS. A Short Sketch for Schools. By Anna Buckland. With Glossary. 18mo. 1s.

CHERRY.—LECTURES ON THE GROWTH OF CRIMINAL LAW IN ANCIENT COMMUNITIES. By R. R. Cherry, LL.D., Reid Professor of Constitutional and Criminal Law in the University of Dublin. 8vo. 5s. net.

DICEY.—INTRODUCTION TO THE STUDY OF THE LAW OF THE CONSTITUTION. By A. V. Dicey, B.C.L., Vinerian Professor of English Law in the University of Oxford. 3d Ed. 8vo. 12s. 6d.

DILKE.—PROBLEMS OF GREATER BRITAIN. By the Right Hon. Sir Charles Wentworth Dilke. With Maps. 4th Ed. Ex. cr. 8vo. 12s. 6d.

DONISTHORPE.—INDIVIDUALISM: A System of Politics. By Wordsworth Donisthorpe. 8vo. 14s.

ENGLISH CITIZEN, THE.—A Series of Short Books on his Rights and Responsibilities. Edited by Henry Craik, LL.D. Cr. 8vo. 3s. 6d. each.

CENTRAL GOVERNMENT. By H. D. Traill, D.C.L.

THE ELECTORATE AND THE LEGISLATURE. By Spencer Walpole.

THE POOR LAW. By Rev. T. W. Fowle, M.A. New Ed. With Appendix.

THE NATIONAL BUDGET; THE NATIONAL DEBT; TAXES AND RATES. By A. J. Wilson.

THE STATE IN RELATION TO LABOUR. By W. Stanley Jevons, LL.D.

THE STATE AND THE CHURCH. By the Hon. Arthur Elliot.

FOREIGN RELATIONS. By Spencer Walpole.

THE STATE IN ITS RELATION TO TRADE. By Sir T. H. Farrer, Bart.

LOCAL GOVERNMENT. By M. D. Chalmers, M.A.

THE STATE IN ITS RELATION TO EDUCATION. By Henry Craik, LL.D.

THE LAND LAWS. By Sir F. Pollock, Bart., Professor of Jurisprudence in the University of Oxford.

COLONIES AND DEPENDENCIES. Part I. INDIA. By J. S. Cotton, M.A. II. THE COLONIES. By E. J. Payne, M.A.

JUSTICE AND POLICE. By F. W. Maitland.

THE PUNISHMENT AND PREVENTION OF CRIME. By Colonel Sir Edmund du Cane, K.C.B., Chairman of Commissioners of Prisons.

FISKE.—CIVIL GOVERNMENT IN THE UNITED STATES CONSIDERED WITH SOME REFERENCE TO ITS ORIGINS. By John Fiske, formerly Lecturer on Philosophy at Harvard University. Cr. 8vo. 6s. 6d.

HOLMES.—THE COMMON LAW. By O. W. Holmes, Jun. Demy 8vo. 12s.

MAITLAND.—PLEAS OF THE CROWN FOR THE COUNTY OF GLOUCESTER BEFORE THE ABBOT OF READING AND HIS FELLOW JUSTICES ITINERANT, IN THE FIFTH YEAR OF THE REIGN OF KING HENRY THE THIRD, AND THE YEAR OF GRACE 1221. By F. W. MAITLAND. 8vo. 7s. 6d

PATERSON.—Works by JAMES PATERSON, Barrister-at-Law.

COMMENTARIES ON THE LIBERTY OF THE SUBJECT, AND THE LAWS OF ENGLAND RELATING TO THE SECURITY OF THE PERSON. Cheaper Issue. Two Vols. Cr. 8vo. 21s.

THE LIBERTY OF THE PRESS, SPEECH, AND PUBLIC WORSHIP. Being Commentaries on the Liberty of the Subject and the Laws of England. Cr. 8vo. 12s.

PHILLIMORE.—PRIVATE LAW AMONG THE ROMANS. From the Pandects. By J. G. PHILLIMORE, Q.C. 8vo. 16s.

POLLOCK.—ESSAYS IN JURISPRUDENCE AND ETHICS. By Sir FREDERICK POLLOCK, Bart., Corpus Christi Professor of Jurisprudence in the University of Oxford. 8vo. 10s. 6d.

INTRODUCTION TO THE HISTORY OF THE SCIENCE OF POLITICS. By the same. Cr. 8vo. 2s. 6d.

RICHEY.—THE IRISH LAND LAWS. By ALEX. G. RICHEY, Q.C., Deputy Regius Professor of Feudal English Law in the University of Dublin. Cr. 8vo. 3s. 6d.

SIDGWICK.—THE ELEMENTS OF POLITICS. By HENRY SIDGWICK, LL.D. 8vo. *[In the Press.*

STEPHEN.—Works by Sir J. FITZJAMES STEPHEN, Q.C., K.C.S.I.

A DIGEST OF THE LAW OF EVIDENCE. 5th Ed., revised and enlarged. Cr. 8vo. 6s.

A DIGEST OF THE CRIMINAL LAW: CRIMES AND PUNISHMENTS. 4th Ed., revised. 8vo. 16s.

A DIGEST OF THE LAW OF CRIMINAL PROCEDURE IN INDICTABLE OFFENCES. By Sir J. F. STEPHEN, K.C.S.I., and H. STEPHEN, LL.M., of the Inner Temple, Barrister-at-Law. 8vo. 12s. 6d.

A HISTORY OF THE CRIMINAL LAW OF ENGLAND. Three Vols. 8vo. 48s.

GENERAL VIEW OF THE CRIMINAL LAW OF ENGLAND. 2d Ed. 8vo. 14s. The first edition of this work was published in 1863. The new edition is substantially a new work, intended as a text-book on the Criminal Law for University and other Students, adapted to the present day.

ANTHROPOLOGY.

DAWKINS.—EARLY MAN IN BRITAIN AND HIS PLACE IN THE TERTIARY PERIOD. By Prof. W. BOYD DAWKINS. Medium 8vo. 25s.

FRAZER.—THE GOLDEN BOUGH. A Study in Comparative Religion. By J. G. FRAZER, M.A., Fellow of Trinity College, Cambridge. 2 vols. 8vo. 28s.

M'LENNAN.—THE PATRIARCHAL THEORY. Based on the papers of the late JOHN F. M'LENNAN. Edited by DONALD M'LENNAN, M.A., Barrister-at-Law. 8vo. 14s.

STUDIES IN ANCIENT HISTORY. Comprising a Reprint of "Primitive Marriage." An inquiry into the origin of the form of capture in Marriage Ceremonies. 8vo. 16s.

TYLOR.—ANTHROPOLOGY. An Introduction to the Study of Man and Civilisation. By E. B. TYLOR, F.R.S. Illustrated. Cr. 8vo. 7s. 6d.

WESTERMARCK.—THE HISTORY OF HUMAN MARRIAGE. By Dr EDWARD WESTERMARCK. *[In preparation.*

EDUCATION.

ARNOLD.—REPORTS ON ELEMENTARY SCHOOLS. 1852-1882. By MATTHEW ARNOLD, D.C.L. Edited by the Right Hon. Sir FRANCIS SANDFORD, K.C.B. Cheaper Issue. Cr. 8vo. 3s. 6d.

HIGHER SCHOOLS AND UNIVERSITIES IN GERMANY. By the same. Crown 8vo. 6s.

BALL.—THE STUDENT'S GUIDE TO THE BAR. By Walter W. R. Ball, M.A., Fellow and Assistant Tutor of Trinity College, Cambridge. 4th Ed., revised. Cr. 8vo. 2s. 6d.

*BLAKISTON.—THE TEACHER. Hints on School Management. A Handbook for Managers, Teachers' Assistants, and Pupil Teachers. By J. R. Blakiston. Cr. 8vo. 2s. 6d. (Recommended by the London, Birmingham, and Leicester School Boards.)

CALDERWOOD.—ON TEACHING. By Prof. Henry Calderwood. New Ed. Ex. fcap. 8vo. 2s. 6d.

FEARON.—SCHOOL INSPECTION. By D. R. Fearon. 6th Ed. Cr. 8vo. 2s. 6d.

FITCH.—NOTES ON AMERICAN SCHOOLS AND TRAINING COLLEGES. Reprinted from the Report of the English Education Department for 1888-89, with permission of the Controller of H.M.'s Stationery Office. By J. G. Fitch, M.A. Gl. 8vo. 2s. 6d.

GEIKIE.—THE TEACHING OF GEOGRAPHY. A Practical Handbook for the use of Teachers. By Archibald Geikie, F.R.S., Director-General of the Geological Survey of the United Kingdom. Cr. 8vo. 2s.

GLADSTONE.—SPELLING REFORM FROM A NATIONAL POINT OF VIEW. By J. H. Gladstone. Cr. 8vo. 1s. 6d.

HERTEL.—OVERPRESSURE IN HIGH SCHOOLS IN DENMARK. By Dr. Hertel. Translated by C. G. Sörensen. With Introduction by Sir J. Crichton-Browne, F.R.S. Cr. 8vo. 3s. 6d.

TODHUNTER.—THE CONFLICT OF STUDIES. By Isaac Todhunter, F.R.S. 8vo. 10s. 6d.

TECHNICAL KNOWLEDGE.

(See also MECHANICS, LAW, and MEDICINE.)

Civil and Mechanical Engineering; Military and Naval Science; Agriculture; Domestic Economy; Book-Keeping.

CIVIL AND MECHANICAL ENGINEERING.

ALEXANDER and THOMSON.—ELEMENTARY APPLIED MECHANICS. By T. Alexander, Professor of Civil Engineering, Trinity College, Dublin, and A. W. Thomson, Professor at College of Science, Poona, India. Part II. Transverse Stress. Cr. 8vo. 10s. 6d.

CHALMERS.—GRAPHICAL DETERMINATION OF FORCES IN ENGINEERING STRUCTURES. By J. B. Chalmers, C.E. Illustrated. 8vo. 24s.

COTTERILL.—APPLIED MECHANICS: An Elementary General Introduction to the Theory of Structures and Machines. By J. H. Cotterill, F.R.S., Professor of Applied Mechanics in the Royal Naval College, Greenwich. 2d Ed. 8vo. 18s.

COTTERILL and SLADE.—LESSONS IN APPLIED MECHANICS. By Prof. J. H. Cotterill and J. H. Slade. Fcap. 8vo. 5s. 6d.

GRAHAM.—GEOMETRY OF POSITION. By R. H. Graham. Cr. 8vo. 7s. 6d.

KENNEDY.—THE MECHANICS OF MACHINERY. By A. B. W. Kennedy, F.R.S. Illustrated. Cr. 8vo. 12s. 6d.

WHITHAM.—STEAM-ENGINE DESIGN. For the Use of Mechanical Engineers, Students, and Draughtsmen. By J. M. Whitham, Professor of Engineering, Arkansas Industrial University. Illustrated. 8vo. 25s.

YOUNG.—SIMPLE PRACTICAL METHODS OF CALCULATING STRAINS ON GIRDERS, ARCHES, AND TRUSSES. With a Supplementary Essay on Economy in Suspension Bridges. By E. W. Young, C.E. With Diagrams. 8vo. 7s. 6d.

MILITARY AND NAVAL SCIENCE.

AITKEN.—THE GROWTH OF THE RECRUIT AND YOUNG SOLDIER. With a view to the selection of "Growing Lads" for the Army, and a Regulated System of Training for Recruits. By Sir W. Aitken, F.R.S., Professor of Pathology in the Army Medical School. Cr. 8vo. 8s. 6d.

ARMY PRELIMINARY EXAMINATION, 1882-1889, Specimens of Papers set at the. With Answers to the Mathematical Questions. Subjects: Arithmetic, Algebra, Euclid, Geometrical Drawing, Geography, French, English Dictation. Cr. 8vo. 8s. 6d.

MATTHEWS.—MANUAL OF LOGARITHMS. By G. F. MATTHEWS, B.A. 8vo. 5s. net.

MERCUR.—ELEMENTS OF THE ART OF WAR. Prepared for the use of Cadets of the United States Military Academy. By JAMES MERCUR, Professor of Civil Engineering at the United States Academy, West Point, New York. 2d Ed., revised and corrected. 8vo. 17s.

PALMER.—TEXT-BOOK OF PRACTICAL LOGARITHMS AND TRIGONOMETRY. By J. H. PALMER, Head Schoolmaster, R.N., H.M.S. *Cambridge*, Devonport. Gl. 8vo. 4s. 6d.

ROBINSON.—TREATISE ON MARINE SURVEYING. Prepared for the use of younger Naval Officers. With Questions for Examinations and Exercises principally from the Papers of the Royal Naval College. With the results. By Rev. JOHN L. ROBINSON, Chaplain and Instructor in the Royal Naval College, Greenwich. Illustrated. Cr. 8vo. 7s. 6d.

SANDHURST MATHEMATICAL PAPERS, for Admission into the Royal Military College, 1881-1889. Edited by E. J. BROOKSMITH, B.A., Instructor in Mathematics at the Royal Military Academy, Woolwich. Cr. 8vo. 3s. 6d.

SHORTLAND.—NAUTICAL SURVEYING. By the late Vice-Admiral SHORTLAND, LL.D. 8vo. 21s.

THOMSON.—POPULAR LECTURES AND ADDRESSES. By Sir WILLIAM THOMSON, LL.D., P.R.S. In 3 vols. Illustrated. Cr. 8vo. Vol. III. Papers on Navigation. [*In the Press.*

WILKINSON.—THE BRAIN OF AN ARMY. A Popular Account of the German General Staff. By SPENSER WILKINSON. Cr. 8vo. 2s. 6d.

WOLSELEY.—Works by General Viscount WOLSELEY, G.C.M.G.

THE SOLDIER'S POCKET-BOOK FOR FIELD SERVICE. 5th Ed., revised and enlarged. 16mo. Roan. 5s.

FIELD POCKET-BOOK FOR THE AUXILIARY FORCES. 16mo. 1s. 6d.

WOOLWICH MATHEMATICAL PAPERS, for Admission into the Royal Military Academy, Woolwich, 1880-1888 inclusive. Edited by E. J. BROOKSMITH, B.A., Instructor in Mathematics at the Royal Military Academy, Woolwich. Cr. 8vo. 6s.

AGRICULTURE.

FRANKLAND.—AGRICULTURAL CHEMICAL ANALYSIS, A Handbook of. By PERCY F. FRANKLAND, F.R.S., Professor of Chemistry, University College, Dundee. Founded upon *Leitfaden für die Agriculture Chemiche Analyse*, von Dr. F. KROCKER. Cr. 8vo. 7s. 6d.

HARTIG.—TEXT-BOOK OF THE DISEASES OF TREES. By Dr. ROBERT HARTIG. Translated by WM. SOMERVILLE, B.Sc., D.Œ., Lecturer on Forestry in the University of Edinburgh. Edited, with Introduction, by Prof. H. MARSHALL WARD. 8vo. [*In preparation.*

LASLETT.—TIMBER AND TIMBER TREES, NATIVE AND FOREIGN. By THOMAS LASLETT. Cr. 8vo. 8s. 6d.

SMITH.—DISEASES OF FIELD AND GARDEN CROPS, CHIEFLY SUCH AS ARE CAUSED BY FUNGI. By WORTHINGTON G. SMITH, F.L.S. Illustrated. Fcap. 8vo. 4s. 6d.

TANNER.—*ELEMENTARY LESSONS IN THE SCIENCE OF AGRICULTURAL PRACTICE. By HENRY TANNER, F.C.S., M.R.A.C., Examiner in the Principles of Agriculture under the Government Department of Science. Fcap. 8vo. 8s. 6d.

*FIRST PRINCIPLES OF AGRICULTURE. By the same. 18mo. 1s.

THE PRINCIPLES OF AGRICULTURE. By the same. A Series of Reading Books for use in Elementary Schools. Ex. fcap. 8vo.

*I. The Alphabet of the Principles of Agriculture. 6d.

*II. Further Steps in the Principles of Agriculture. 1s.

*III. Elementary School Readings on the Principles of Agriculture for the third stage. 1s.

WARD.—TIMBER AND SOME OF ITS DISEASES. By H. Marshall Ward, M.A., F.L.S., F.R.S., Fellow of Christ's College, Cambridge, Professor of Botany at the Royal Indian Engineering College, Cooper's Hill. With Illustrations. Cr. 8vo. 6s.

DOMESTIC ECONOMY.

***BARKER.**—FIRST LESSONS IN THE PRINCIPLES OF COOKING. By Lady Barker. 18mo. 1s.

***BERNERS.**—FIRST LESSONS ON HEALTH. By J. Berners. 18mo. 1s.

***COOKERY BOOK.**—THE MIDDLE CLASS COOKERY BOOK. Edited by the Manchester School of Domestic Cookery. Fcap. 8vo. 1s. 6d.

CRAVEN.—A GUIDE TO DISTRICT NURSES. By Mrs. Dacre Craven (*née* Florence Sarah Lees), Hon. Associate of the Order of St. John of Jerusalem, etc. Cr. 8vo. 2s. 6d.

FREDERICK.—HINTS TO HOUSEWIVES ON SEVERAL POINTS, PARTICULARLY ON THE PREPARATION OF ECONOMICAL AND TASTEFUL DISHES. By Mrs. Frederick. Cr. 8vo. 1s.

***GRAND'HOMME.**—CUTTING-OUT AND DRESSMAKING. From the French of Mdlle. E. Grand'homme. With Diagrams. 18mo. 1s.

JEX-BLAKE.—THE CARE OF INFANTS. A Manual for Mothers and Nurses. By Sophia Jex-Blake, M.D., Lecturer on Hygiene at the London School of Medicine for Women. 18mo. 1s.

RATHBONE.—THE HISTORY AND PROGRESS OF DISTRICT NURSING FROM ITS COMMENCEMENT IN THE YEAR 1859 TO THE PRESENT DATE, including the foundation by the Queen of the Queen Victoria Jubilee Institute for Nursing the Poor in their own Homes. By William Rathbone, M.P. Cr. 8vo. 2s. 6d.

***TEGETMEIER.**—HOUSEHOLD MANAGEMENT AND COOKERY. With an Appendix of Recipes used by the Teachers of the National School of Cookery. By W. B. Tegetmeier. Compiled at the request of the School Board for London. 18mo. 1s.

***WRIGHT.**—THE SCHOOL COOKERY-BOOK. Compiled and Edited by C. E. Guthrie Wright, Hon. Sec. to the Edinburgh School of Cookery. 18mo. 1s.

BOOK-KEEPING.

***THORNTON.**—FIRST LESSONS IN BOOK-KEEPING. By J. Thornton. Cr. 8vo. 2s. 6d. KEY. Oblong 4to. 10s. 6d.

***PRIMER OF BOOK-KEEPING.** By the same. 18mo. 1s. KEY. 8vo. 2s. 6d.

GEOGRAPHY.

(See also PHYSICAL GEOGRAPHY.)

BARTHOLOMEW.—*THE ELEMENTARY SCHOOL ATLAS. By John Bartholomew, F.R.G.S. 4to. 1s.

***MACMILLAN'S SCHOOL ATLAS, PHYSICAL AND POLITICAL.** Consisting of 80 Maps and complete Index. By the same. Prepared for the use of Senior Pupils. Royal 4to. 8s. 6d. Half-morocco. 10s. 6d.

THE LIBRARY REFERENCE ATLAS OF THE WORLD. By the same. A Complete Series of 84 Modern Maps. With Geographical Index to 100,000 places. Half-morocco. Gilt edges. Folio. £2:12:6 net. Also issued in parts, 5s. each net. Geographical Index, 7s. 6d. net. Part I., April 1891.

***CLARKE.**—CLASS-BOOK OF GEOGRAPHY. By C. B. Clarke, F.R.S. New Ed., revised 1889, with 18 Maps. Fcap. 8vo. 3s. Sewed, 2s. 6d.

GEIKIE.—Works by Archibald Geikie, F.R.S., Director-General of the Geological Survey of the United Kingdom.

***THE TEACHING OF GEOGRAPHY.** A Practical Handbook for the use of Teachers. Cr. 8vo. 2s.

***GEOGRAPHY OF THE BRITISH ISLES.** 18mo. 1s.

***GREEN.**—A SHORT GEOGRAPHY OF THE BRITISH ISLANDS. By John Richard Green and A. S. Green. With Maps. Fcap. 8vo. 3s. 6d.

***GROVE.**—A PRIMER OF GEOGRAPHY. By Sir George Grove, D.C.L. Illustrated. 18mo. 1s.

KIEPERT.—A MANUAL OF ANCIENT GEOGRAPHY. By Dr. H. Kiepert. Cr. 8vo. 5s.

MACMILLAN'S GEOGRAPHICAL SERIES.—Edited by Archibald Geikie, F.R.S., Director-General of the Geological Survey of the United Kingdom.

*THE TEACHING OF GEOGRAPHY. A Practical Handbook for the Use of Teachers. By Archibald Geikie, F.R.S. Cr. 8vo. 2s.

*MAPS AND MAP-DRAWING. By W. A. Elderton. 18mo. 1s.

*GEOGRAPHY OF THE BRITISH ISLES. By A. Geikie, F.R.S. 18mo. 1s.

*AN ELEMENTARY CLASS-BOOK OF GENERAL GEOGRAPHY. By H. R. Mill, D.Sc., Lecturer on Physiography and on Commercial Geography in the Heriot-Watt College, Edinburgh. Illustrated. Cr. 8vo. 3s. 6d.

*GEOGRAPHY OF EUROPE. By J. Sime, M.A. Illustrated. Gl. 8vo. 3s.

*ELEMENTARY GEOGRAPHY OF INDIA, BURMA, AND CEYLON. By H. F. Blanford, F.G.S. Gl. 8vo. 2s. 6d.

GEOGRAPHY OF NORTH AMERICA. By Prof. N. S. Shaler. [*In preparation.*

GEOGRAPHY OF THE BRITISH COLONIES. By G. M. Dawson and A. Sutherland. [*In preparation.*

*** Other volumes will be announced in due course.

STRACHEY.—LECTURES ON GEOGRAPHY. By General Richard Strachey, R.E. Cr. 8vo. 4s. 6d.

***TOZER.**—A PRIMER OF CLASSICAL GEOGRAPHY. By H. F. Tozer, M.A. 18mo. 1s.

HISTORY.

ARNOLD.—THE SECOND PUNIC WAR. Being Chapters from THE HISTORY OF ROME, by the late Thomas Arnold, D.D., Headmaster of Rugby. Edited, with Notes, by W. T. Arnold, M.A. With 8 Maps. Cr. 8vo. 5s.

ARNOLD.—THE ROMAN SYSTEM OF PROVINCIAL ADMINISTRATION TO THE ACCESSION OF CONSTANTINE THE GREAT. By W. T. Arnold, M.A. Cr. 8vo. 6s.

***BEESLY.**—STORIES FROM THE HISTORY OF ROME. By Mrs. Beesly. Fcap. 8vo. 2s. 6d.

BRYCE.—Works by James Bryce, M.P., D.C.L., Regius Professor of Civil Law in the University of Oxford.

THE HOLY ROMAN EMPIRE. 9th Ed. Cr. 8vo. 7s. 6d.

*** Also a *Library Edition*. Demy 8vo. 14s.

THE AMERICAN COMMONWEALTH. 2 vols. Ex. cr. 8vo. 25s. Part I. The National Government. Part II. The State Governments. Part III. The Party System. Part IV. Public Opinion. Part V. Illustrations and Reflections. Part VI. Social Institutions.

***BUCKLEY.**—A HISTORY OF ENGLAND FOR BEGINNERS. By Arabella B. Buckley. With Maps and Tables. Gl. 8vo. 3s.

BURY.—A HISTORY OF THE LATER ROMAN EMPIRE FROM ARCADIUS TO IRENE, A.D. 395-800. By John B. Bury, M.A., Fellow of Trinity College, Dublin. 2 vols. 8vo. 32s.

CASSEL.—MANUAL OF JEWISH HISTORY AND LITERATURE. By Dr. D. Cassel. Translated by Mrs. Henry Lucas. Fcap. 8vo. 2s. 6d.

ENGLISH STATESMEN, TWELVE. Cr. 8vo. 2s. 6d. each.

William the Conqueror. By Edward A. Freeman, D.C.L., LL.D.

Henry II. By Mrs. J. R. Green.

Edward I. By F. York Powell. [*In preparation.*

Henry VII. By James Gairdner.

CARDINAL WOLSEY. By Professor M. CREIGHTON.
ELIZABETH. By E. S. BEESLY. [*In preparation.*
OLIVER CROMWELL. By FREDERIC HARRISON.
WILLIAM III. By H. D. TRAILL.
WALPOLE. By JOHN MORLEY.
CHATHAM. By JOHN MORLEY. [*In preparation.*
PITT. By JOHN MORLEY. [*In preparation.*
PEEL. By J. R. THURSFIELD.

FISKE.—Works by JOHN FISKE, formerly Lecturer on Philosophy at Harvard University.

THE CRITICAL PERIOD IN AMERICAN HISTORY, 1783–1789. Ex. cr. 8vo. 10s. 6d.

THE BEGINNINGS OF NEW ENGLAND; or, The Puritan Theocracy in its Relations to Civil and Religious Liberty. Cr. 8vo. 7s. 6d.

FREEMAN.—Works by EDWARD A. FREEMAN, D.C.L., Regius Professor of Modern History in the University of Oxford, etc.

*OLD ENGLISH HISTORY. With Maps. Ex. fcap. 8vo. 6s.

A SCHOOL HISTORY OF ROME. Cr. 8vo. [*In preparation.*

METHODS OF HISTORICAL STUDY. 8vo. 10s. 6d.

THE CHIEF PERIODS OF EUROPEAN HISTORY. Six Lectures. With an Essay on Greek Cities under Roman Rule. 8vo. 10s. 6d.

HISTORICAL ESSAYS. First Series. 4th Ed. 8vo. 10s. 6d.

HISTORICAL ESSAYS. Second Series. 3d Ed., with additional Essays. 8vo. 10s. 6d.

HISTORICAL ESSAYS. Third Series. 8vo. 12s.

THE GROWTH OF THE ENGLISH CONSTITUTION FROM THE EARLIEST TIMES. 4th Ed. Cr. 8vo. 5s.

*GENERAL SKETCH OF EUROPEAN HISTORY. Enlarged, with Maps, etc. 18mo. 3s. 6d.

*PRIMER OF EUROPEAN HISTORY. 18mo. 1s. (*History Primers.*)

FRIEDMANN.—ANNE BOLEYN. A Chapter of English History, 1527–1536. By PAUL FRIEDMANN. 2 vols. 8vo. 28s.

FYFFE.—A SCHOOL HISTORY OF GREECE. By C. A. FYFFE, M.A., late Fellow of University College, Oxford. Cr. 8vo. [*In preparation.*

GIBBINS.—THE HISTORY OF COMMERCE IN EUROPE. By H. de B. GIBBINS, M.A. With Maps. 18mo. 2s. 6d.

GREEN.—Works by JOHN RICHARD GREEN, LL.D., late Honorary Fellow of Jesus College, Oxford.

*A SHORT HISTORY OF THE ENGLISH PEOPLE. New and Revised Ed. With Maps, Genealogical Tables, and Chronological Annals. Cr. 8vo. 8s. 6d. 151st Thousand.

*Also the same in Four Parts. With the corresponding portion of Mr. Tait's "Analysis." Crown 8vo. 3s. each. Part I. 607–1265. Part II. 1204–1553. Part III. 1540–1689. Part IV. 1660–1873.

HISTORY OF THE ENGLISH PEOPLE. In four vols. 8vo. 16s. each.

Vol. I.—Early England, 449–1071; Foreign Kings, 1071–1214; The Charter, 1214–1291; The Parliament, 1307–1461. With 8 Maps.

Vol. II.—The Monarchy, 1461–1540; The Reformation, 1540–1603.

Vol. III.—Puritan England, 1603–1660; The Revolution, 1660–1688. With four Maps.

Vol. IV.—The Revolution, 1688–1760; Modern England, 1760–1815. With Maps and Index.

THE MAKING OF ENGLAND. With Maps. 8vo. 16s.

THE CONQUEST OF ENGLAND. With Maps and Portrait. 8vo. 18s.

*ANALYSIS OF ENGLISH HISTORY, based on Green's "Short History of the English People." By C. W. A. TAIT, M.A., Assistant Master at Clifton College. Revised and Enlarged Ed. Crown 8vo. 4s. 6d.

*READINGS FROM ENGLISH HISTORY. Selected and Edited by JOHN RICHARD GREEN. Three Parts. Gl. 8vo. 1s. 6d. each. I. Hengist to Cressy. II. Cressy to Cromwell. III. Cromwell to Balaklava.

GUEST.—LECTURES ON THE HISTORY OF ENGLAND. By M. J. Guest. With Maps. Cr. 8vo. 6s.

***HISTORICAL COURSE FOR SCHOOLS.**—Edited by E. A. Freeman, D.C.L., Regius Professor of Modern History in the University of Oxford. 18mo.

GENERAL SKETCH OF EUROPEAN HISTORY. By E. A. Freeman, D.C.L. New Ed., revised and enlarged. With Chronological Table, Maps, and Index. 3s. 6d.

HISTORY OF ENGLAND. By Edith Thompson. New Ed., revised and enlarged. With Coloured Maps. 2s. 6d.

HISTORY OF SCOTLAND. By Margaret Macarthur. 2s.

HISTORY OF ITALY. By Rev. W. Hunt, M.A. New Ed. With Coloured Maps. 3s. 6d.

HISTORY OF GERMANY. By J. Sime, M.A. New Ed., revised. 3s.

HISTORY OF AMERICA. By John A. Doyle. With Maps. 4s. 6d.

HISTORY OF EUROPEAN COLONIES. By E. J. Payne, M.A. With Maps. 4s. 6d.

HISTORY OF FRANCE. By Charlotte M. Yonge. With Maps. 3s. 6d.

HISTORY OF GREECE. By Edward A. Freeman, D.C.L. [*In preparation.*

HISTORY OF ROME. By Edward A. Freeman, D.C.L. [*In preparation.*

***HISTORY PRIMERS.**—Edited by John Richard Green, LL.D. 18mo. 1s. each.

ROME. By Bishop Creighton. Maps.

GREECE. By C. A. Fyffe, M.A., late Fellow of University College, Oxford. Maps.

EUROPE. By E. A. Freeman, D.C.L. Maps.

FRANCE. By Charlotte M. Yonge.

INDIAN HISTORY: ASIATIC AND EUROPEAN. By J. Talboys Wheeler.

GREEK ANTIQUITIES. By Rev. J. P. Mahaffy, D.D. Illustrated.

CLASSICAL GEOGRAPHY. By H. F. Tozer, M.A.

GEOGRAPHY. By Sir G. Grove, D.C.L. Maps.

ROMAN ANTIQUITIES. By Prof. Wilkins, Litt.D. Illustrated.

ANALYSIS OF ENGLISH HISTORY. By Prof. T. F. Tout, M.A.

HOLE.—A GENEALOGICAL STEMMA OF THE KINGS OF ENGLAND AND FRANCE. By Rev. C. Hole. On Sheet. 1s.

JENNINGS.—CHRONOLOGICAL TABLES. A synchronistic arrangement of the events of Ancient History (with an Index). By Rev. Arthur C. Jennings. 8vo. 5s.

LABBERTON.—NEW HISTORICAL ATLAS AND GENERAL HISTORY. By R. H. Labberton. 4to. New Ed., revised and enlarged. 15s.

LETHBRIDGE.—A SHORT MANUAL OF THE HISTORY OF INDIA. With an Account of India as it is. The Soil, Climate, and Productions; the People, their Races, Religions, Public Works, and Industries; the Civil Services, and System of Administration. By Sir Roper Lethbridge, Fellow of the Calcutta University. With Maps. Cr. 8vo. 5s.

MAHAFFY.—GREEK LIFE AND THOUGHT FROM THE AGE OF ALEXANDER TO THE ROMAN CONQUEST. By Rev. J. P. Mahaffy, D.D., Fellow of Trinity College, Dublin. Cr. 8vo. 12s. 6d.

THE GREEK WORLD UNDER ROMAN SWAY. From Plutarch to Polybius. By the same Author. Cr. 8vo. 10s. 6d.

MARRIOTT.—THE MAKERS OF MODERN ITALY: Mazzini, Cavour, Garibaldi. Three Lectures. By J. A. R. Marriott, M.A., Lecturer in Modern History and Political Economy, Oxford. Cr. 8vo. 1s. 6d.

MICHELET.—A SUMMARY OF MODERN HISTORY. By M. Michelet. Translated by M. C. M. Simpson. Gl. 8vo. 4s. 6d.

NORGATE.—ENGLAND UNDER THE ANGEVIN KINGS. By Kate Norgate. With Maps and Plans. 2 vols. 8vo. 32s.

OTTÉ.—SCANDINAVIAN HISTORY. By E. C. Otté. With Maps. Gl. 8vo. 6s.

SEELEY.—Works by J. R. Seeley, M.A., Regius Professor of Modern History in the University of Cambridge.

THE EXPANSION OF ENGLAND. Crown 8vo. 4s. 6d.

OUR COLONIAL EXPANSION. Extracts from the above. Cr. 8vo. Sewed. 1s.

***TAIT.**—ANALYSIS OF ENGLISH HISTORY, based on Green's "Short History of the English People." By C. W. A. Tait, M.A., Assistant Master at Clifton. Revised and Enlarged Ed. Cr. 8vo. 4s. 6d.

WHEELER.—Works by J. Talboys Wheeler.

*A PRIMER OF INDIAN HISTORY. Asiatic and European. 18mo. 1s.

*COLLEGE HISTORY OF INDIA, ASIATIC AND EUROPEAN. With Maps. Cr. 8vo. 8s.; sewed, 2s. 6d.

A SHORT HISTORY OF INDIA AND OF THE FRONTIER STATES OF AFGHANISTAN, NEPAUL, AND BURMA. With Maps. Cr. 8vo. 12s.

YONGE.—Works by Charlotte M. Yonge.

CAMEOS FROM ENGLISH HISTORY. Ex. fcap. 8vo. 5s. each. (1) FROM ROLLO TO EDWARD II. (2) THE WARS IN FRANCE. (3) THE WARS OF THE ROSES. (4) REFORMATION TIMES. (5) ENGLAND AND SPAIN. (6) FORTY YEARS OF STUART RULE (1603–1643). (7) REBELLION AND RESTORATION (1642–1678).

EUROPEAN HISTORY. Narrated in a Series of Historical Selections from the Best Authorities. Edited and arranged by E. M. Sewell and C. M. Yonge. Cr. 8vo. First Series, 1003–1154. 6s. Second Series, 1088–1228. 6s.

THE VICTORIAN HALF CENTURY—A JUBILEE BOOK. With a New Portrait of the Queen. Cr. 8vo. Paper covers, 1s. Cloth, 1s. 6d.

ART.

***ANDERSON.**—LINEAR PERSPECTIVE AND MODEL DRAWING. A School and Art Class Manual, with Questions and Exercises for Examination, and Examples of Examination Papers. By Laurence Anderson. Illustrated. 8vo. 2s.

COLLIER.—A PRIMER OF ART. By the Hon. John Collier. Illustrated. 18mo. 1s.

COOK.—THE NATIONAL GALLERY, A POPULAR HANDBOOK TO. By Edward T. Cook, with a preface by John Ruskin, LL.D., and Selections from his Writings. 3d Ed. Cr. 8vo. Half-morocco, 14s.

*** Also an Edition on large paper, limited to 250 copies. 2 vols. 8vo.

DELAMOTTE.—A BEGINNER'S DRAWING BOOK. By P. H. Delamotte, F.S.A. Progressively arranged. New Ed., improved. Cr. 8vo. 3s. 6d.

ELLIS.—SKETCHING FROM NATURE. A Handbook for Students and Amateurs. By Tristram J. Ellis. Illustrated by H. Stacy Marks, R.A., and the Author. New Ed., revised and enlarged. Cr. 8vo. 3s. 6d.

GROVE.—A DICTIONARY OF MUSIC AND MUSICIANS. A.D. 1450–1889. Edited by Sir George Grove, D.C.L. In four vols. 8vo. Price 21s. each. Also in Parts.

Parts I.–XIV., Parts XIX.–XXII., 3s. 6d. each. Parts XV., XVI., 7s. Parts XVII., XVIII., 7s. Parts XXIII.–XXV. (Appendix), 9s.

A COMPLETE INDEX TO THE ABOVE. By Mrs. E. Wodehouse. 8vo. 7s. 6d.

HUNT.—TALKS ABOUT ART. By William Hunt. With a Letter from Sir J. E. Millais, Bart., R.A. Cr. 8vo. 3s. 6d.

MELDOLA.—THE CHEMISTRY OF PHOTOGRAPHY. By Raphael Meldola, F.R.S., Professor of Chemistry in the Technical College, Finsbury. Cr. 8vo. 6s.

TAYLOR.—A PRIMER OF PIANOFORTE-PLAYING. By Franklin Taylor. Edited by Sir George Grove. 18mo. 1s.

TAYLOR.—A SYSTEM OF SIGHT-SINGING FROM THE ESTABLISHED MUSICAL NOTATION; based on the Principle of Tonic Relation, and Illustrated by Extracts from the Works of the Great Masters. By Sedley Taylor. 8vo. 5s. net.

TYRWHITT.—OUR SKETCHING CLUB. Letters and Studies on Landscape Art. By Rev. R. St. John Tyrwhitt. With an authorised Reproduction of the Lessons and Woodcuts in Prof. Ruskin's "Elements of Drawing." 4th Ed. Cr. 8vo. 7s. 6d.

DIVINITY.

ABBOTT.—BIBLE LESSONS. By Rev. Edwin A. Abbott, D.D. Cr. 8vo. 4s. 6d.

ABBOTT–RUSHBROOKE.—THE COMMON TRADITION OF THE SYNOPTIC GOSPELS, in the Text of the Revised Version. By Rev. Edwin A. Abbott, D.D., and W. G. Rushbrooke, M.L. Cr. 8vo. 8s. 6d.

ARNOLD.—Works by Matthew Arnold.

A BIBLE-READING FOR SCHOOLS,—THE GREAT PROPHECY OF ISRAEL'S RESTORATION (Isaiah, Chapters xl.-lxvi.) Arranged and Edited for Young Learners. 18mo. 1s.

ISAIAH XL.-LXVI. With the Shorter Prophecies allied to it. Arranged and Edited, with Notes. Cr. 8vo. 5s.

ISAIAH OF JERUSALEM, IN THE AUTHORISED ENGLISH VERSION. With Introduction, Corrections and Notes. Cr. 8vo. 4s. 6d.

BENHAM.—A COMPANION TO THE LECTIONARY. Being a Commentary on the Proper Lessons for Sundays and Holy Days. By Rev. W. Benham, B.D. Cr. 8vo. 4s. 6d.

CASSEL.—MANUAL OF JEWISH HISTORY AND LITERATURE; preceded by a BRIEF SUMMARY OF BIBLE HISTORY. By Dr. D. Cassel. Translated by Mrs. H. Lucas. Fcap. 8vo. 2s. 6d.

CHURCH.—STORIES FROM THE BIBLE. By Rev. A J. Church, M.A. Illustrated. Cr. 8vo. 5s.

***CROSS.**—BIBLE READINGS SELECTED FROM THE PENTATEUCH AND THE BOOK OF JOSHUA. By Rev. John A. Cross. 2d Ed., enlarged, with Notes. Gl. 8vo. 2s. 6d.

DRUMMOND.—INTRODUCTION TO THE STUDY OF THEOLOGY. By James Drummond, LL.D., Professor of Theology in Manchester New College, London. Cr. 8vo. 5s.

FARRAR.—Works by the Venerable Archdeacon F. W. Farrar, D.D., F.R.S., Archdeacon and Canon of Westminster.

THE HISTORY OF INTERPRETATION. Being the Bampton Lectures, 1885. 8vo. 16s.

THE MESSAGES OF THE BOOKS. Being Discourses and Notes on the Books of the New Testament. 8vo. 14s.

***GASKOIN.**—THE CHILDREN'S TREASURY OF BIBLE STORIES. By Mrs. Herman Gaskoin. Edited with Preface by Rev. G. F. Maclear, D.D. 18mo. 1s. each. Part I.—Old Testament History. Part II.—New Testament. Part III.—The Apostles: St. James the Great, St. Paul, and St. John the Divine.

GOLDEN TREASURY PSALTER.—Students' Edition. Being an Edition of "The Psalms chronologically arranged, by Four Friends," with briefer Notes. 18mo. 3s. 6d.

GREEK TESTAMENT.—Edited, with Introduction and Appendices, by Bishop Westcott and Dr. F. J. A. Hort. Two Vols. Cr. 8vo. 10s. 6d. each. Vol. I. The Text. Vol. II. Introduction and Appendix.

SCHOOL EDITION OF TEXT. 12mo. Cloth, 4s. 6d.; Roan, red edges, 5s. 6d. 18mo. Morocco, gilt edges, 6s. 6d.

***GREEK TESTAMENT, SCHOOL READINGS IN THE.** Being the outline of the life of our Lord, as given by St. Mark, with additions from the Text of the other Evangelists. Arranged and Edited, with Notes and Vocabulary, by Rev. A. Calvert, M.A. Fcap. 8vo. 2s. 6d.

*THE GOSPEL ACCORDING TO ST. MATTHEW. Being the Greek Text as revised by Bishop WESTCOTT and Dr. HORT. With Introduction and Notes by Rev. A. SLOMAN, M.A., Headmaster of Birkenhead School. Fcap. 8vo. 2s. 6d.

THE GOSPEL ACCORDING TO ST. MARK. Being the Greek Text as revised by Bishop WESTCOTT and Dr. HORT. With Introduction and Notes by Rev. J. O. F. MURRAY, M.A., Lecturer at Emmanuel College, Cambridge. Fcap. 8vo. [*In preparation.*

*THE GOSPEL ACCORDING TO ST. LUKE. Being the Greek Text as revised by Bishop WESTCOTT and Dr. HORT. With Introduction and Notes by Rev. JOHN BOND, M.A. Fcap. 8vo. 2s. 6d.

*THE ACTS OF THE APOSTLES. Being the Greek Text as revised by Bishop WESTCOTT and Dr. HORT. With Explanatory Notes by T. E. PAGE, M.A., Assistant Master at the Charterhouse. Fcap. 8vo. 3s. 6d.

GWATKIN.—CHURCH HISTORY TO THE BEGINNING OF THE MIDDLE AGES. By H. M. GWATKIN, M.A. 8vo. [*In preparation.*

HARDWICK.—Works by Archdeacon HARDWICK.

A HISTORY OF THE CHRISTIAN CHURCH. Middle Age. From Gregory the Great to the Excommunication of Luther. Edited by W. STUBBS, D.D., Bishop of Oxford. With 4 Maps. Cr. 8vo. 10s. 6d.

A HISTORY OF THE CHRISTIAN CHURCH DURING THE REFORMATION. 9th Ed. Edited by Bishop STUBBS. Cr. 8vo. 10s. 6d.

HOOLE.—THE CLASSICAL ELEMENT IN THE NEW TESTAMENT. Considered as a proof of its Genuineness, with an Appendix on the Oldest Authorities used in the Formation of the Canon. By CHARLES H. HOOLE, M.A., Student of Christ Church, Oxford. 8vo. 10s. 6d.

JENNINGS and LOWE.—THE PSALMS, WITH INTRODUCTIONS AND CRITICAL NOTES. By A. C. JENNINGS, M.A.; assisted in parts by W. H. LOWE, M.A. In 2 vols. 2d Ed., revised. Cr. 8vo. 10s. 6d. each.

KIRKPATRICK.—THE MINOR PROPHETS. Warburtonian Lectures. By Rev. Prof. KIRKPATRICK. [*In preparation.*

KUENEN.—PENTATEUCH AND BOOK OF JOSHUA: An Historico-Critical Inquiry into the Origin and Composition of the Hexateuch. By A. KUENEN. Translated by P. H. WICKSTEED, M.A. 8vo. 14s.

LIGHTFOOT.—Works by the Right Rev. J. B. LIGHTFOOT, D.D., late Bishop of Durham.

ST. PAUL'S EPISTLE TO THE GALATIANS. A Revised Text, with Introduction, Notes, and Dissertations. 10th Ed., revised. 8vo. 12s.

ST. PAUL'S EPISTLE TO THE PHILIPPIANS. A Revised Text, with Introduction, Notes, and Dissertations. 9th Ed., revised. 8vo. 12s.

ST. PAUL'S EPISTLES TO THE COLOSSIANS AND TO PHILEMON. A Revised Text, with Introductions, Notes, and Dissertations. 8th Ed., revised. 8vo. 12s.

THE APOSTOLIC FATHERS. Part I. ST. CLEMENT OF ROME. A Revised Text, with Introductions, Notes, Dissertations, and Translations. 2 vols. 8vo. 32s.

THE APOSTOLIC FATHERS. Part II. ST. IGNATIUS—ST. POLYCARP. Revised Texts, with Introductions, Notes, Dissertations, and Translations. 2d Ed. 3 vols. 8vo. 48s.

THE APOSTOLIC FATHERS. Abridged Edition. With short Introductions, Greek Text, and English Translation. 8vo. [*In the Press.*

ESSAYS ON THE WORK ENTITLED "SUPERNATURAL RELIGION." (Reprinted from the *Contemporary Review*.) 8vo. 10s. 6d.

MACLEAR.—Works by the Rev. G. F. MACLEAR, D.D., Warden of St. Augustine's College, Canterbury.

ELEMENTARY THEOLOGICAL CLASS-BOOKS.

*A SHILLING BOOK OF OLD TESTAMENT HISTORY. With Map. 18mo.

*A SHILLING BOOK OF NEW TESTAMENT HISTORY. With Map. 18mo.

These works have been carefully abridged from the Author's large manuals.

*A CLASS-BOOK OF OLD TESTAMENT HISTORY. Maps. 18mo. 4s. 6d.

MACLEAR.—Works by Rev. G. F. MACLEAR, D.D.—*continued.*

*A CLASS-BOOK OF NEW TESTAMENT HISTORY, including the Connection of the Old and New Testaments. With maps. 18mo. 5s. 6d.

AN INTRODUCTION TO THE THIRTY-NINE ARTICLES. [*In the Press.*

*AN INTRODUCTION TO THE CREEDS. 18mo. 2s. 6d.

*A CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND. 18mo. 1s. 6d.

*A FIRST CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND. With Scripture Proofs. 18mo. 6d.

*A MANUAL OF INSTRUCTION FOR CONFIRMATION AND FIRST COMMUNION. WITH PRAYERS AND DEVOTIONS. 32mo. 2s.

MAURICE.—THE LORD'S PRAYER, THE CREED, AND THE COMMANDMENTS. To which is added the Order of the Scriptures. By Rev. F. D. MAURICE, M.A. 18mo. 1s.

THE PENTATEUCH AND BOOK OF JOSHUA: An Historico-Critical Inquiry into the Origin and Composition of the Hexateuch. By A. KUENEN, Professor of Theology at Leiden. Translated by P. H. WICKSTEED, M.A. 8vo. 14s.

PROCTER.—A HISTORY OF THE BOOK OF COMMON PRAYER, with a Rationale of its Offices. By Rev. F. PROCTER. 18th Ed. Cr. 8vo. 10s. 6d.

***PROCTER and MACLEAR.**—AN ELEMENTARY INTRODUCTION TO THE BOOK OF COMMON PRAYER. Rearranged and supplemented by an Explanation of the Morning and Evening Prayer and the Litany. By Rev. F. PROCTER and Rev. Dr. MACLEAR. New Edition, containing the Communion Service and the Confirmation and Baptismal Offices. 18mo. 2s. 6d.

THE PSALMS, CHRONOLOGICALLY ARRANGED. By Four Friends. New Ed. Cr. 8vo. 5s. net.

THE PSALMS, WITH INTRODUCTIONS AND CRITICAL NOTES. By A. C. JENNINGS, M.A., Jesus College, Cambridge; assisted in parts by W. H. LOWE, M.A., Hebrew Lecturer at Christ's College, Cambridge. In 2 vols. 2d Ed., revised. Cr. 8vo. 10s. 6d. each.

RYLE.—AN INTRODUCTION TO THE CANON OF THE OLD TESTAMENT. By Rev. H. E. RYLE, Hulsean Professor of Divinity in the University of Cambridge. Cr. 8vo. [*In preparation.*

SIMPSON.—AN EPITOME OF THE HISTORY OF THE CHRISTIAN CHURCH DURING THE FIRST THREE CENTURIES, AND OF THE REFORMATION IN ENGLAND. By Rev. WILLIAM SIMPSON, M.A. 7th Ed. Fcap. 8vo. 3s. 6d.

ST. JAMES' EPISTLE.—The Greek Text, with Introduction and Notes. By Rev. JOSEPH MAYOR, M.A., Professor of Moral Philosophy in King's College, London. 8vo. [*In the Press.*

ST. JOHN'S EPISTLES.—The Greek Text, with Notes and Essays. By Right Rev. B. F. WESTCOTT, D.D., Bishop of Durham. 2d Ed., revised. 8vo. 12s. 6d.

ST. PAUL'S EPISTLES.—THE EPISTLE TO THE ROMANS. Edited by the Very Rev. C. J. VAUGHAN, D.D., Dean of Llandaff. 5th Ed. Cr. 8vo. 7s. 6d.

THE TWO EPISTLES TO THE CORINTHIANS, A COMMENTARY ON. By the late Rev. W. KAY, D.D., Rector of Great Leghs, Essex. 8vo. 9s.

THE EPISTLE TO THE GALATIANS. Edited by the Right Rev. J. B. LIGHTFOOT, D.D. 10th Ed. 8vo. 12s.

THE EPISTLE TO THE PHILIPPIANS. By the Same Editor. 9th Ed. 8vo. 12s.

THE EPISTLE TO THE PHILIPPIANS, with Translation, Paraphrase, and Notes for English Readers. By the Very Rev. C. J. VAUGHAN, D.D. Cr. 8vo. 5s.

THE EPISTLE TO THE COLOSSIANS AND TO PHILEMON. By the Right Rev. J. B. LIGHTFOOT, D.D. 8th Ed. 8vo. 12s.

THE EPISTLES TO THE EPHESIANS, THE COLOSSIANS, AND PHILEMON; with Introductions and Notes, and an Essay on the Traces of Foreign Elements in the Theology of these Epistles. By Rev. J. LLEWELYN DAVIES, M.A. 8vo. 7s. 6d.

THE EPISTLE TO THE THESSALONIANS, COMMENTARY ON THE GREEK TEXT. By JOHN EADIE, D.D. Edited by Rev. W. YOUNG, M.A., with Preface by Prof. CAIRNS. 8vo. 12s.

THE EPISTLE TO THE HEBREWS.—In Greek and English. With Critical and Explanatory Notes. Edited by Rev. F. RENDALL, M.A. Cr. 8vo. 6s.

THE ENGLISH TEXT, WITH COMMENTARY. By the same Editor. Cr. 8vo. 7s. 6d.

THE GREEK TEXT. With Notes by C. J. VAUGHAN, D.D., Dean of Llandaff. Cr. 8vo. 7s. 6d.

THE GREEK TEXT. With Notes and Essays by the Right Rev. Bishop WESTCOTT, D.D. 8vo. 14s.

VAUGHAN.—THE CHURCH OF THE FIRST DAYS. Comprising the Church of Jerusalem, the Church of the Gentiles, the Church of the World. By C. J. VAUGHAN, D.D., Dean of Llandaff. New Ed. Cr. 8vo. 10s. 6d.

WESTCOTT.—Works by the Right Rev. BROOKE FOSS WESTCOTT, D.D., Bishop of Durham.

A GENERAL SURVEY OF THE HISTORY OF THE CANON OF THE NEW TESTAMENT DURING THE FIRST FOUR CENTURIES. 6th Ed. With Preface on "Supernatural Religion." Cr. 8vo. 10s. 6d.

INTRODUCTION TO THE STUDY OF THE FOUR GOSPELS. 7th Ed. Cr. 8vo. 10s. 6d.

THE BIBLE IN THE CHURCH. A Popular Account of the Collection and Reception of the Holy Scriptures in the Christian Churches. 18mo. 4s. 6d.

THE EPISTLES OF ST. JOHN. The Greek Text, with Notes and Essays. 2d Ed., revised. 8vo. 12s. 6d.

THE EPISTLE TO THE HEBREWS. The Greek Text, with Notes and Essays. 8vo. 14s.

SOME THOUGHTS FROM THE ORDINAL. Cr. 8vo. 1s. 6d.

WESTCOTT and HORT.—THE NEW TESTAMENT IN THE ORIGINAL GREEK. The Text, revised by the Right Rev. Bishop WESTCOTT and Dr. F. J. A. HORT. 2 vols. Cr. 8vo. 10s. 6d. each. Vol. I. Text. Vol. II. Introduction and Appendix.

SCHOOL EDITION OF TEXT. 12mo. 4s. 6d.; Roan, red edges, 5s. 6d. 18mo. Morocco, gilt edges, 6s. 6d.

WILSON.—THE BIBLE STUDENT'S GUIDE to the more correct understanding of the English Translation of the Old Testament, by reference to the original Hebrew. By WILLIAM WILSON, D.D., Canon of Winchester. 2d Ed., carefully revised. 4to. 25s.

WRIGHT.—THE COMPOSITION OF THE FOUR GOSPELS. A Critical Enquiry. By Rev. ARTHUR WRIGHT, M.A., Fellow and Tutor of Queen's College, Cambridge. Cr. 8vo. 5s.

WRIGHT.—THE BIBLE WORD-BOOK: A Glossary of Archaic Words and Phrases in the Authorised Version of the Bible and the Book of Common Prayer. By W. ALDIS WRIGHT, M.A., Vice-Master of Trinity College, Cambridge. 2d Ed., revised and enlarged. Cr. 8vo. 7s. 6d.

***YONGE.**—SCRIPTURE READINGS FOR SCHOOLS AND FAMILIES. By CHARLOTTE M. YONGE. In Five Vols. Ex. fcap. 8vo. 1s. 6d. each. With Comments. 3s. 6d. each.

FIRST SERIES.—GENESIS TO DEUTERONOMY. SECOND SERIES.—FROM JOSHUA TO SOLOMON. THIRD SERIES.—THE KINGS AND THE PROPHETS. FOURTH SERIES.—THE GOSPEL TIMES. FIFTH SERIES.—APOSTOLIC TIMES.

ZECHARIAH—THE HEBREW STUDENT'S COMMENTARY ON ZECHARIAH, HEBREW AND LXX. With Excursus on Syllable-dividing, Metheg, Initial Dagesh, and Siman Rapheh. By W. H. LOWE, M.A., Hebrew Lecturer at Christ's College, Cambridge. 8vo. 10s. 6d.

Printed by R. & R. CLARK, *Edinburgh.*

The *Times* says: "Both artists and writers have done their best to give it a high place in English periodical literature, and the result is *a magazine which has no rival in England.*"

The *Guardian* describes it as: "Full of good matter both in the way of writing and drawing. . . . A capital magazine for all tables and all times."

The English Illustrated Magazine, 1884.

A Handsome Volume, of 792 pages, with 428 Illustrations, price 7s. 6d.

The Volume contains a Complete Series of Descriptive Sketches by the Author of "John Halifax, Gentleman," with Illustrations by C. NAPIER HEMY; a Complete Historical Novel by CHARLOTTE M. YONGE; and numerous Short Stories and Essays on Popular Subjects by well-known Writers.

The English Illustrated Magazine, 1885.

A Handsome Volume, of 840 pages, with nearly 500 Illustrations, price 8s.

The Volume contains a Complete Novel, "A FAMILY AFFAIR," by HUGH CONWAY, Author of "Called Back," and also Complete Stories and Essays by BRET HARTE, the Author of "John Herring," ARCHIBALD FORBES, J. HENRY SHORTHOUSE, HENRY IRVING, Mrs. OLIPHANT, and others, besides numerous interesting miscellaneous Articles by the First Authors of the Day.

The English Illustrated Magazine, 1886.

A Handsome Volume, of 832 pages, containing nearly 500 Illustrations, price 8s.

Among the Chief Contributors to the Volume are the following:—

THE AUTHOR OF "John Halifax, Gentleman."
D. CHRISTIE MURRAY.
MARGARET VELEY.
W. E. NORRIS.
GRANT ALLEN.
CLEMENTINA BLACK.
Mrs. MOLESWORTH.
WILKIE COLLINS.
Mrs. OLIPHANT.
KATHARINE S. MACQUOID.

Rev. CANON AINGER, ALFRED T. WATSON, H. SUTHERLAND EDWARDS, Dr. MORELL MACKENZIE, A. C. SWINBURNE, HENRY W. LUCY, EDMUND GOSSE, The Right Rev. The BISHOP OF RIPON, RANDOLPH CALDECOTT, Miss L. ALMA-TADEMA, and "Days with Sir ROGER DE COVERLEY," with pictures by HUGH THOMSON.

MACMILLAN AND CO., LONDON.

The English Illustrated Magazine, 1887.

A Handsome Volume, of 832 pages, and containing nearly 500 Illustrations, price 8s.

The Volume contains the following Complete Stories and Serials:—

Marzio's Crucifix. By F. Marion Crawford. **A Secret Inheritance.** By B. L. Farjeon. **Jacquetta.** By the Author of "John Herring." **Gerald.** By Stanley J. Weyman. **An Unknown Country.** By the Author of "John Halifax, Gentleman." With Illustrations by F. Noel Paton. **A Siege Baby.** By J. S. Winter. **Miss Falkland.** By Clementina Black.

The English Illustrated Magazine, 1888.

A Handsome Volume, of 832 pp., with nearly 500 Illustrations, price 8s.

Among the chief Contents of the Volume are the following Complete Stories and Serials:—

Coaching Days and Coaching Ways. By W. O. Tristram. With Illustrations by H. Railton and Hugh Thomson.

The Story of Jael. By the Author of "Mehalah."

Lil: A Liverpool Child. By Agnes C. Maitland.

The Patagonia. By Henry James.

Family Portraits. By S. J. Weyman.

The Mediation of Ralph Hardelot. By Prof. W. Minto.

That Girl in Black. By Mrs. Molesworth.

Glimpses of Old English Homes. By Elizabeth Balch.

Pagodas, Aurioles, and Umbrellas. By C. F. Gordon Cumming.

The Magic Fan. By John Strange Winter.

The English Illustrated Magazine, 1889.

A Handsome Volume, of 900 pp., with nearly 500 Illustrations; price 8s.

Among the chief Contents of the Volume are the following Complete Stories and Serials:—

Sant' Ilario. By F. Marion Crawford.

The House of the Wolf. By Stanley J. Weyman.

Glimpses of Old English Homes. By Elizabeth Balch.

One Night—The Better Man. By Arthur Paterson.

How the "Crayture" got on the Strength. And other Sketches. By Archibald Forbes.

La Belle Americaine. By W. E. Norris.

Success. By Katharine S. Macquoid.

Jenny Harlowe. By W. Clark Russell.

The English Illustrated Magazine, 1890.

A Handsome Volume, of 900 pp., with nearly 550 Illustrations, price 8s.

Among the chief Contents of the Volume are the following Complete Stories and Serials:—

The Ring of Amasis. By the Earl of Lytton. **The Glittering Plain: or, the Land of Living Men.** By William Morris. **The Old Brown Mare.** By W. E. Norris. **My Journey to Texas.** By Arthur Paterson. **A Glimpse of Highclere Castle—A Glimpse of Osterley Park.** By Elizabeth Balch. **For the Cause.** By Stanley J. Weyman. **Morised.** By the Marchioness of Carmarthen. **Overland from India.** By Sir Donald Mackenzie Wallace, K.C.I.E. **The Doll's House and After.** By Walter Besant. **La Mulette, Anno 1814.** By W. Clark Russell.

MACMILLAN AND CO., LONDON.

IV.60.5.91.